



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2020 – 02:32 PM BST

PDB ID : 5L2X
Title : Crystal structure of human PrimPol ternary complex
Authors : Rechkoblit, O.; Gupta, Y.K.; Malik, R.; Rajashankar, K.R.; Johnson, R.E.;
Prakash, L.; Prakash, S.; Aggarwal, A.K.
Deposited on : 2016-08-02
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	<div> <div>8%</div> <div>64%</div> <div>13%</div> <div>23%</div> </div>
1	B	353	<div> <div>5%</div> <div>59%</div> <div>10%</div> <div>29%</div> </div>
2	C	17	<div> <div>29%</div> <div>24%</div> <div>41%</div> <div>6%</div> <div>29%</div> </div>
2	G	17	<div> <div>12%</div> <div>35%</div> <div>6%</div> <div>6%</div> <div>53%</div> </div>
3	D	13	<div> <div>15%</div> <div>15%</div> <div>38%</div> <div>46%</div> </div>
3	H	13	<div> <div>23%</div> <div>38%</div> <div>8%</div> <div>8%</div> <div>46%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed primase/polymerase protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	0	0
			2137	1380	358	389	10			
1	B	251	Total	C	N	O	S	0	0	0
			2010	1302	333	366	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	ARG	GLN	conflict	UNP Q96LW4
B	168	ARG	GLN	conflict	UNP Q96LW4


- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*CP*GP*CP*(5IU)P*AP*CP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	12	Total	C	I	N	O	P	0	0	0
			241	114	1	44	70	12			
2	G	8	Total	C	I	N	O	P	0	0	0
			159	75	1	26	49	8			

- Molecule 3 is a DNA chain called DNA (5'-D(P*GP*GP*TP*AP*GP*CP*(DDG))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	7	Total	C	N	O	P	0	0	0
			147	69	30	41	7			
3	H	7	Total	C	N	O	P	0	0	0
			147	69	30	41	7			

- Molecule 4 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C₁₀H₁₆N₅O₁₂P₃).





- Molecule 1: DNA-directed primase/polymerase protein







- Molecule 2: DNA (5'-D(P*TP*CP*GP*CP*(5IU)P*AP*CP*C)-3')



- Molecule 3: DNA (5'-D(P*GP*GP*TP*AP*GP*CP*(DDG))-3')



- Molecule 3: DNA (5'-D(P*GP*GP*TP*AP*GP*CP*(DDG))-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.72Å 65.21Å 72.46Å 67.89° 85.22° 86.65°	Depositor
Resolution (Å)	39.35 – 2.20 56.58 – 2.04	Depositor EDS
% Data completeness (in resolution range)	89.3 (39.35-2.20) 81.9 (56.58-2.04)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.01	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.05Å)	Xtriage
Refinement program	PHENIX (dev_2264: ???)	Depositor
R, R_{free}	0.206 , 0.255 0.205 , 0.253	Depositor DCC
R_{free} test set	2115 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 59.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5001	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5IU, CA, DTP, DDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2187	0.58	0/2967
1	B	0.47	0/2058	0.67	1/2787 (0.0%)
2	C	0.72	0/246	0.92	0/373
2	G	0.88	0/153	1.00	0/230
3	D	0.77	0/141	1.05	0/216
3	H	0.64	0/141	0.85	0/216
All	All	0.50	0/4926	0.68	1/6789 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	298	ILE	C-N-CA	-5.50	110.74	122.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	262	ASP	Peptide
1	B	71	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2137	0	2031	29	0
1	B	2010	0	1945	23	0
2	C	241	0	132	7	0
2	G	159	0	88	1	0
3	D	147	0	79	4	0
3	H	147	0	79	1	0
4	A	30	0	12	0	0
4	B	30	0	12	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	41	0	0	0	0
6	B	52	0	0	2	0
6	C	2	0	0	0	0
6	G	2	0	0	0	0
6	H	1	0	0	0	0
All	All	5001	0	4378	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:HIS:HD2	1:A:346:CYS:H	1.22	0.88
1:A:108:VAL:HG12	1:A:335:VAL:HG21	1.65	0.79
1:A:191:LYS:NZ	1:B:339:ASP:OD2	2.21	0.73
1:A:184:HIS:CD2	1:A:346:CYS:H	2.08	0.69
1:B:94:LYS:O	1:B:96:LEU:N	2.27	0.67
1:A:308:GLU:H	1:A:308:GLU:CD	1.98	0.66
1:B:91:LYS:O	1:B:93:ARG:N	2.29	0.66
1:A:54:PHE:HD2	1:A:80:VAL:HG11	1.62	0.65
1:B:196:ALA:HB2	1:B:263:LEU:HB2	1.82	0.61
1:B:143:GLN:HE22	1:B:149:ASN:HB3	1.65	0.61
3:D:10:DT:H2"	3:D:11:DA:C8	2.37	0.60
2:C:7:5IU:H2"	2:C:8:DA:C8	2.38	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LYS:HA	1:A:298:ILE:HD11	1.86	0.58
1:A:162:THR:HG22	1:A:302:VAL:HG21	1.87	0.57
3:D:8:DG:H2''	3:D:9:DG:C8	2.41	0.56
1:A:54:PHE:CD2	1:A:80:VAL:HG11	2.40	0.55
1:A:48:GLN:HB2	1:A:78:TYR:CZ	2.43	0.53
1:B:69:CYS:O	1:B:298:ILE:HD13	2.08	0.53
1:A:271:ASN:HD22	1:A:271:ASN:H	1.57	0.53
1:A:184:HIS:HD2	1:A:346:CYS:N	2.00	0.52
3:D:9:DG:H2''	3:D:10:DT:OP2	2.08	0.52
2:G:7:5IU:H2''	2:G:8:DA:C8	2.44	0.52
1:A:62:VAL:CG1	1:A:80:VAL:HG13	2.40	0.52
1:A:285:THR:HG22	2:C:7:5IU:OP1	2.10	0.51
1:B:291:ARG:HG2	1:B:304:LEU:HD12	1.94	0.50
1:A:143:GLN:NE2	1:A:149:ASN:OD1	2.44	0.50
2:C:7:5IU:H2''	2:C:8:DA:N7	2.27	0.50
1:A:93:ARG:HA	1:A:93:ARG:NH1	2.27	0.49
1:A:315:ILE:O	1:A:325:GLN:NE2	2.46	0.49
1:B:325:GLN:NE2	6:B:506:HOH:O	2.45	0.49
1:A:196:ALA:HA	1:A:199:LEU:HD12	1.92	0.49
1:B:68:GLU:HG3	1:B:298:ILE:HB	1.95	0.49
1:B:162:THR:HG22	1:B:302:VAL:HG21	1.94	0.49
1:B:297:LYS:HB2	1:B:300:LYS:HG2	1.95	0.47
1:A:190:ARG:HG2	1:A:281:LEU:HD13	1.95	0.47
1:B:71:VAL:CB	1:B:298:ILE:HG21	2.44	0.47
1:B:67:LEU:HD13	1:B:99:CYS:SG	2.55	0.47
2:C:9:DC:H1'	2:C:10:DC:H5''	1.96	0.47
1:A:116:GLU:HA	1:A:166:PHE:O	2.15	0.47
3:D:8:DG:H2''	3:D:9:DG:H8	1.80	0.47
1:A:128:LYS:O	1:A:131:VAL:HG22	2.15	0.47
1:A:175:HIS:O	1:A:177:VAL:HG23	2.16	0.46
1:B:159:ASP:OD2	1:B:160:SER:N	2.48	0.46
2:C:12:DC:H2''	2:C:13:DA:C8	2.51	0.45
1:A:73:ASP:OD2	1:A:75:GLN:HB2	2.17	0.45
1:B:299:GLY:C	1:B:300:LYS:HD2	2.37	0.45
1:B:81:THR:HG23	1:B:82:THR:O	2.18	0.44
1:B:262:ASP:OD1	6:B:501:HOH:O	2.21	0.44
1:B:270:ASN:OD1	1:B:274:GLU:HG2	2.18	0.43
1:A:191:LYS:HE2	1:A:191:LYS:HB2	1.87	0.43
1:B:83:TYR:CE1	1:B:331:LEU:HD13	2.53	0.43
2:C:6:DC:H3'	2:C:7:5IU:I5	2.90	0.42
1:B:263:LEU:HB3	1:B:266:LEU:HD12	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ASP:C	1:A:200:LEU:H	2.23	0.42
1:B:59:LYS:HB3	1:B:59:LYS:HE3	1.82	0.42
3:H:13:DC:H2''	3:H:14:DDG:H5'	2.02	0.42
1:A:277:LEU:HA	1:A:277:LEU:HD23	1.74	0.41
1:B:289:ASN:N	1:B:289:ASN:HD22	2.18	0.41
1:A:48:GLN:HB3	2:C:4:DC:H5''	2.03	0.41
1:A:137:TYR:OH	1:A:192:ILE:O	2.28	0.41
1:A:120:PRO:O	1:A:123:PRO:HD3	2.21	0.40
1:B:116:GLU:HA	1:B:166:PHE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/353 (75%)	253 (96%)	11 (4%)	1 (0%)	34	37
1	B	247/353 (70%)	232 (94%)	11 (4%)	4 (2%)	9	7
All	All	512/706 (72%)	485 (95%)	22 (4%)	5 (1%)	15	14

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	93	ARG
1	B	95	ASN
1	A	71	VAL
1	B	92	SER
1	B	70	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/319 (70%)	220 (99%)	3 (1%)	69	81
1	B	217/319 (68%)	211 (97%)	6 (3%)	43	56
All	All	440/638 (69%)	431 (98%)	9 (2%)	55	69

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	CYS
1	A	264	SER
1	A	295	SER
1	B	73	ASP
1	B	92	SER
1	B	97	LEU
1	B	149	ASN
1	B	289	ASN
1	B	301	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	A	53	ASN
1	A	143	GLN
1	A	184	HIS
1	A	271	ASN
1	B	143	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	DDG	D	14	3,2	17,23,24	1.21	2 (11%)	15,33,36	2.80	7 (46%)
3	DDG	H	14	3,2	17,23,24	1.32	2 (11%)	15,33,36	2.99	8 (53%)
2	5IU	C	7	3,2	14,21,22	0.86	0	16,30,33	2.11	4 (25%)
2	5IU	G	7	3,2	14,21,22	0.88	0	16,30,33	2.25	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DDG	D	14	3,2	-	0/3/18/19	0/3/3/3
3	DDG	H	14	3,2	-	1/3/18/19	0/3/3/3
2	5IU	C	7	3,2	-	1/4/21/22	0/2/2/2
2	5IU	G	7	3,2	-	1/4/21/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	14	DDG	C6-C5	4.08	1.48	1.41
3	D	14	DDG	C6-C5	3.74	1.47	1.41
3	D	14	DDG	C5-C4	2.35	1.47	1.40
3	H	14	DDG	C5-C4	2.35	1.47	1.40

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	7	5IU	C4-N3-C2	7.06	121.11	115.14
2	C	7	5IU	C4-N3-C2	6.68	120.78	115.14
3	D	14	DDG	C2-N3-C4	5.29	121.39	115.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	14	DDG	C6-C5-C4	-5.28	115.75	120.80
3	H	14	DDG	C2-N3-C4	4.96	121.02	115.36
3	D	14	DDG	C6-N1-C2	4.34	122.83	115.93
3	H	14	DDG	C6-N1-C2	4.30	122.76	115.93
2	G	7	5IU	C6-C5-I5	4.03	121.79	118.52
3	D	14	DDG	C6-C5-C4	-4.00	116.98	120.80
3	D	14	DDG	N3-C2-N1	-4.00	121.89	127.22
3	D	14	DDG	C5-C6-N1	-3.98	117.99	123.43
3	H	14	DDG	C5-C6-N1	-3.64	118.45	123.43
3	H	14	DDG	N3-C2-N1	-3.59	122.44	127.22
2	C	7	5IU	C5-C4-N3	-3.15	119.06	123.27
3	D	14	DDG	C4-C5-N7	-3.11	106.16	109.40
3	H	14	DDG	O4'-C4'-C5'	-3.08	104.45	109.52
3	H	14	DDG	C4-C5-N7	-2.96	106.31	109.40
2	G	7	5IU	C5-C4-N3	-2.83	119.48	123.27
3	D	14	DDG	C3'-C2'-C1'	2.56	105.73	102.78
2	C	7	5IU	C6-C5-I5	2.54	120.58	118.52
3	H	14	DDG	C2'-C3'-C4'	2.33	107.09	102.72
2	C	7	5IU	C5-C6-N1	-2.01	119.57	120.70

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	7	5IU	O4'-C1'-N1-C6
2	G	7	5IU	O4'-C1'-N1-C6
3	H	14	DDG	O4'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	14	DDG	1	0
2	C	7	5IU	4	0
2	G	7	5IU	1	0

5.5 Carbohydrates

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DTP	A	401	5	26,32,32	0.82	1 (3%)	30,50,50	1.54	6 (20%)
4	DTP	B	401	5	26,32,32	1.01	2 (7%)	30,50,50	1.46	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTP	A	401	5	-	6/18/34/34	0/3/3/3
4	DTP	B	401	5	-	6/18/34/34	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	401	DTP	C5-C4	2.51	1.47	1.40
4	A	401	DTP	C5-C4	2.05	1.46	1.40
4	B	401	DTP	C2-N3	2.00	1.35	1.32

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	DTP	N3-C2-N1	-3.73	122.85	128.68
4	A	401	DTP	C2'-C1'-N9	-3.71	105.72	114.27
4	B	401	DTP	C2'-C1'-N9	-3.48	106.25	114.27
4	B	401	DTP	N3-C2-N1	-3.30	123.52	128.68
4	B	401	DTP	C4-C5-N7	-3.27	105.99	109.40
4	B	401	DTP	O2G-PG-O3B	3.19	115.33	104.64
4	A	401	DTP	O2G-PG-O3B	2.93	114.48	104.64
4	A	401	DTP	PA-O3A-PB	-2.65	123.73	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	DTP	PB-O3B-PG	-2.16	125.41	132.83
4	B	401	DTP	PB-O3B-PG	-2.16	125.42	132.83
4	A	401	DTP	O3B-PG-O1G	-2.14	99.31	111.19

There are no chirality outliers.


All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	401	DTP	PB-O3B-PG-O2G
4	B	401	DTP	PB-O3B-PG-O2G
4	A	401	DTP	PA-O3A-PB-O1B
4	A	401	DTP	PB-O3A-PA-O1A
4	B	401	DTP	PB-O3B-PG-O3G
4	B	401	DTP	PA-O3A-PB-O2B
4	B	401	DTP	PB-O3A-PA-O1A
4	B	401	DTP	PA-O3A-PB-O1B
4	B	401	DTP	PB-O3B-PG-O1G
4	A	401	DTP	PB-O3B-PG-O3G
4	A	401	DTP	PA-O3A-PB-O2B
4	A	401	DTP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	271/353 (76%)	0.84	27 (9%)	7 6	35, 52, 84, 98	0
1	B	251/353 (71%)	0.70	16 (6%)	19 18	31, 43, 69, 85	0
2	C	11/17 (64%)	1.98	5 (45%)	0 0	52, 97, 121, 123	0
2	G	7/17 (41%)	1.01	2 (28%)	0 0	45, 49, 103, 115	0
3	D	6/13 (46%)	2.05	2 (33%)	0 0	68, 83, 107, 123	0
3	H	6/13 (46%)	2.56	3 (50%)	0 0	68, 83, 106, 120	0
All	All	552/766 (72%)	0.83	55 (9%)	7 6	31, 48, 85, 123	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	261	PRO	13.1
1	A	262	ASP	8.3
1	A	200	LEU	7.1
3	H	8	DG	6.6
3	D	8	DG	6.5
1	A	72	GLY	5.9
1	A	198	ASP	5.5
1	A	35	PRO	5.5
1	A	12	ILE	5.4
2	C	13	DA	4.8
1	B	72	GLY	4.6
2	C	12	DC	4.3
1	A	163	ASP	4.3
1	A	71	VAL	4.2
1	A	16	ALA	4.1
1	A	199	LEU	3.9
1	B	38	PRO	3.9
2	C	10	DC	3.9
3	H	10	DT	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	9	DG	3.5
1	B	261	PRO	3.5
3	D	9	DG	3.5
1	A	299	GLY	3.4
1	A	70	LYS	3.3
1	B	39	PRO	3.3
1	B	95	ASN	3.3
1	A	263	LEU	3.1
1	A	38	PRO	3.0
2	C	11	DA	3.0
1	B	316	GLN	3.0
1	A	337	PHE	2.9
2	G	10	DC	2.7
1	A	39	PRO	2.7
1	A	315	ILE	2.7
1	A	179	PHE	2.6
2	C	2	DA	2.6
2	G	9	DC	2.6
1	B	94	LYS	2.6
1	A	264	SER	2.5
1	B	148	VAL	2.5
1	B	178	ALA	2.5
1	A	344	LEU	2.5
1	A	194	GLN	2.4
1	A	1	MET	2.4
1	B	192	ILE	2.3
1	B	96	LEU	2.3
1	A	343	ILE	2.3
1	B	142	LEU	2.2
1	A	69	CYS	2.2
1	A	193	LEU	2.2
1	A	302	VAL	2.1
1	B	317	SER	2.1
1	B	70	LYS	2.1
1	B	146	TYR	2.0
1	B	141	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	DDG	D	14	21/22	0.81	0.19	47,51,78,85	0
2	5IU	C	7	20/21	0.81	0.17	80,91,112,345	0
3	DDG	H	14	21/22	0.82	0.18	37,45,73,88	0
2	5IU	G	7	20/21	0.85	0.19	62,68,91,139	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	DTP	A	401	30/30	0.90	0.16	37,44,52,58	0
4	DTP	B	401	30/30	0.94	0.17	30,37,43,47	0
5	CA	B	402	1/1	0.98	0.10	36,36,36,36	0
5	CA	A	402	1/1	0.99	0.07	44,44,44,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DTP A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DTP B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.