



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 28, 2021 – 06:09 PM JST

PDB ID : 6L5G  
Title : Crystal structure of yak lactoperoxidase with disordered heme moiety at 2.50 Å resolution  
Authors : Singh, P.K.; Rani, C.; Sharma, P.; Sharma, S.; Singh, T.P.  
Deposited on : 2019-10-23  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.16
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	 5% 81% 17% .
2	B	2	 50% 50%

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	S	0	0	0
			4770	3037	847	860	26			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0
3	A	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	0	1
			50	38	1	5	6		


- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total K 2 2	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula:  $\text{Zn}$ ).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	354	Total	O	0	0
			354	354		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Lactoperoxidase



#### • Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.82Å 84.71Å 99.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.38 – 2.50 64.37 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (64.38-2.50) 99.9 (64.37-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.20	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.185 , 0.239 0.191 , 0.240	Depositor DCC
$R_{free}$ test set	1139 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtriage
Anisotropy	0.699	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5278	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, NAG, K, HEM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/4898	0.83	1/6645 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	122	ASN	CB-CA-C	5.15	120.70	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4770	0	4688	91	0
2	B	28	0	25	0	0
3	A	42	0	39	1	0
4	A	1	0	0	0	0
5	A	50	0	10	6	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	30	0	40	4	0
9	A	354	0	0	7	0
All	All	5278	0	4802	96	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:CYS:CB	1:A:170:PRO:HA	1.69	1.23
1:A:167:CYS:HB3	1:A:170:PRO:CA	1.75	1.15
5:A:607[A]:HEM:O2D	9:A:703:HOH:O	1.84	0.93
1:A:167:CYS:HB3	1:A:170:PRO:HA	0.93	0.92
1:A:116:GLU:OE2	8:A:612:GOL:O2	1.88	0.91
1:A:120:GLY:HA3	1:A:126:LYS:NZ	1.88	0.88
1:A:123:GLU:HB3	1:A:126:LYS:HG2	1.56	0.88
1:A:360:ARG:NH1	1:A:371:GLU:O	2.08	0.87
1:A:351:HIS:HD1	1:A:437:ASN:HD21	1.26	0.83
1:A:227:LEU:HD22	1:A:270:LEU:HD22	1.65	0.79
1:A:82:ILE:HD13	1:A:480:LEU:HD13	1.68	0.76
1:A:167:CYS:CB	1:A:170:PRO:CA	2.48	0.75
1:A:170:PRO:HB2	1:A:171:PRO:CD	2.18	0.72
1:A:150:LYS:HZ2	1:A:154:GLN:HE22	1.38	0.70
1:A:170:PRO:HB2	1:A:171:PRO:HD2	1.73	0.69
1:A:150:LYS:NZ	1:A:154:GLN:HE22	1.90	0.68
1:A:119:LEU:HD11	1:A:138:ASN:HD22	1.60	0.66
1:A:377:HIS:HA	1:A:380:PHE:CE2	2.31	0.66
1:A:360:ARG:NH2	1:A:389:ASP:OD2	2.30	0.65
5:A:607[B]:HEM:O2D	9:A:701:HOH:O	0.65	0.64
1:A:77:GLU:HG2	1:A:145:PRO:HB3	1.78	0.64
1:A:120:GLY:HA3	1:A:126:LYS:HZ3	1.60	0.64
1:A:123:GLU:HB3	1:A:126:LYS:CG	2.29	0.62
1:A:120:GLY:HA3	1:A:126:LYS:HZ2	1.65	0.61
1:A:118:GLU:OE1	1:A:118:GLU:HA	2.01	0.61
1:A:570:ASN:HD22	1:A:575:ASP:HB3	1.65	0.61
1:A:119:LEU:O	1:A:123:GLU:OE1	2.19	0.61
1:A:130:GLU:OE1	1:A:426:HIS:ND1	2.34	0.61
1:A:423:GLN:OE1	8:A:612:GOL:H32	2.00	0.60
1:A:126:LYS:HE2	1:A:161:PHE:CE2	2.38	0.59
1:A:120:GLY:HA2	1:A:123:GLU:HB2	1.84	0.58
1:A:333:ASN:HD22	1:A:334:SER:N	2.01	0.57
1:A:130:GLU:CD	1:A:426:HIS:HD1	2.08	0.57
1:A:551:ARG:HD3	1:A:584:LYS:HA	1.86	0.57
1:A:349:PHE:HA	1:A:497:ASN:HD21	1.71	0.56
1:A:63:GLN:O	1:A:64:ARG:HB2	2.05	0.56
1:A:330:GLY:HA3	9:A:1002:HOH:O	2.05	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:ILE:HD13	1:A:480:LEU:CD1	2.35	0.56
1:A:94:GLN:H	8:A:614:GOL:H32	1.70	0.55
1:A:333:ASN:HD22	1:A:333:ASN:C	2.09	0.55
1:A:8:ALA:HB3	1:A:9:PRO:HD3	1.89	0.54
1:A:13:VAL:HG22	1:A:14:LYS:O	2.06	0.54
1:A:63:GLN:O	1:A:64:ARG:CB	2.57	0.53
1:A:120:GLY:CA	1:A:126:LYS:NZ	2.68	0.52
1:A:545:GLN:HB2	9:A:845:HOH:O	2.10	0.52
1:A:189:ALA:HB2	1:A:304:ILE:HD12	1.93	0.51
1:A:167:CYS:SG	1:A:170:PRO:HA	2.49	0.51
1:A:221:ASP:HB2	1:A:226:TYR:CZ	2.44	0.51
1:A:95:ASN:ND2	3:A:601:NAG:H83	2.26	0.51
1:A:170:PRO:CB	1:A:171:PRO:CD	2.89	0.50
1:A:120:GLY:HA2	1:A:123:GLU:CB	2.41	0.50
1:A:82:ILE:CD1	1:A:480:LEU:HD13	2.38	0.49
1:A:118:GLU:C	1:A:119:LEU:HG	2.32	0.49
1:A:118:GLU:O	1:A:120:GLY:N	2.35	0.49
1:A:120:GLY:CA	1:A:126:LYS:HZ2	2.25	0.49
1:A:123:GLU:HG3	1:A:125:SER:H	1.77	0.48
1:A:112:ASP:OD2	5:A:607[B]:HEM:O1D	2.31	0.48
1:A:77:GLU:HG2	1:A:145:PRO:CB	2.44	0.47
1:A:541:ARG:O	1:A:545:GLN:HG3	2.14	0.47
1:A:530:TRP:CE2	1:A:531:GLU:HG3	2.49	0.47
1:A:425:THR:HG21	9:A:1007:HOH:O	2.15	0.47
1:A:130:GLU:OE1	1:A:426:HIS:CE1	2.69	0.46
1:A:167:CYS:HB3	1:A:170:PRO:N	2.25	0.46
1:A:169:THR:O	1:A:169:THR:HG22	2.16	0.46
1:A:313:LEU:N	1:A:314:PRO:CD	2.79	0.46
1:A:150:LYS:NZ	1:A:154:GLN:NE2	2.63	0.45
1:A:303:PHE:O	1:A:307:ILE:HG12	2.16	0.45
1:A:138:ASN:O	1:A:161:PHE:HA	2.17	0.45
1:A:237:CYS:HA	1:A:381:PHE:O	2.18	0.44
1:A:118:GLU:C	1:A:120:GLY:H	2.19	0.43
1:A:511:LEU:HA	1:A:511:LEU:HD12	1.90	0.43
1:A:343:PHE:CD1	1:A:518:GLN:HG2	2.53	0.43
5:A:607[A]:HEM:HBD1	9:A:701:HOH:O	2.19	0.42
1:A:50:GLU:OE2	8:A:615:GOL:H32	2.19	0.42
1:A:148:ASP:O	1:A:151:LEU:HB2	2.20	0.42
1:A:98:LEU:HD23	1:A:98:LEU:HA	1.89	0.42
1:A:518:GLN:O	1:A:522:ILE:HG23	2.20	0.42
5:A:607[B]:HEM:CGD	9:A:701:HOH:O	1.65	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:THR:O	1:A:170:PRO:O	2.38	0.41
1:A:260:ILE:HG23	1:A:261:LEU:HD13	2.02	0.41
1:A:358:VAL:HB	1:A:379:LEU:CD1	2.51	0.41
1:A:191:LEU:HA	1:A:253:ASP:HB2	2.03	0.41
1:A:529:TRP:CD1	1:A:531:GLU:HB2	2.55	0.41
1:A:63:GLN:H	1:A:63:GLN:CD	2.23	0.41
1:A:229:PHE:HB3	1:A:247:PRO:HG2	2.03	0.40
1:A:423:GLN:HA	1:A:423:GLN:HE21	1.86	0.40
1:A:313:LEU:HA	1:A:313:LEU:HD23	1.96	0.40
1:A:130:GLU:CD	1:A:426:HIS:ND1	2.72	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	593/595 (100%)	558 (94%)	27 (5%)	8 (1%)	12	21

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	ARG
1	A	169	THR
1	A	170	PRO
1	A	118	GLU
1	A	119	LEU
1	A	173	GLN
1	A	56	ALA
1	A	7	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	518/518 (100%)	478 (92%)	40 (8%)	13	25

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	TRP
1	A	6	CYS
1	A	12	LEU
1	A	14	LYS
1	A	57	LEU
1	A	89	GLU
1	A	98	LEU
1	A	119	LEU
1	A	122	ASN
1	A	124	HIS
1	A	125	SER
1	A	126	LYS
1	A	153	THR
1	A	173	GLN
1	A	175	LEU
1	A	187	LEU
1	A	198	SER
1	A	201	SER
1	A	203	LEU
1	A	232	LYS
1	A	233	LYS
1	A	261	LEU
1	A	268	LEU
1	A	276	LEU
1	A	292	LEU
1	A	317	LEU
1	A	322	GLN
1	A	333	ASN
1	A	347	PHE
1	A	360	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	376	LEU
1	A	464	LEU
1	A	465	LYS
1	A	471	LEU
1	A	480	LEU
1	A	511	LEU
1	A	513	CYS
1	A	520	GLN
1	A	564	LEU
1	A	593	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
1	A	147	ASN
1	A	154	GLN
1	A	333	ASN
1	A	468	GLN
1	A	497	ASN
1	A	570	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	B	1	1,2	14,14,15	0.58	0	17,19,21	0.98	0
2	NAG	B	2	2	14,14,15	0.66	0	17,19,21	1.51	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	O4-C4-C5	3.31	117.51	109.30
2	B	2	NAG	C1-O5-C5	2.45	115.51	112.19
2	B	2	NAG	C3-C4-C5	-2.34	106.06	110.24
2	B	2	NAG	O5-C1-C2	-2.28	107.69	111.29
2	B	2	NAG	C4-C3-C2	-2.09	107.95	111.02

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	GOL	A	611	-	5,5,5	0.17	0	5,5,5	0.46	0
5	HEM	A	607[B]	-	27,50,50	1.13	1 (3%)	17,82,82	1.36	3 (17%)
3	NAG	A	605	1	14,14,15	0.63	0	17,19,21	1.52	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	HEM	A	607[A]	6	27,50,50	1.14	1 (3%)	17,82,82	1.54	3 (17%)
8	GOL	A	613	-	5,5,5	0.17	0	5,5,5	0.46	0
8	GOL	A	614	-	5,5,5	0.18	0	5,5,5	0.48	0
3	NAG	A	604	1	14,14,15	0.70	0	17,19,21	1.60	3 (17%)
8	GOL	A	612	-	5,5,5	0.19	0	5,5,5	0.46	0
8	GOL	A	615	-	5,5,5	0.15	0	5,5,5	0.35	0
3	NAG	A	601	1	14,14,15	0.59	0	17,19,21	1.47	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	611	-	-	4/4/4/4	-
5	HEM	A	607[B]	-	-	0/6/54/54	-
3	NAG	A	605	1	-	0/6/23/26	0/1/1/1
5	HEM	A	607[A]	6	-	0/6/54/54	-
8	GOL	A	613	-	-	4/4/4/4	-
8	GOL	A	614	-	-	4/4/4/4	-
3	NAG	A	604	1	-	0/6/23/26	0/1/1/1
8	GOL	A	612	-	-	3/4/4/4	-
8	GOL	A	615	-	-	2/4/4/4	-
3	NAG	A	601	1	-	3/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	607[B]	HEM	C3B-C2B	-3.12	1.36	1.40
5	A	607[A]	HEM	C3B-C2B	-3.12	1.36	1.40

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	607[A]	HEM	CBD-CAD-C3D	-4.36	104.45	112.48
3	A	601	NAG	O5-C5-C6	-3.17	102.23	107.20
5	A	607[B]	HEM	CBD-CAD-C3D	-3.12	106.72	112.48
3	A	605	NAG	C3-C4-C5	-3.10	104.71	110.24
3	A	601	NAG	C8-C7-N2	3.02	121.22	116.10
3	A	605	NAG	C1-O5-C5	3.02	116.28	112.19

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	604	NAG	O3-C3-C2	-2.72	103.83	109.47
3	A	604	NAG	O7-C7-N2	2.46	126.47	121.95
5	A	607[B]	HEM	C4A-C3A-C2A	2.23	108.55	107.00
5	A	607[A]	HEM	C4A-C3A-C2A	2.23	108.55	107.00
3	A	604	NAG	O5-C5-C6	-2.14	103.85	107.20
3	A	605	NAG	C1-C2-N2	-2.09	106.92	110.49
5	A	607[B]	HEM	CMD-C2D-C1D	2.08	131.66	128.46
5	A	607[A]	HEM	CMD-C2D-C1D	2.08	131.66	128.46

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	611	GOL	O1-C1-C2-C3
8	A	613	GOL	O1-C1-C2-C3
8	A	614	GOL	O1-C1-C2-C3
8	A	614	GOL	C1-C2-C3-O3
8	A	612	GOL	O1-C1-C2-C3
8	A	615	GOL	O1-C1-C2-C3
3	A	601	NAG	C8-C7-N2-C2
3	A	601	NAG	O7-C7-N2-C2
3	A	601	NAG	O5-C5-C6-O6
8	A	613	GOL	O1-C1-C2-O2
8	A	611	GOL	C1-C2-C3-O3
8	A	613	GOL	C1-C2-C3-O3
8	A	611	GOL	O1-C1-C2-O2
8	A	612	GOL	O1-C1-C2-O2
8	A	614	GOL	O1-C1-C2-O2
8	A	614	GOL	O2-C2-C3-O3
8	A	613	GOL	O2-C2-C3-O3
8	A	615	GOL	O1-C1-C2-O2
8	A	611	GOL	O2-C2-C3-O3
8	A	612	GOL	O2-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	607[B]	HEM	3	0
5	A	607[A]	HEM	3	0
8	A	614	GOL	1	0


*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	612	GOL	2	0
8	A	615	GOL	1	0
3	A	601	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## Ligand HEM A 607 (A)





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	595/595 (100%)	0.19	28 (4%)	31 33	18, 29, 75, 163	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	GLY	19.7
1	A	3	GLU	17.2
1	A	2	TRP	16.3
1	A	122	ASN	14.8
1	A	6	CYS	13.5
1	A	169	THR	12.2
1	A	4	VAL	11.2
1	A	1	SER	9.6
1	A	595	ASN	8.9
1	A	119	LEU	7.1
1	A	9	PRO	7.0
1	A	168	PRO	6.5
1	A	7	GLY	6.3
1	A	120	GLY	6.1
1	A	594	GLU	6.0
1	A	121	SER	5.3
1	A	172	TYR	5.2
1	A	11	PRO	4.9
1	A	13	VAL	4.9
1	A	167	CYS	4.9
1	A	593	ARG	3.5
1	A	171	PRO	3.4
1	A	8	ALA	3.2
1	A	10	VAL	2.6
1	A	124	HIS	2.6
1	A	170	PRO	2.5
1	A	127	THR	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	123	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.80	0.23	53,58,66,67	0
2	NAG	B	1	14/15	0.92	0.16	31,37,42,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	GOL	A	613	6/6	0.79	0.28	40,46,46,48	0
8	GOL	A	614	6/6	0.79	0.29	43,46,48,48	0
3	NAG	A	601	14/15	0.84	0.24	56,61,70,81	0
8	GOL	A	615	6/6	0.87	0.25	44,46,47,47	0
3	NAG	A	604	14/15	0.88	0.17	36,43,57,61	0
8	GOL	A	612	6/6	0.88	0.24	51,53,54,55	0
3	NAG	A	605	14/15	0.90	0.15	41,43,48,48	0
8	GOL	A	611	6/6	0.92	0.17	46,47,49,50	0
5	HEM	A	607[B]	43/43	0.95	0.14	19,21,22,22	7
5	HEM	A	607[A]	43/43	0.95	0.14	19,21,24,26	7
6	K	A	608	1/1	0.96	0.12	25,25,25,25	0
4	CA	A	606	1/1	0.99	0.06	19,19,19,19	0
7	ZN	A	610	1/1	0.99	0.09	35,35,35,35	0
6	K	A	609	1/1	0.99	0.13	22,22,22,22	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around NAG A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAG A 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAG A 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around HEM A 607 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 607 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.