



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 14, 2021 – 12:22 PM EDT

PDB ID : 7L5I  
Title : Crystal Structure of Haemophilus influenzae MtsZ at pH 7.0  
Authors : Struwe, M.A.; Luo, Z.; Kappler, U.; Kobe, B.  
Deposited on : 2020-12-22  
Resolution : 1.73 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.18  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	813	 90% 6%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12793 atoms, of which 5964 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trimethylamine-N-oxide reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	763	Total	C	H	N	O	S	0	4	0
			11840	3791	5871	1026	1117	35			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MET	-	initiating methionine	UNP A0A2S9RK57
A	14	SER	-	expression tag	UNP A0A2S9RK57
A	15	TYR	-	expression tag	UNP A0A2S9RK57
A	16	TYR	-	expression tag	UNP A0A2S9RK57
A	17	HIS	-	expression tag	UNP A0A2S9RK57
A	18	HIS	-	expression tag	UNP A0A2S9RK57
A	19	HIS	-	expression tag	UNP A0A2S9RK57
A	20	HIS	-	expression tag	UNP A0A2S9RK57
A	21	HIS	-	expression tag	UNP A0A2S9RK57
A	22	HIS	-	expression tag	UNP A0A2S9RK57
A	23	ASP	-	expression tag	UNP A0A2S9RK57
A	24	TYR	-	expression tag	UNP A0A2S9RK57
A	25	ASP	-	expression tag	UNP A0A2S9RK57
A	26	ILE	-	expression tag	UNP A0A2S9RK57
A	27	PRO	-	expression tag	UNP A0A2S9RK57
A	28	THR	-	expression tag	UNP A0A2S9RK57
A	29	THR	-	expression tag	UNP A0A2S9RK57
A	30	GLU	-	expression tag	UNP A0A2S9RK57
A	31	ASN	-	expression tag	UNP A0A2S9RK57
A	32	LEU	-	expression tag	UNP A0A2S9RK57
A	33	TYR	-	expression tag	UNP A0A2S9RK57
A	34	PHE	-	expression tag	UNP A0A2S9RK57
A	35	GLN	-	expression tag	UNP A0A2S9RK57
A	36	GLY	-	expression tag	UNP A0A2S9RK57
A	37	ALA	-	expression tag	UNP A0A2S9RK57
A	38	MET	-	expression tag	UNP A0A2S9RK57
A	39	GLY	-	expression tag	UNP A0A2S9RK57

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	40	SER	-	expression tag	UNP A0A2S9RK57

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula:  $C_8H_{18}N_2O_4S$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		
2	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		
2	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		

- Molecule 3 is 2-AMINO-5,6-DIMERCAPTO-7-METHYL-3,7,8A,9-TETRAHYDRO-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-4-ONE GUANOSINE DINUCLEOTIDE (three-letter code: MGD) (formula:  $C_{20}H_{26}N_{10}O_{13}P_2S_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	S	0	0
			68	20	21	10	13	2	2		
3	A	1	Total	C	H	N	O	P	S	0	0
			68	20	21	10	13	2	2		

- Molecule 4 is MOLYBDENUM ATOM (three-letter code: MO) (formula: Mo) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mo	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is OXYGEN ATOM (three-letter code: O) (formula: O) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O	0	1
			2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	717	Total 717	O 717	0	0



- Molecule 1: Trimethylamine-N-oxide reductase

Protein	Residue	Score	Category
Protein A	Q823	0.95	High
	V824	0.92	High
	A825	0.90	High
	GLY	0.88	High
	LVS	0.85	High
	ALA	0.82	High
	GLY	0.80	High
	ALA	0.78	High
	LVS	0.75	High
	THR	0.72	High
Protein B	TRP	0.70	High
	LEU	0.68	High
	ASP	0.65	High
	THR	0.62	High
	THR	0.60	High
	S434	0.58	High
	K435	0.55	High
	Y470	0.52	High
	W471	0.50	High
	H479	0.48	High
Protein C	W503	0.45	High
	M527	0.42	High
	A528	0.40	High
	G529	0.38	High
	D530	0.35	High
	M539	0.32	High
	E548	0.30	High
	K608	0.28	High
	D648	0.25	High
	K652	0.22	High
Protein D	E666	0.20	High
	K725	0.18	High
	R736	0.15	High
	E758	0.12	High
	M765	0.10	High
	TRR	0.08	High
	LEU	0.05	High
	GLY	0.02	High
	GLU	0.00	High
	SER	-0.02	High
Protein E	LVS	-0.05	High
	P773	-0.08	High
	L792	-0.10	High
	T800	-0.12	High
	V820	-0.15	High
	E371	-0.18	High
	Y398	-0.20	High
	K354	-0.22	High
	M237	-0.25	High
	T231	-0.28	High
D219	-0.30	High	
Protein F	I214	-0.32	High
	F157	-0.35	High
	R123	-0.38	High
	M99	-0.40	High
	C97	-0.42	High
	T50	-0.45	High
	A43	-0.48	High
	GLU	-0.50	High
	LVS	-0.52	High
	SER	-0.55	High
GLY	-0.58	High	
Protein G	ASP	-0.60	High
	THR	-0.62	High
	ASP	-0.65	High
	HIS	-0.68	High
	HIS	-0.70	High
	TRP	-0.72	High
	THR	-0.75	High
	ALA	-0.78	High
	GLY	-0.80	High
	TYR	-0.82	High
Protein H	SER	-0.85	High
	MET	-0.88	High
	GLU	-0.90	High
	LEU	-0.92	High
	TYR	-0.95	High
	PHE	-0.98	High
	GLN	-1.00	High
	GLY	-1.02	High
	ALA	-1.05	High
	MET	-1.08	High
GLY	-1.10	High	
Protein I	LVS	-1.12	High
	SER	-1.15	High
	GLU	-1.18	High
	ASP	-1.20	High
	THR	-1.22	High
	ASP	-1.25	High
	HIS	-1.28	High
	HIS	-1.30	High
	HIS	-1.32	High
	TYR	-1.35	High
Protein J	ASP	-1.38	High
	ILE	-1.40	High
	PRO	-1.42	High
	THR	-1.45	High
	THR	-1.48	High
	GLU	-1.50	High
	ASN	-1.52	High
	LEU	-1.55	High
	TYR	-1.58	High
	PHE	-1.60	High
Protein K	GLN	-1.62	High
	GLY	-1.65	High
	ALA	-1.68	High
	MET	-1.70	High
	GLY	-1.72	High
	SER	-1.75	High
	LVS	-1.78	High
	GLU	-1.80	High
	A43	-1.82	High
	T50	-1.85	High
Protein L	C97	-1.88	High
	M99	-1.90	High
	R123	-1.92	High
	F157	-1.95	High
	I214	-1.98	High
	D219	-2.00	High
	T231	-2.02	High
	M237	-2.05	High
	K314		

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.20Å 93.68Å 132.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.84 – 1.73 46.84 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.84-1.73) 99.3 (46.84-1.73)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.00 (at 1.73Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.132 , 0.160 0.132 , 0.160	Depositor DCC
$R_{free}$ test set	2000 reflections (2.65%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.512	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	12793	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MGD, O, MO, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/6120	0.64	3/8305 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	ARG	NE-CZ-NH2	8.25	124.42	120.30
1	A	123	ARG	NE-CZ-NH1	-7.63	116.49	120.30
1	A	527	MET	CG-SD-CE	6.04	109.87	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5969	5871	5869	17	0
2	A	45	51	51	1	0
3	A	94	42	39	1	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	2	0	0	1	0
7	A	717	0	0	5	0
All	All	6829	5964	5959	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:LYS:NZ	7:A:1001:HOH:O	1.99	0.95
1:A:725:LYS:NZ	7:A:1008:HOH:O	2.29	0.63
2:A:901:EPE:H82	6:A:908[A]:O:O	2.05	0.56
1:A:97[A]:CYS:SG	1:A:123:ARG:HD2	2.46	0.55
1:A:214[B]:ILE:HD11	1:A:398:TYR:CZ	2.46	0.50

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	761/813 (94%)	744 (98%)	15 (2%)	2 (0%)	41 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	PHE
1	A	237	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	633/671 (94%)	624 (99%)	9 (1%)	67 50

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	758	GLU
1	A	800	THR
1	A	470	TYR
1	A	471	TRP
1	A	479	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MGD	A	903	4	41,52,52	4.86	25 (60%)	43,81,81	3.07	19 (44%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EPE	A	902	-	15,15,15	0.90	1 (6%)	18,20,20	1.82	5 (27%)
2	EPE	A	901	-	15,15,15	0.79	1 (6%)	18,20,20	1.64	4 (22%)
2	EPE	A	907	-	15,15,15	0.77	1 (6%)	18,20,20	2.04	7 (38%)
3	MGD	A	904	4	41,52,52	4.97	24 (58%)	43,81,81	3.46	19 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MGD	A	903	4	-	1/18/66/66	0/6/6/6
2	EPE	A	902	-	-	2/9/19/19	0/1/1/1
2	EPE	A	901	-	-	6/9/19/19	0/1/1/1
2	EPE	A	907	-	-	5/9/19/19	0/1/1/1
3	MGD	A	904	4	-	5/18/66/66	0/6/6/6

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	904	MGD	O11-C11	12.00	1.59	1.43
3	A	903	MGD	O11-C11	11.97	1.59	1.43
3	A	904	MGD	O11-C23	-10.20	1.29	1.43
3	A	903	MGD	C23-C14	-9.21	1.46	1.53
3	A	903	MGD	C2'-C3'	-9.16	1.28	1.53

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	MGD	O11-C23-C14	14.68	118.76	108.96
3	A	903	MGD	O11-C23-N22	-9.40	98.90	108.57
3	A	903	MGD	C17-C16-C21	6.00	119.90	114.57
3	A	903	MGD	C16-C21-N22	5.98	123.60	118.13
3	A	904	MGD	C17-C16-C21	5.66	119.60	114.57

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	907	EPE	C10-C9-N1-C6

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	907	EPE	C8-C7-N4-C3
2	A	907	EPE	S-C10-C9-N1
3	A	904	MGD	PA-O3B-PB-O5'
3	A	904	MGD	C5'-O5'-PB-O1B


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	903	MGD	1	0
2	A	901	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	763/813 (93%)	-0.52	3 (0%) 92 94	11, 18, 34, 82	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	825	ALA	10.0
1	A	824	VAL	3.1
1	A	420	PRO	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EPE	A	901	15/15	0.81	0.26	27,40,93,112	32
5	CL	A	906	1/1	0.85	0.06	57,57,57,57	0
6	O	A	908[A]	1/1	0.85	0.27	27,27,27,27	1
6	O	A	908[B]	1/1	0.85	0.27	18,18,18,18	1


*Continued on next page...*



*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EPE	A	907	15/15	0.90	0.16	29,59,71,82	0
2	EPE	A	902	15/15	0.92	0.17	26,42,51,54	0
3	MGD	A	904	47/47	0.98	0.09	10,13,16,18	0
3	MGD	A	903	47/47	0.98	0.12	9,13,17,20	0
4	MO	A	905	1/1	1.00	0.08	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around O A 908 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MGD A 904:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)


**Electron density around MGD A 903:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MO A 905:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**6.5 Other polymers** ⓘ

There are no such residues in this entry.