



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2021 – 05:19 PM JST

PDB ID : 6L9M  
Title : H2-Ld complexed with AH1 peptide  
Authors : Wei, P.C.; Yin, L.  
Deposited on : 2019-11-10  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.18
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	<div> <div>3%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>
1	D	278	<div> <div>3%</div> <div>79%</div> <div>17%</div> <div>.</div> </div>
1	G	278	<div> <div>2%</div> <div>78%</div> <div>19%</div> <div>.</div> </div>
1	J	278	<div> <div>2%</div> <div>80%</div> <div>17%</div> <div>.</div> </div>
2	B	99	<div> <div>%</div> <div>73%</div> <div>22%</div> <div>5%</div> </div>
2	E	99	<div> <div>69%</div> <div>26%</div> <div>..</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	99	<div> <div>2%</div> <div>71%</div> <div>23%</div> <div>6%</div> </div>
2	K	99	<div> <div>2%</div> <div>68%</div> <div>26%</div> <div>6%</div> </div>
3	C	9	<div> <div>11%</div> <div>89%</div> <div>11%</div> </div>
3	F	9	<div> <div>89%</div> <div>11%</div> </div>
3	I	9	<div> <div>11%</div> <div>78%</div> <div>22%</div> </div>
3	L	9	<div> <div>78%</div> <div>22%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H2-Ld.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	S	41	0	0
			2268	1437	395	426	10			
1	D	278	Total	C	N	O	S	41	0	0
			2268	1437	395	426	10			
1	G	278	Total	C	N	O	S	41	0	0
			2268	1437	395	426	10			
1	J	278	Total	C	N	O	S	41	0	0
			2268	1437	395	426	10			

- Molecule 2 is a protein called b2m.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	10	0	0
			821	524	138	152	7			
2	E	99	Total	C	N	O	S	10	0	0
			821	524	138	152	7			
2	H	99	Total	C	N	O	S	10	0	0
			821	524	138	152	7			
2	K	99	Total	C	N	O	S	10	0	0
			821	524	138	152	7			

- Molecule 3 is a protein called SER-PRO-SER-TYR-VAL-TYR-HIS-GLN-PHE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			81	54	12	15			
3	F	9	Total	C	N	O	0	0	0
			81	54	12	15			
3	I	9	Total	C	N	O	0	0	0
			81	54	12	15			
3	L	9	Total	C	N	O	0	0	0
			81	54	12	15			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total 32	O 32	0	0
4	B	12	Total 12	O 12	0	0
4	D	36	Total 36	O 36	0	0
4	E	21	Total 21	O 21	0	0
4	G	38	Total 38	O 38	0	0
4	H	19	Total 19	O 19	0	0
4	J	32	Total 32	O 32	0	0
4	K	19	Total 19	O 19	0	0
4	L	2	Total 2	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: H2-Ld




#### • Molecule 1: H2-Ld



#### • Molecule 1: H2-Ld




#### • Molecule 1: H2-Ld






- Molecule 2: b2m




- Molecule 2: b2m





- Molecule 2: b2m





- Molecule 2: b2m



- Molecule 3: SER-PRO-SER-TYR-VAL-TYR-HIS-GLN-PHE



- Molecule 3: SER-PRO-SER-TYR-VAL-TYR-HIS-GLN-PHE




- Molecule 3: SER-PRO-SER-TYR-VAL-TYR-HIS-GLN-PHE





- Molecule 3: SER-PRO-SER-TYR-VAL-TYR-HIS-GLN-PHE

Chain L:  78% 22%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.87Å 88.39Å 105.71Å 80.97° 75.96° 88.24°	Depositor
Resolution (Å)	19.97 – 2.60 19.96 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.7 (19.97-2.60) 95.7 (19.96-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.08 (at 2.59Å)	Xtriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, $R_{free}$	0.213 , 0.249 0.204 , 0.246	Depositor DCC
$R_{free}$ test set	1866 reflections (3.53%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtriage
Anisotropy	0.539	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 35.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.018 for h,-k,h-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12891	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.60	2/2337 (0.1%)	0.73	8/3180 (0.3%)
1	D	0.68	6/2337 (0.3%)	0.98	9/3180 (0.3%)
1	G	0.64	3/2337 (0.1%)	0.70	5/3180 (0.2%)
1	J	1.04	7/2337 (0.3%)	0.92	19/3180 (0.6%)
2	B	0.90	4/847 (0.5%)	0.75	3/1148 (0.3%)
2	E	1.01	5/847 (0.6%)	0.78	4/1148 (0.3%)
2	H	0.91	4/847 (0.5%)	0.73	2/1148 (0.2%)
2	K	1.28	10/847 (1.2%)	0.86	5/1148 (0.4%)
3	C	0.34	0/85	0.43	0/114
3	F	0.34	0/85	0.43	0/114
3	I	0.37	0/85	0.46	0/114
3	L	0.41	0/85	0.49	0/114
All	All	0.83	41/13076 (0.3%)	0.82	55/17768 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	0	1

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	22	TYR	CE2-CZ	-17.10	1.16	1.38
1	J	22	TYR	CG-CD2	-17.00	1.17	1.39
1	J	22	TYR	CE1-CZ	-16.81	1.16	1.38
1	J	22	TYR	CG-CD1	-16.59	1.17	1.39
1	J	223	GLU	CD-OE2	-15.01	1.09	1.25

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	35	ARG	NE-CZ-NH1	33.45	137.02	120.30
1	J	223	GLU	OE1-CD-OE2	-17.34	102.50	123.30
1	D	35	ARG	NE-CZ-NH2	-14.88	112.86	120.30
1	J	79	ARG	NE-CZ-NH2	11.54	126.07	120.30
1	G	35	ARG	NE-CZ-NH1	10.39	125.50	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	34	HIS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2268	0	2131	40	0
1	D	2268	0	2131	54	0
1	G	2268	0	2131	40	0
1	J	2268	0	2131	39	0
2	B	821	0	796	17	0
2	E	821	0	796	20	0
2	H	821	0	796	26	0
2	K	821	0	796	26	0
3	C	81	0	70	1	0
3	F	81	0	70	1	0
3	I	81	0	70	2	0
3	L	81	0	70	1	0
4	A	32	0	0	0	0
4	B	12	0	0	0	0
4	D	36	0	0	2	0
4	E	21	0	0	0	0
4	G	38	0	0	4	0
4	H	19	0	0	0	0
4	J	32	0	0	3	0
4	K	19	0	0	1	0
4	L	2	0	0	0	0
All	All	12891	0	11988	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:K:70:PHE:HD2	2:K:78:TYR:CE1	1.74	1.04
1:A:96:GLN:HE22	2:B:60:TRP:HB3	1.18	1.02
1:D:59:TYR:HE1	1:D:63:ILE:HD13	1.22	1.01
2:K:70:PHE:CD2	2:K:78:TYR:HE1	1.84	0.94
1:D:59:TYR:CE1	1:D:63:ILE:HD13	2.02	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/278 (99%)	269 (98%)	7 (2%)	0	100	100
1	D	276/278 (99%)	267 (97%)	9 (3%)	0	100	100
1	G	276/278 (99%)	268 (97%)	8 (3%)	0	100	100
1	J	276/278 (99%)	269 (98%)	7 (2%)	0	100	100
2	B	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
2	E	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
2	H	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
2	K	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	I	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	L	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	1520/1544 (98%)	1473 (97%)	47 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	216 (93%)	16 (7%)	15	31
1	D	232/232 (100%)	215 (93%)	17 (7%)	14	28
1	G	232/232 (100%)	214 (92%)	18 (8%)	12	25
1	J	232/232 (100%)	213 (92%)	19 (8%)	11	22
2	B	94/94 (100%)	86 (92%)	8 (8%)	10	21
2	E	94/94 (100%)	86 (92%)	8 (8%)	10	21
2	H	94/94 (100%)	85 (90%)	9 (10%)	8	16
2	K	94/94 (100%)	86 (92%)	8 (8%)	10	21
3	C	9/9 (100%)	9 (100%)	0	100	100
3	F	9/9 (100%)	9 (100%)	0	100	100
3	I	9/9 (100%)	9 (100%)	0	100	100
3	L	9/9 (100%)	8 (89%)	1 (11%)	6	11
All	All	1340/1340 (100%)	1236 (92%)	104 (8%)	12	25

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	172	LEU
2	H	67	HIS
2	K	70	PHE
1	G	178	THR
1	G	258	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	188	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	H	67	HIS
1	J	96	GLN
1	D	96	GLN
1	A	96	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	278/278 (100%)	-0.12	9 (3%) 47 40	16, 34, 74, 108	10 (3%)
1	D	278/278 (100%)	-0.09	7 (2%) 57 51	15, 33, 74, 109	10 (3%)
1	G	278/278 (100%)	-0.10	6 (2%) 62 56	17, 34, 74, 108	10 (3%)
1	J	278/278 (100%)	-0.11	5 (1%) 68 64	14, 32, 73, 108	10 (3%)
2	B	99/99 (100%)	-0.30	1 (1%) 82 80	15, 30, 64, 89	3 (3%)
2	E	99/99 (100%)	-0.27	0 100 100	17, 30, 63, 88	3 (3%)
2	H	99/99 (100%)	-0.08	2 (2%) 65 60	18, 32, 65, 90	3 (3%)
2	K	99/99 (100%)	-0.18	2 (2%) 65 60	16, 31, 65, 93	3 (3%)
3	C	9/9 (100%)	0.14	1 (11%) 5 3	28, 38, 57, 84	0
3	F	9/9 (100%)	0.14	0 100 100	33, 35, 58, 92	0
3	I	9/9 (100%)	0.14	1 (11%) 5 3	21, 34, 62, 83	0
3	L	9/9 (100%)	-0.24	0 100 100	22, 30, 37, 66	0
All	All	1544/1544 (100%)	-0.13	34 (2%) 62 56	14, 33, 73, 109	52 (3%)

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	54	GLN	4.2
1	A	277	PRO	4.1
1	G	277	PRO	4.1
1	D	277	PRO	3.9
1	A	107	GLY	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.