



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 02:05 AM BST

PDB ID : 3LBS
Title : Crystal structure of the cytoplasmic tail of (pro)renin receptor as a MBP fusion (Maltose-bound form)
Authors : Zhang, Y.; Garavito, R.M.
Deposited on : 2010-01-08
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>.</div> </div> </div>
1	B	384	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>..</div> </div> </div>
2	C	2	<div> <div></div> <div>100%</div> </div>
2	D	2	<div> <div></div> <div>100%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6166 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, Renin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2891	1862	470	553	6			
1	B	376	Total	C	N	O	S	0	0	0
			2923	1880	475	561	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	-	expression tag	UNP P0AEX9
A	2	THR	-	expression tag	UNP P0AEX9
A	137	ARG	LYS	conflict	UNP P0AEX9
A	365	ALA	-	linker	UNP P0AEX9
A	366	ASP	-	linker	UNP P0AEX9
B	1	GLU	-	expression tag	UNP P0AEX9
B	2	THR	-	expression tag	UNP P0AEX9
B	137	ARG	LYS	conflict	UNP P0AEX9
B	365	ALA	-	linker	UNP P0AEX9
B	366	ASP	-	linker	UNP P0AEX9

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	176	Total 176	O 176	0	0
3	B	130	Total 130	O 130	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltose-binding periplasmic protein, Renin receptor




- Molecule 1: Maltose-binding periplasmic protein, Renin receptor



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.95Å 96.78Å 191.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.60 – 2.15 35.60 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.2 (35.60-2.15) 96.3 (35.60-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.82 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.212 , 0.278 0.211 , 0.279	Depositor DCC
R_{free} test set	1330 reflections (3.17%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6166	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1508e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	1/2963 (0.0%)	0.88	2/4025 (0.0%)
1	B	0.87	1/2995 (0.0%)	0.83	0/4068
All	All	0.94	2/5958 (0.0%)	0.86	2/8093 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	TRP	CB-CG	6.88	1.62	1.50
1	B	288	GLU	CG-CD	5.06	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	LEU	CA-CB-CG	5.88	128.82	115.30
1	A	316	ARG	NE-CZ-NH1	5.84	123.22	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2891	0	2848	50	0
1	B	2923	0	2876	56	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	176	0	0	8	0
3	B	130	0	0	4	0
All	All	6166	0	5766	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:LEU:CD2	1:A:60:ILE:HD11	1.83	1.08
1:A:43:LEU:HD21	1:A:60:ILE:HD11	1.36	1.05
1:A:43:LEU:HD21	1:A:60:ILE:CD1	1.87	1.04
1:A:297:LYS:HE3	3:A:457:HOH:O	1.64	0.97
1:A:178:ILE:HD12	1:B:370:ASP:HA	1.49	0.95
1:B:43:LEU:HD13	1:B:60:ILE:HD11	1.51	0.91
1:A:321:MET:HA	1:A:321:MET:HE2	1.65	0.77
1:A:68:GLY:HA3	1:A:332:ASN:O	1.89	0.72
1:A:372:ILE:HA	1:A:375:ARG:CD	2.20	0.72
1:A:297:LYS:CE	3:A:457:HOH:O	2.30	0.72
1:B:43:LEU:CD1	1:B:60:ILE:HD11	2.19	0.70
1:B:246:VAL:CG2	1:B:326:LYS:HE3	2.22	0.69
1:A:370:ASP:HA	1:B:178:ILE:HD12	1.75	0.69
1:A:43:LEU:HD21	1:A:60:ILE:HD12	1.72	0.69
1:A:178:ILE:HD12	1:B:370:ASP:CA	2.22	0.67
1:A:178:ILE:CD1	1:B:370:ASP:HA	2.24	0.67
1:A:79:ILE:HD12	1:A:81:PRO:HD3	1.75	0.67
1:A:321:MET:HA	1:A:321:MET:CE	2.24	0.67
1:B:375:ARG:HB3	1:B:377:THR:HG22	1.77	0.67
1:A:43:LEU:CD2	1:A:60:ILE:CD1	2.59	0.66
1:B:68:GLY:HA3	1:B:332:ASN:O	1.96	0.66
1:B:356:THR:HG23	1:B:359:GLU:CB	2.27	0.65
1:A:370:ASP:HB3	1:B:178:ILE:HG13	1.79	0.64
1:B:256:LYS:HG2	1:B:326:LYS:O	1.98	0.63
1:B:121:LEU:HD11	1:B:223:ALA:HA	1.80	0.62
1:B:43:LEU:HD13	1:B:60:ILE:CD1	2.26	0.62
1:A:342:ALA:HB2	1:A:373:ILE:HD11	1.84	0.59
1:B:356:THR:HG23	1:B:359:GLU:HB2	1.85	0.57
1:B:164:ASP:OD2	1:B:251:LYS:HD3	2.05	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:ILE:HA	1:A:375:ARG:HD3	1.87	0.55
1:B:349:ASN:HB3	1:B:355:GLN:HB2	1.89	0.54
1:B:159:PRO:HG3	1:B:257:PRO:HA	1.88	0.54
1:A:43:LEU:HD23	1:A:60:ILE:HD11	1.80	0.54
1:B:102:LYS:NZ	3:B:473:HOH:O	2.40	0.54
1:A:15:LYS:O	1:A:297:LYS:HD3	2.07	0.53
1:B:254:PRO:HB3	3:B:485:HOH:O	2.08	0.53
1:A:82:ASP:O	1:A:86:GLN:HG3	2.11	0.51
1:A:370:ASP:HA	1:B:178:ILE:CD1	2.39	0.51
1:A:372:ILE:HA	1:A:375:ARG:HD2	1.92	0.51
1:B:128:THR:OG1	1:B:131:GLU:HG2	2.12	0.50
1:A:15:LYS:C	1:A:297:LYS:HD3	2.32	0.49
1:B:356:THR:HG23	1:B:359:GLU:HB3	1.94	0.49
1:B:249:THR:HB	1:B:253:GLN:O	2.12	0.49
1:A:62:TRP:CD1	1:A:66:ARG:HG3	2.47	0.49
1:B:140:LYS:HG2	1:B:144:LYS:O	2.13	0.48
1:A:4:GLU:N	1:A:6:LYS:HG2	2.28	0.48
1:A:297:LYS:CG	3:A:457:HOH:O	2.61	0.48
1:B:117:TYR:O	1:B:242:TYR:HA	2.14	0.48
1:B:307:TYR:CE2	1:B:311:LEU:HD11	2.48	0.48
1:B:122:LEU:HD11	1:B:135:LEU:HD21	1.95	0.47
1:B:190:ALA:O	1:B:193:THR:HG22	2.14	0.47
1:B:219:LYS:NZ	1:B:221:GLU:OE2	2.43	0.47
1:A:277:LYS:HE2	1:A:281:GLU:OE2	2.14	0.47
1:B:172:GLU:O	1:B:173:ASN:C	2.52	0.47
1:A:211:SER:HB2	3:A:523:HOH:O	2.16	0.46
1:A:46:LYS:NZ	3:A:489:HOH:O	2.48	0.46
1:B:148:MET:O	1:B:225:THR:HA	2.15	0.46
1:B:42:LYS:O	1:B:46:LYS:HG2	2.16	0.45
1:B:137:ARG:HB3	1:B:137:ARG:CZ	2.46	0.45
1:A:337:SER:HB3	3:A:433:HOH:O	2.16	0.45
1:A:92:PHE:CZ	1:A:110:VAL:HG21	2.52	0.45
1:A:7:LEU:HB2	1:A:35:VAL:HG22	1.99	0.44
1:A:123:PRO:HD2	3:A:467:HOH:O	2.17	0.44
1:A:374:TYR:HH	1:B:374:TYR:HH	1.58	0.44
1:B:190:ALA:HA	1:B:193:THR:HG22	1.99	0.44
1:B:67:PHE:HB3	1:B:104:ILE:HD12	2.00	0.44
1:B:62:TRP:CD1	1:B:66:ARG:HG3	2.52	0.44
1:A:15:LYS:O	1:A:297:LYS:CD	2.66	0.44
1:A:61:PHE:HA	1:A:263:SER:O	2.18	0.44
1:B:344:ARG:HD3	3:B:495:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:LYS:HE3	1:B:219:LYS:HB2	1.61	0.43
1:A:90:TYR:HA	1:A:91:PRO:HD3	1.92	0.43
1:B:277:LYS:HB2	3:B:419:HOH:O	2.18	0.43
1:B:192:LEU:O	1:B:196:VAL:HG23	2.19	0.43
1:A:370:ASP:O	1:A:375:ARG:NH2	2.52	0.43
1:A:117:TYR:CE2	1:A:125:PRO:HD3	2.54	0.42
1:B:196:VAL:HG12	1:B:200:LYS:HD2	2.00	0.42
1:B:121:LEU:HD11	1:B:223:ALA:CB	2.49	0.42
1:B:130:GLU:N	1:B:130:GLU:OE1	2.50	0.42
1:A:372:ILE:HD12	1:A:375:ARG:HD3	2.00	0.42
1:B:247:LEU:HB2	1:B:255:SER:HB2	2.00	0.42
1:B:167:TYR:OH	1:B:180:ASP:OD1	2.27	0.42
1:A:85:PHE:HA	1:A:88:LYS:HE2	2.02	0.42
1:B:79:ILE:HD12	1:B:81:PRO:HD3	2.01	0.42
1:B:89:LEU:HD22	1:B:94:TRP:CZ2	2.55	0.42
1:A:342:ALA:HB1	1:A:364:ALA:HA	2.02	0.41
1:B:233:SER:O	1:B:236:ASP:HB2	2.19	0.41
1:A:167:TYR:OH	1:A:180:ASP:OD1	2.32	0.41
1:A:169:PHE:CD2	1:A:333:ILE:HD11	2.56	0.41
1:B:121:LEU:HD11	1:B:223:ALA:CA	2.50	0.41
1:B:154:PRO:HG3	1:B:344:ARG:CA	2.50	0.41
1:A:329:ILE:HD12	1:A:329:ILE:HA	1.96	0.41
1:B:3:GLU:N	1:B:6:LYS:HE2	2.35	0.41
1:B:297:LYS:HA	1:B:298:PRO:HD3	1.95	0.41
1:A:175:LYS:HB3	1:A:175:LYS:HE3	1.61	0.41
1:A:291:GLU:O	1:A:295:LYS:HG3	2.20	0.41
1:A:355:GLN:HG2	3:A:445:HOH:O	2.20	0.41
1:B:121:LEU:C	1:B:121:LEU:HD12	2.41	0.41
1:B:356:THR:CG2	1:B:359:GLU:HB2	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/384 (96%)	359 (97%)	11 (3%)	0	100	100
1	B	374/384 (97%)	352 (94%)	20 (5%)	2 (0%)	29	22
All	All	744/768 (97%)	711 (96%)	31 (4%)	2 (0%)	41	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	173	ASN
1	B	120	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/309 (96%)	289 (97%)	8 (3%)	44	46
1	B	301/309 (97%)	292 (97%)	9 (3%)	41	40
All	All	598/618 (97%)	581 (97%)	17 (3%)	43	44

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	LEU
1	A	102	LYS
1	A	179	LYS
1	A	239	LYS
1	A	258	PHE
1	A	328	GLU
1	A	358	ASP
1	A	371	SER
1	B	131	GLU
1	B	137	ARG
1	B	139	LEU
1	B	172	GLU
1	B	179	LYS
1	B	233	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	253	GLN
1	B	258	PHE
1	B	356	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	B	378	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	0.60	0	17,17,17	1.45	3 (17%)
2	GLC	C	2	2	11,11,12	0.59	0	15,15,17	1.20	1 (6%)
2	GLC	D	1	2	12,12,12	0.51	0	17,17,17	1.53	3 (17%)
2	GLC	D	2	2	11,11,12	0.65	0	15,15,17	1.22	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	1/2/19/22	0/1/1/1
2	GLC	D	1	2	-	1/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GLC	C1-O5-C5	3.48	120.23	113.66
2	D	1	GLC	O5-C1-C2	3.24	116.06	110.28
2	C	1	GLC	C1-O5-C5	3.20	119.70	113.66
2	D	2	GLC	C6-C5-C4	-2.90	106.21	113.00
2	C	1	GLC	O2-C2-C3	-2.47	104.64	110.35
2	C	2	GLC	C2-C3-C4	-2.42	106.71	110.89
2	C	1	GLC	O5-C1-C2	2.40	114.57	110.28
2	D	1	GLC	O4-C4-C3	-2.23	105.18	110.35

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	GLC	C4-C5-C6-O6
2	D	1	GLC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/384 (96%)	-0.06	4 (1%) 80 85	15, 27, 44, 61	0
1	B	376/384 (97%)	0.22	20 (5%) 26 35	20, 35, 55, 65	0
All	All	748/768 (97%)	0.08	24 (3%) 47 56	15, 31, 51, 65	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	GLU	4.2
1	B	3	GLU	3.8
1	B	121	LEU	3.5
1	B	173	ASN	3.3
1	A	4	GLU	3.3
1	B	172	GLU	3.1
1	A	174	GLY	3.1
1	B	377	THR	3.1
1	B	137	ARG	3.0
1	B	124	ASN	2.8
1	B	174	GLY	2.8
1	B	141	ALA	2.8
1	B	122	LEU	2.7
1	A	271	PRO	2.6
1	B	378	ASN	2.5
1	B	354	ARG	2.5
1	B	241	ASN	2.4
1	B	175	LYS	2.4
1	B	143	GLY	2.2
1	B	171	TYR	2.1
1	B	324	ALA	2.1
1	B	123	PRO	2.1
1	A	269	ALA	2.1
1	B	112	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	0.95	0.18	14,18,21,25	0
2	GLC	D	1	12/12	0.97	0.17	18,22,26,26	0
2	GLC	C	2	11/12	0.97	0.20	17,18,20,20	0
2	GLC	D	2	11/12	0.98	0.16	14,20,22,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.