



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 01:03 AM BST

PDB ID : 6LCT  
Title : Crystal structure of catalytic inactive chloroplast resolvase NtMOC1 in complex with Holliday junction  
Authors : Yan, J.J.; Hong, S.X.; Guan, Z.Y.; Yin, P.  
Deposited on : 2019-11-19  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	<div> <div>11%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>6%</div> </div> </div>
1	B	169	<div> <div>11%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>
2	C	18	<div> <div>6%</div> <div> <div></div> <div>61%</div> <div>39%</div> </div> </div>
3	D	18	<div> <div>11%</div> <div> <div></div> <div>61%</div> <div>39%</div> </div> </div>
4	E	18	<div> <div>11%</div> <div> <div></div> <div>67%</div> <div>33%</div> </div> </div>
5	F	18	<div> <div>6%</div> <div> <div></div> <div>67%</div> <div>33%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	B	301	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NtMOC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	159	Total	C	N	O	S	0	0	0
			1196	779	207	208	2			
1	B	159	Total	C	N	O	S	0	0	0
			1169	756	202	210	1			

- Molecule 2 is a DNA chain called DNA (5'-D(\*AP\*CP\*AP\*AP\*CP\*AP\*GP\*AP\*GP\*GP\*AP\*TP\*GP\*GP\*AP\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	18	Total	C	N	O	P	0	0	0
			372	177	78	100	17			

- Molecule 3 is a DNA chain called DNA (5'-D(\*GP\*CP\*CP\*TP\*TP\*GP\*CP\*TP\*GP\*GP\*GP\*AP\*CP\*AP\*TP\*CP\*TP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	18	Total	C	N	O	P	0	0	0
			363	175	62	109	17			


- Molecule 4 is a DNA chain called DNA (5'-D(P\*AP\*GP\*CP\*TP\*CP\*CP\*AP\*TP\*CP\*CP\*AP\*GP\*CP\*AP\*AP\*GP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	18	Total	C	N	O	P	0	0	0
			366	173	70	105	18			

- Molecule 5 is a DNA chain called DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*GP\*TP\*CP\*CP\*CP\*TP\*CP\*TP\*GP\*TP\*TP\*GP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	18	Total	C	N	O	P	0	0	0
			364	176	61	110	17			

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	6	Total	O	0	0
			6	6		
7	B	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NtMOC1




- Molecule 1: NtMOC1



- Molecule 2: DNA (5'-D(\*AP\*CP\*AP\*AP\*CP\*AP\*GP\*AP\*GP\*GP\*AP\*TP\*GP\*GP\*AP\*GP\*CP\*T)-3')




- Molecule 3: DNA (5'-D(\*GP\*CP\*CP\*TP\*TP\*GP\*CP\*TP\*GP\*GP\*GP\*AP\*CP\*AP\*TP\*CP\*TP\*T)-3')



- Molecule 4: DNA (5'-D(P\*AP\*GP\*CP\*TP\*CP\*CP\*AP\*TP\*CP\*CP\*AP\*GP\*CP\*AP\*AP\*G  
P\*GP\*C)-3')

Chain E: 



- Molecule 5: DNA (5'-D(\*AP\*AP\*GP\*AP\*TP\*GP\*TP\*CP\*CP\*CP\*TP\*CP\*TP\*GP\*TP\*TP  
\*GP\*T)-3')

Chain F: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.05Å 89.81Å 68.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.35 – 2.55 45.35 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.6 (45.35-2.55) 99.6 (45.35-2.55)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.220 , 0.245 0.220 , 0.245	Depositor DCC
$R_{free}$ test set	1223 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.3	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 70.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3849	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/1229	0.40	0/1670
1	B	0.24	0/1200	0.41	0/1636
2	C	0.52	0/420	0.79	0/648
3	D	0.52	0/405	0.93	0/624
4	E	0.53	0/410	0.83	0/629
5	F	0.53	0/406	1.00	0/625
All	All	0.38	0/4070	0.66	0/5832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1196	0	1211	12	0
1	B	1169	0	1162	8	0
2	C	372	0	199	7	0
3	D	363	0	203	5	0
4	E	366	0	201	6	0
5	F	364	0	207	3	0
6	A	5	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	5	0	0	0	0
7	A	6	0	0	0	0
7	B	3	0	0	0	0
All	All	3849	0	3183	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:LYS:NZ	1:B:181:PRO:O	2.31	0.63
1:A:151:ARG:NH1	6:A:301:SO4:O4	2.35	0.59
5:F:7:DT:H2'	5:F:8:DC:C6	2.39	0.57
4:E:15:DA:H2''	4:E:16:DG:H5''	1.86	0.56
3:D:4:DT:H2''	3:D:5:DT:H5''	1.89	0.54
1:A:132:PRO:HG2	1:A:244:LEU:HD11	1.92	0.51
3:D:16:DC:H2'	3:D:17:DT:C6	2.46	0.51
1:A:145:GLY:N	2:C:13:DG:OP2	2.33	0.51
1:A:250:ARG:H	1:A:250:ARG:HH21	1.61	0.49
4:E:16:DG:H2''	4:E:17:DG:C8	2.48	0.49
1:B:113:ILE:HB	1:B:172:VAL:HG22	1.93	0.49
1:A:112:VAL:HG21	1:A:267:LYS:HB2	1.95	0.48
2:C:3:DA:H2'	2:C:4:DA:C8	2.48	0.48
1:B:199:ILE:HD13	1:B:209:VAL:HG11	1.96	0.47
1:A:158:VAL:HG12	1:A:201:ILE:HG23	1.95	0.47
1:B:139:HIS:ND1	1:B:150:LYS:HE3	2.29	0.47
2:C:3:DA:H2''	2:C:4:DA:O5'	2.15	0.47
1:A:250:ARG:H	1:A:250:ARG:NH2	2.12	0.46
4:E:2:DG:H2''	4:E:3:DC:H5'	1.98	0.46
5:F:1:DA:H2''	5:F:2:DA:H5''	1.98	0.45
1:B:145:GLY:N	3:D:13:DC:OP2	2.41	0.45
1:A:185:LYS:HE3	2:C:11:DA:OP2	2.16	0.45
1:B:185:LYS:HG3	3:D:10:DG:OP2	2.19	0.43
1:A:127:LYS:HB2	1:A:130:GLN:HB2	2.01	0.43
2:C:2:DC:H2''	2:C:3:DA:C8	2.55	0.42
4:E:16:DG:H2''	4:E:17:DG:H8	1.85	0.42
3:D:2:DC:H42	4:E:17:DG:H1	1.67	0.42
1:A:185:LYS:HB3	1:B:188:TRP:CD2	2.55	0.41
1:A:244:LEU:HB3	1:A:247:LEU:HD12	2.02	0.41
1:B:195:TYR:O	1:B:199:ILE:HG12	2.20	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:14:DG:H2''	2:C:15:DA:C8	2.56	0.41
5:F:14:DG:H2'	5:F:15:DT:C6	2.56	0.41
2:C:3:DA:H4'	2:C:4:DA:OP1	2.20	0.41
4:E:14:DA:H2'	4:E:15:DA:C8	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/169 (92%)	153 (99%)	2 (1%)	0	100	100
1	B	157/169 (93%)	154 (98%)	3 (2%)	0	100	100
All	All	312/338 (92%)	307 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/134 (91%)	118 (97%)	4 (3%)	38	51
1	B	117/134 (87%)	109 (93%)	8 (7%)	16	20
All	All	239/268 (89%)	227 (95%)	12 (5%)	24	33

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	LYS
1	A	231	ASP
1	A	240	LEU
1	A	250	ARG
1	B	141	LYS
1	B	212	VAL
1	B	214	SER
1	B	227	ARG
1	B	231	ASP
1	B	235	GLN
1	B	239	GLN
1	B	240	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	SO4	A	301	-	4,4,4	0.13	0	6,6,6	0.06	0
6	SO4	B	301	-	4,4,4	0.17	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	301	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	159/169 (94%)	1.00	19 (11%) 4 5	56, 98, 155, 181	0
1	B	159/169 (94%)	0.95	19 (11%) 4 5	56, 95, 187, 221	0
2	C	18/18 (100%)	0.55	1 (5%) 24 29	74, 111, 160, 179	0
3	D	18/18 (100%)	0.73	2 (11%) 5 7	69, 119, 152, 174	0
4	E	18/18 (100%)	0.76	2 (11%) 5 7	74, 135, 173, 178	0
5	F	18/18 (100%)	0.78	1 (5%) 24 29	77, 124, 155, 171	0
All	All	390/410 (95%)	0.92	44 (11%) 5 6	56, 101, 171, 221	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	LYS	6.5
1	B	219	SER	5.1
1	A	250	ARG	5.0
1	B	224	THR	4.1
3	D	1	DG	3.9
1	B	217	TRP	3.9
1	B	235	GLN	3.8
1	A	251	LYS	3.8
4	E	1	DA	3.6
1	B	244	LEU	3.6
1	A	239	GLN	3.4
1	B	216	ALA	3.4
1	B	228	SER	3.4
1	B	226	GLU	3.4
1	A	252	LYS	3.1
1	A	238	SER	2.8
1	A	221	PHE	2.7
1	B	223	LEU	2.7
1	A	241	PHE	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
5	F	18	DT	2.7
1	A	232	TYR	2.7
1	B	231	ASP	2.7
3	D	2	DC	2.6
1	B	239	GLN	2.6
1	A	261	LEU	2.5
1	B	119	THR	2.5
4	E	18	DC	2.4
1	A	195	TYR	2.4
1	A	257	ALA	2.4
1	A	192	GLY	2.4
1	B	194	GLY	2.4
1	B	241	PHE	2.4
1	A	115	VAL	2.3
2	C	18	DT	2.3
1	B	164	PHE	2.3
1	A	143	LEU	2.3
1	A	190	SER	2.2
1	A	189	TRP	2.2
1	A	188	TRP	2.2
1	B	188	TRP	2.2
1	A	191	GLY	2.1
1	B	196	GLY	2.1
1	B	201	ILE	2.0
1	B	198	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	SO4	B	301	5/5	0.69	0.46	142,143,145,146	0
6	SO4	A	301	5/5	0.87	0.45	136,136,139,140	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SO4 B 301:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SO4 A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.