



wwPDB EM Validation Summary Report ⓘ

Nov 15, 2022 – 02:25 PM JST

PDB ID : 6LDI
EMDB ID : EMD-0874
Title : The cryo-EM structure of E. coli CueR transcription activation complex
Authors : Fang, C.L.; Zhang, Y.
Deposited on : 2019-11-21
Resolution : 3.69 Å(reported)
Based on initial models : 6B6H, 4WLS

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	
1	B	329	
2	C	1342	
3	D	1416	
4	E	91	
5	F	633	
6	1	50	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	2	50	<div><div></div><div>52%</div><div>48%</div></div>
8	3	5	<div><div></div><div>100%</div></div>
9	G	139	<div><div></div><div>24%</div><div>78%</div><div>13%</div><div>9%</div></div>
9	H	139	<div><div></div><div>27%</div><div>73%</div><div>19%</div><div>8%</div></div>

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 32870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	230	Total	C	N	O	S	0	0
			1784	1112	317	349	6		
1	B	227	Total	C	N	O	S	0	0
			1748	1090	306	346	6		

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	1338	Total	C	N	O	S	0	0
			10522	6612	1823	2044	43		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	1331	Total	C	N	O	S	0	0
			10290	6482	1814	1945	49		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1408	LEU	-	expression tag	UNP P0A8T7
D	1409	GLU	-	expression tag	UNP P0A8T7
D	1410	VAL	-	expression tag	UNP P0A8T7
D	1411	HIS	-	expression tag	UNP P0A8T7
D	1412	HIS	-	expression tag	UNP P0A8T7
D	1413	HIS	-	expression tag	UNP P0A8T7
D	1414	HIS	-	expression tag	UNP P0A8T7
D	1415	HIS	-	expression tag	UNP P0A8T7
D	1416	HIS	-	expression tag	UNP P0A8T7

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	76	Total	C	N	O	S	0	0
			605	368	115	121	1		

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	F	465	Total	C	N	O	S	0	0
			3748	2352	663	710	23		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	initiating methionine	UNP P00579
F	-18	GLY	-	expression tag	UNP P00579
F	-17	SER	-	expression tag	UNP P00579
F	-16	SER	-	expression tag	UNP P00579
F	-15	HIS	-	expression tag	UNP P00579
F	-14	HIS	-	expression tag	UNP P00579
F	-13	HIS	-	expression tag	UNP P00579
F	-12	HIS	-	expression tag	UNP P00579
F	-11	HIS	-	expression tag	UNP P00579
F	-10	HIS	-	expression tag	UNP P00579
F	-9	SER	-	expression tag	UNP P00579
F	-8	SER	-	expression tag	UNP P00579
F	-7	GLY	-	expression tag	UNP P00579
F	-6	LEU	-	expression tag	UNP P00579
F	-5	VAL	-	expression tag	UNP P00579
F	-4	PRO	-	expression tag	UNP P00579
F	-3	ARG	-	expression tag	UNP P00579
F	-2	GLY	-	expression tag	UNP P00579
F	-1	SER	-	expression tag	UNP P00579
F	0	HIS	-	expression tag	UNP P00579

- Molecule 6 is a DNA chain called DNA (50-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	1	50	Total	C	N	O	P	0	0
			1022	488	181	304	49		

- Molecule 7 is a DNA chain called DNA (50-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
7	2	50	Total	C	N	O	P	0	0
			1035	490	206	290	49		

- Molecule 8 is a RNA chain called RNA (5'-R(*CP*UP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
8	3	5	Total	C	N	O	P	0	0
			102	47	18	33	4		

- Molecule 9 is a protein called HTH-type transcriptional regulator CueR.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	G	127	Total	C	N	O	S	0	0
			1009	621	183	200	5		
9	H	128	Total	C	N	O	S	0	0
			1001	620	182	194	5		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-3	GLY	-	expression tag	UNP P0A9G4
G	-2	ALA	-	expression tag	UNP P0A9G4
G	-1	MET	-	expression tag	UNP P0A9G4
G	0	ALA	-	expression tag	UNP P0A9G4
H	-3	GLY	-	expression tag	UNP P0A9G4
H	-2	ALA	-	expression tag	UNP P0A9G4
H	-1	MET	-	expression tag	UNP P0A9G4
H	0	ALA	-	expression tag	UNP P0A9G4

- Molecule 10 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
10	D	2	Total	Zn	0
			2	2	

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
11	D	1	Total	Mg	0
			1	1	


- Molecule 12 is SILVER ION (three-letter code: AG) (formula: Ag) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
12	H	1	Total 1	Ag 1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha





- Molecule 2: DNA-directed RNA polymerase subunit beta













• Molecule 3: DNA-directed RNA polymerase subunit beta'



- Molecule 4: DNA-directed RNA polymerase subunit omega



- Molecule 5: RNA polymerase sigma factor RpoD




- Molecule 6: DNA (50-MER)



- Molecule 7: DNA (50-MER)






- Molecule 8: RNA (5'-R(*CP*UP*CP*GP*A)-3')

Chain 3: 100%

There are no outlier residues recorded for this chain.


- Molecule 9: HTH-type transcriptional regulator CueR

Chain G: 24% 78% 13% 9%



- Molecule 9: HTH-type transcriptional regulator CueR

Chain H: 27% 73% 19% 8%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	184524	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	56	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.124	Depositor
Minimum map value	-0.063	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	392.1, 392.1, 392.1	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.307, 1.307, 1.307	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AG, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1806	0.49	0/2446
1	B	0.29	0/1770	0.53	0/2400
2	C	0.34	0/10691	0.51	1/14425 (0.0%)
3	D	0.34	1/10445 (0.0%)	0.54	2/14105 (0.0%)
4	E	0.29	0/607	0.49	0/817
5	F	0.31	1/3798 (0.0%)	0.53	2/5105 (0.0%)
6	1	0.82	0/1144	1.01	0/1765
7	2	0.84	0/1166	0.90	2/1800 (0.1%)
8	3	1.09	0/113	1.06	0/174
9	G	0.26	0/1022	0.49	0/1379
9	H	0.27	0/1014	0.53	0/1368
All	All	0.39	2/33576 (0.0%)	0.57	7/45784 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
3	D	0	2
5	F	0	3
All	All	0	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1020	TRP	CD2-CE2	6.14	1.48	1.41
5	F	315	TRP	CD2-CE2	6.13	1.48	1.41

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1020	TRP	CD2-CE3-CZ3	8.14	129.39	118.80
5	F	315	TRP	CD2-CE3-CZ3	8.13	129.37	118.80
5	F	315	TRP	CE2-CD2-CE3	5.14	124.87	118.70
7	2	14	DC	O4'-C4'-C3'	-5.09	102.46	104.50
3	D	1020	TRP	CE2-CD2-CE3	5.08	124.79	118.70

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	1324	ASN	Peptide
3	D	1184	ASP	Peptide
3	D	76	LYS	Peptide
5	F	258	GLN	Peptide
5	F	396	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1784	0	1811	17	0
1	B	1748	0	1757	23	0
2	C	10522	0	10513	86	0
3	D	10290	0	10460	85	0
4	E	605	0	612	4	0
5	F	3748	0	3791	30	0
6	1	1022	0	567	14	0
7	2	1035	0	560	21	0
8	3	102	0	55	0	0
9	G	1009	0	1002	13	0
9	H	1001	0	991	18	0
10	D	2	0	0	0	0
11	D	1	0	0	0	0
12	H	1	0	0	0	0
All	All	32870	0	32119	284	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 284 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:1:46:DT:OP2	9:H:18:ARG:NH1	2.27	0.68
5:F:259:PHE:HB2	5:F:261:LEU:HG	1.76	0.67
2:C:180:ARG:HH22	2:C:465:ARG:HE	1.43	0.65
7:2:38:DA:H5'	9:H:60:LEU:HD23	1.77	0.65
2:C:401:GLY:O	2:C:405:PHE:HB2	1.96	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/329 (69%)	219 (96%)	9 (4%)	0	100	100
1	B	225/329 (68%)	211 (94%)	14 (6%)	0	100	100
2	C	1336/1342 (100%)	1265 (95%)	71 (5%)	0	100	100
3	D	1325/1416 (94%)	1233 (93%)	92 (7%)	0	100	100
4	E	74/91 (81%)	70 (95%)	4 (5%)	0	100	100
5	F	461/633 (73%)	438 (95%)	23 (5%)	0	100	100
9	G	125/139 (90%)	117 (94%)	8 (6%)	0	100	100
9	H	126/139 (91%)	117 (93%)	9 (7%)	0	100	100
All	All	3900/4418 (88%)	3670 (94%)	230 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/286 (69%)	198 (100%)	0	100	100
1	B	193/286 (68%)	193 (100%)	0	100	100
2	C	1149/1157 (99%)	1148 (100%)	1 (0%)	93	98
3	D	1109/1177 (94%)	1107 (100%)	2 (0%)	93	97
4	E	65/75 (87%)	65 (100%)	0	100	100
5	F	408/557 (73%)	407 (100%)	1 (0%)	93	97
9	G	112/118 (95%)	112 (100%)	0	100	100
9	H	108/118 (92%)	108 (100%)	0	100	100
All	All	3342/3774 (89%)	3338 (100%)	4 (0%)	93	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	306	THR
3	D	580	TRP
3	D	1261	LEU
5	F	148	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
9	G	68	ASN
9	H	110	ASN
9	H	55	GLN
3	D	665	GLN
5	F	461	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	3	4/5 (80%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-0874. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 130




Y Index: 138




Z Index: 144


6.3.2 Raw map



X Index: 152



Y Index: 141




Z Index: 144

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_0874_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 219 nm³; this corresponds to an approximate mass of 198 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.271 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.271 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.69	-	-
Author-provided FSC curve	3.67	4.20	3.75
Unmasked-calculated*	4.23	6.66	4.30

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.23 differs from the reported value 3.69 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-0874 and PDB model 6LDI. Per-residue inclusion information can be found in section [3](#) on page [8](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 75% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7513	<div></div> 0.4710
1	<div></div> 0.9051	<div></div> 0.4500
2	<div></div> 0.8850	<div></div> 0.4420
3	<div></div> 0.9412	<div></div> 0.5100
A	<div></div> 0.8055	<div></div> 0.5080
B	<div></div> 0.7477	<div></div> 0.4760
C	<div></div> 0.7948	<div></div> 0.4950
D	<div></div> 0.7598	<div></div> 0.4810
E	<div></div> 0.7334	<div></div> 0.4910
F	<div></div> 0.6145	<div></div> 0.4120
G	<div></div> 0.5538	<div></div> 0.3970
H	<div></div> 0.5168	<div></div> 0.3690

1.0

0.0

<0.0