



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 10, 2022 – 10:16 AM EST

PDB ID : 7LFC  
Title : Structure of importin  $\alpha 3$  bound to p50 NLS  
Authors : Florio, T.J.; Lokareddy, R.K.; Cingolani, G.  
Deposited on : 2021-01-16  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.25  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.25


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	
2	B	14	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	0	0
			3164	2009	540	602	13			

- Molecule 2 is a protein called Nuclear factor NF-kappa-B p105 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	S	0	0	0
			102	63	22	16	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Importin subunit alpha-3



- Molecule 2: Nuclear factor NF-kappa-B p105 subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.65Å 59.07Å 86.09Å 90.00° 96.34° 90.00°	Depositor
Resolution (Å)	14.93 – 2.10 14.93 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.7 (14.93-2.10) 98.7 (14.93-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.18	Depositor
R, $R_{free}$	0.187 , 0.220 0.195 , 0.229	Depositor DCC
$R_{free}$ test set	1381 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.9	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3371	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/3222	0.58	1/4405 (0.0%)
2	B	0.58	0/102	0.74	0/133
All	All	0.44	0/3324	0.59	1/4538 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	ARG	CG-CD-NE	-5.55	100.15	111.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3164	0	3179	35	0
2	B	102	0	111	12	0
3	A	103	0	0	1	0
3	B	2	0	0	0	0
All	All	3371	0	3290	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:GLU:N	1:A:148:GLU:OE1	2.00	0.93
1:A:276:GLN:HA	1:A:279:MET:HE3	1.56	0.85
2:B:436:ARG:HB3	2:B:436:ARG:HH11	1.45	0.81
1:A:229:ARG:CD	2:B:436:ARG:HD2	2.18	0.73
1:A:229:ARG:HD2	2:B:436:ARG:HD2	1.72	0.71
1:A:378:ASP:OD1	1:A:379:LYS:NZ	2.23	0.67
1:A:426:GLN:O	1:A:430:VAL:HG23	1.94	0.67
1:A:102:ASP:OD2	1:A:104:ASN:N	2.28	0.65
1:A:460:GLU:HA	1:A:463:GLU:HG3	1.81	0.62
2:B:436:ARG:HH11	2:B:436:ARG:CB	2.11	0.62
1:A:395:LEU:HD13	1:A:407:LEU:HD21	1.81	0.62
1:A:257:ASN:ND2	3:A:601:HOH:O	2.24	0.61
1:A:350:LEU:HD13	1:A:373:ILE:HD11	1.82	0.61
1:A:222:TRP:CE2	2:B:438:ARG:HD3	2.35	0.61
2:B:435:GLN:HA	2:B:435:GLN:OE1	1.98	0.59
1:A:386:LYS:HG3	1:A:427:VAL:HG22	1.84	0.58
1:A:414:PRO:HA	1:A:456:CYS:SG	2.44	0.58
1:A:229:ARG:HH22	2:B:434:VAL:HG13	1.68	0.57
1:A:175:GLU:HG3	1:A:216:PHE:CD1	2.42	0.54
1:A:97:LYS:HA	2:B:442:MET:CE	2.38	0.53
1:A:276:GLN:HA	1:A:279:MET:CE	2.36	0.52
2:B:436:ARG:HH11	2:B:436:ARG:CG	2.24	0.50
1:A:298:VAL:HG21	1:A:339:LYS:HD3	1.93	0.49
1:A:461:LYS:O	1:A:465:LEU:HD23	2.12	0.48
1:A:460:GLU:HA	1:A:463:GLU:CG	2.43	0.48
1:A:97:LYS:HA	2:B:442:MET:HE1	1.96	0.47
1:A:448:THR:O	1:A:451:ASN:HB2	2.15	0.46
1:A:417:CYS:HA	1:A:420:LEU:HD23	1.98	0.46
1:A:201:VAL:HG21	1:A:239:THR:HG23	1.98	0.45
1:A:337:HIS:ND1	1:A:338:PRO:HD2	2.31	0.45
1:A:395:LEU:CD1	1:A:407:LEU:HD21	2.47	0.44
2:B:434:VAL:HG13	2:B:434:VAL:O	2.16	0.44
1:A:434:GLY:O	1:A:438:ILE:HG12	2.18	0.44
1:A:358:GLN:HE21	1:A:359:GLN:NE2	2.15	0.44
1:A:394:ASN:O	1:A:397:ILE:HG12	2.18	0.44
2:B:436:ARG:CG	2:B:436:ARG:NH1	2.80	0.43
1:A:80:ALA:O	1:A:88:GLN:HG3	2.19	0.43
1:A:465:LEU:HD12	1:A:473:ILE:CG2	2.49	0.42
1:A:412:VAL:O	1:A:415:PRO:HD2	2.20	0.42
1:A:395:LEU:HD23	1:A:395:LEU:O	2.20	0.42
1:A:125:ARG:HG3	1:A:131:LEU:HD23	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/521 (79%)	406 (98%)	6 (2%)	0	100	100
2	B	10/14 (71%)	9 (90%)	1 (10%)	0	100	100
All	All	422/535 (79%)	415 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/460 (77%)	352 (99%)	2 (1%)	86	90
2	B	11/14 (79%)	9 (82%)	2 (18%)	1	1
All	All	365/474 (77%)	361 (99%)	4 (1%)	73	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ARG
1	A	216	PHE
2	B	435	GLN
2	B	436	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	85	GLN
1	A	358	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	414/521 (79%)	0.57	53 (12%) <b>3</b> <b>4</b>	33, 55, 118, 139	0
2	B	12/14 (85%)	1.59	4 (33%) <b>0</b> <b>0</b>	52, 80, 105, 109	0
All	All	426/535 (79%)	0.60	57 (13%) <b>3</b> <b>4</b>	33, 55, 118, 139	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443	GLU	6.2
2	B	435	GLN	5.8
1	A	482	ASP	5.1
1	A	397	ILE	4.6
1	A	484	PHE	4.6
1	A	464	GLN	4.5
2	B	434	VAL	4.3
1	A	478	TYR	4.3
1	A	462	ILE	4.2
1	A	480	ILE	4.0
1	A	444	ASP	3.7
1	A	463	GLU	3.6
1	A	456	CYS	3.5
1	A	423	LYS	3.5
1	A	103	ARG	3.5
1	A	473	ILE	3.4
1	A	465	LEU	3.4
1	A	447	GLU	3.4
1	A	445	GLU	3.4
1	A	481	ILE	3.3
1	A	440	LYS	3.2
1	A	78	GLN	3.1
1	A	74	GLU	3.0
1	A	452	LEU	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	211	SER	2.9
2	B	432	GLU	2.9
1	A	457	GLY	2.9
1	A	467	ASN	2.9
1	A	474	TYR	2.9
1	A	477	ALA	2.9
1	A	468	HIS	2.8
1	A	472	ASP	2.7
1	A	101	SER	2.7
1	A	454	GLU	2.7
1	A	470	ASN	2.6
1	A	436	SER	2.6
1	A	459	LEU	2.6
1	A	460	GLU	2.6
1	A	229	ARG	2.5
1	A	469	GLU	2.5
1	A	359	GLN	2.5
1	A	409	GLN	2.5
1	A	231	LYS	2.5
1	A	220	VAL	2.5
1	A	471	GLU	2.5
1	A	461	LYS	2.4
1	A	455	GLU	2.4
1	A	451	ASN	2.4
1	A	453	ILE	2.3
1	A	485	PHE	2.2
1	A	127	ASP	2.2
1	A	232	ASP	2.1
1	A	115	ILE	2.1
1	A	83	ASP	2.1
1	A	148	GLU	2.0
2	B	443	PRO	2.0
1	A	104	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.