



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 11:05 pm BST

PDB ID : 2LGP
Title : Solution structure of LA45 from LDLR
Authors : Guttman, M.; Komives, E.A.
Deposited on : 2011-08-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	94	

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:127-A:163 (37)	0.25	2
2	A:177-A:209 (33)	0.16	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 8, 9, 10, 13, 15, 17, 18
2	6, 14, 16
3	4, 11
4	7, 12
5	19, 20

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1310 atoms, of which 604 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Low-density lipoprotein receptor.

Mol	Chain	Residues	Atoms						Trace
1	A	94	Total	C	H	N	O	S	0
			1308	423	604	119	150	12	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	GLY	-	EXPRESSION TAG	UNP P01130
A	122	SER	-	EXPRESSION TAG	UNP P01130

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	
2	A	2	Total	Ca
			2	2

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Low-density lipoprotein receptor




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Low-density lipoprotein receptor



4.2.2 Score per residue for model 2 (medoid)


- Molecule 1: Low-density lipoprotein receptor






4.2.3 Score per residue for model 3

- Molecule 1: Low-density lipoprotein receptor




4.2.4 Score per residue for model 4

- Molecule 1: Low-density lipoprotein receptor




4.2.5 Score per residue for model 5

- Molecule 1: Low-density lipoprotein receptor




4.2.6 Score per residue for model 6

- Molecule 1: Low-density lipoprotein receptor




4.2.7 Score per residue for model 7

- Molecule 1: Low-density lipoprotein receptor




4.2.8 Score per residue for model 8

- Molecule 1: Low-density lipoprotein receptor




4.2.9 Score per residue for model 9

- Molecule 1: Low-density lipoprotein receptor




4.2.10 Score per residue for model 10

- Molecule 1: Low-density lipoprotein receptor




4.2.11 Score per residue for model 11

- Molecule 1: Low-density lipoprotein receptor




4.2.12 Score per residue for model 12

- Molecule 1: Low-density lipoprotein receptor




4.2.13 Score per residue for model 13

- Molecule 1: Low-density lipoprotein receptor




4.2.14 Score per residue for model 14

- Molecule 1: Low-density lipoprotein receptor




4.2.15 Score per residue for model 15

- Molecule 1: Low-density lipoprotein receptor




4.2.16 Score per residue for model 16

- Molecule 1: Low-density lipoprotein receptor




4.2.17 Score per residue for model 17

- Molecule 1: Low-density lipoprotein receptor




4.2.18 Score per residue for model 18

- Molecule 1: Low-density lipoprotein receptor




4.2.19 Score per residue for model 19

- Molecule 1: Low-density lipoprotein receptor



4.2.20 Score per residue for model 20

- Molecule 1: Low-density lipoprotein receptor



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	2
CNSSOLVE	refinement	2
CNSSOLVE	structure solution	2
X-PLOR NIH	refinement	
X-PLOR NIH	geometry optimization	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	997
Number of shifts mapped to atoms	997
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
CA

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	535	440	438	14±3
All	All	10740	8800	8760	288

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:184:LEU:HD13	1:A:205:SER:CB	0.79	2.07	11	5
1:A:132:PHE:CD2	1:A:145:ALA:HB2	0.73	2.18	20	9
1:A:132:PHE:CG	1:A:145:ALA:HB2	0.67	2.25	19	9
1:A:184:LEU:HD22	1:A:205:SER:OG	0.65	1.92	15	5
1:A:144:TRP:O	1:A:146:CYS:N	0.62	2.32	4	11
1:A:184:LEU:HD13	1:A:205:SER:OG	0.60	1.96	17	7
1:A:184:LEU:HD13	1:A:205:SER:HB3	0.59	1.74	5	4
1:A:189:ILE:HD12	1:A:193:TRP:HB2	0.58	1.75	12	19
1:A:156:SER:HA	1:A:159:TRP:CG	0.57	2.34	6	3
1:A:194:ARG:O	1:A:196:ASP:N	0.57	2.37	19	2
1:A:132:PHE:CG	1:A:132:PHE:O	0.56	2.58	12	2
1:A:145:ALA:O	1:A:146:CYS:CB	0.55	2.55	11	12

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:132:PHE:CD2	1:A:132:PHE:O	0.54	2.59	7	2
1:A:203:ASP:O	1:A:205:SER:N	0.54	2.41	19	2
1:A:156:SER:HA	1:A:159:TRP:CD2	0.52	2.39	19	15
1:A:134:CYS:N	1:A:138:THR:O	0.52	2.42	18	7
1:A:196:ASP:OD1	1:A:196:ASP:N	0.52	2.42	20	1
1:A:184:LEU:HD13	1:A:205:SER:HB2	0.51	1.79	11	2
1:A:155:GLY:O	1:A:158:GLU:N	0.51	2.43	7	6
1:A:196:ASP:OD1	1:A:207:GLU:OE2	0.50	2.30	20	1
1:A:155:GLY:O	1:A:157:ASP:N	0.49	2.45	13	10
1:A:184:LEU:C	1:A:186:GLY:H	0.49	2.11	19	1
1:A:136:SER:O	1:A:137:SER:CB	0.49	2.61	11	4
1:A:193:TRP:O	1:A:196:ASP:OD1	0.49	2.31	20	1
1:A:140:ILE:HD13	1:A:152:CYS:SG	0.49	2.48	12	7
1:A:132:PHE:CE2	1:A:133:GLN:O	0.48	2.67	8	9
1:A:132:PHE:N	1:A:140:ILE:O	0.47	2.48	4	3
1:A:129:PRO:O	1:A:130:ALA:HB3	0.47	2.09	19	2
1:A:156:SER:O	1:A:162:ARG:HB2	0.47	2.10	17	7
1:A:143:LEU:HD12	1:A:143:LEU:H	0.47	1.69	4	5
1:A:184:LEU:O	1:A:186:GLY:N	0.47	2.48	19	2
1:A:181:PHE:CE1	1:A:194:ARG:NE	0.46	2.83	10	7
1:A:139:CYS:O	1:A:140:ILE:HG23	0.46	2.10	19	3
1:A:132:PHE:CD2	1:A:145:ALA:CB	0.46	2.98	17	4
1:A:143:LEU:H	1:A:143:LEU:HD12	0.46	1.71	14	3
1:A:193:TRP:CD1	1:A:193:TRP:N	0.45	2.84	16	10
1:A:193:TRP:N	1:A:193:TRP:CD1	0.45	2.84	17	10
1:A:132:PHE:CB	1:A:145:ALA:HB2	0.45	2.42	9	4
1:A:183:CYS:SG	1:A:184:LEU:N	0.45	2.89	20	1
1:A:139:CYS:O	1:A:140:ILE:CG2	0.45	2.64	11	13
1:A:184:LEU:N	1:A:205:SER:OG	0.45	2.50	5	2
1:A:179:PHE:CD1	1:A:179:PHE:N	0.45	2.85	20	2
1:A:132:PHE:CG	1:A:145:ALA:CB	0.44	2.99	17	2
1:A:193:TRP:CE3	1:A:200:ASP:OD2	0.44	2.71	11	2
1:A:146:CYS:N	1:A:157:ASP:OD1	0.44	2.51	7	1
1:A:194:ARG:C	1:A:196:ASP:N	0.43	2.70	20	2
1:A:194:ARG:C	1:A:196:ASP:H	0.43	2.17	20	1
1:A:184:LEU:HD22	1:A:205:SER:HB3	0.43	1.91	5	2
1:A:143:LEU:O	1:A:145:ALA:N	0.43	2.52	1	8
1:A:132:PHE:O	1:A:133:GLN:C	0.43	2.57	7	1
1:A:140:ILE:HG21	1:A:151:ASP:O	0.42	2.13	7	1
1:A:145:ALA:O	1:A:146:CYS:C	0.42	2.57	7	1
1:A:158:GLU:O	1:A:159:TRP:C	0.42	2.57	18	7

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:156:SER:O	1:A:162:ARG:CB	0.41	2.69	18	5
1:A:185:SER:HB3	1:A:201:CYS:SG	0.41	2.55	6	2
1:A:203:ASP:O	1:A:204:LYS:C	0.41	2.58	20	2
1:A:138:THR:HG22	1:A:139:CYS:N	0.41	2.31	14	4
1:A:203:ASP:O	1:A:204:LYS:CB	0.41	2.68	14	2
1:A:147:ASP:O	1:A:148:ASN:CB	0.41	2.68	19	1
1:A:143:LEU:N	1:A:143:LEU:HD12	0.41	2.30	4	1
1:A:143:LEU:HD12	1:A:143:LEU:N	0.41	2.30	12	2
1:A:147:ASP:O	1:A:148:ASN:HB2	0.41	2.16	16	1
1:A:132:PHE:O	1:A:132:PHE:CD2	0.41	2.73	19	1
1:A:138:THR:CG2	1:A:139:CYS:N	0.41	2.84	14	1
1:A:142:GLN:O	1:A:142:GLN:HG2	0.41	2.16	12	1
1:A:156:SER:O	1:A:159:TRP:HB2	0.41	2.16	19	1
1:A:155:GLY:O	1:A:159:TRP:CD1	0.41	2.74	20	1
1:A:132:PHE:CZ	1:A:133:GLN:O	0.40	2.74	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	70/94 (74%)	52±2 (75±3%)	11±2 (16±3%)	6±2 (9±3%)	1	11
All	All	1400/1880 (74%)	1047 (75%)	226 (16%)	127 (9%)	1	11

All 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	148	ASN	20
1	A	156	SER	12
1	A	145	ALA	12
1	A	152	CYS	11
1	A	163	CYS	10
1	A	146	CYS	9
1	A	144	TRP	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	128	GLY	7
1	A	129	PRO	6
1	A	184	LEU	6
1	A	137	SER	5
1	A	180	GLU	3
1	A	195	CYS	2
1	A	185	SER	2
1	A	177	SER	2
1	A	206	ASP	2
1	A	127	CYS	2
1	A	204	LYS	2
1	A	154	ASP	2
1	A	208	GLU	1
1	A	133	GLN	1
1	A	134	CYS	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/81 (77%)	51±3 (83±4%)	11±3 (17±4%)	5	40
All	All	1240/1620 (77%)	1029 (83%)	211 (17%)	5	40

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	191	SER	20
1	A	205	SER	18
1	A	146	CYS	15
1	A	156	SER	14
1	A	127	CYS	13
1	A	151	ASP	11
1	A	200	ASP	10
1	A	147	ASP	9
1	A	183	CYS	9
1	A	137	SER	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	148	ASN	9
1	A	153	GLU	8
1	A	131	SER	6
1	A	187	GLU	6
1	A	194	ARG	6
1	A	163	CYS	6
1	A	161	GLN	5
1	A	154	ASP	5
1	A	203	ASP	4
1	A	196	ASP	4
1	A	149	ASP	3
1	A	142	GLN	3
1	A	209	ASN	2
1	A	185	SER	2
1	A	162	ARG	2
1	A	206	ASP	2
1	A	133	GLN	2
1	A	195	CYS	1
1	A	184	LEU	1
1	A	141	PRO	1
1	A	202	LYS	1
1	A	201	CYS	1
1	A	134	CYS	1
1	A	138	THR	1
1	A	157	ASP	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 91% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	997
Number of shifts mapped to atoms	997
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	94	-0.26 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	86	0.82 ± 0.18	Should be applied
$^{13}\text{C}'$	76	-0.36 ± 0.28	None needed (< 0.5 ppm)
^{15}N	82	0.78 ± 0.74	None needed (imprecise)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 713 atoms were assigned a chemical shift out of a possible 782. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	322/340 (95%)	130/135 (96%)	129/140 (92%)	63/65 (97%)
Sidechain	320/363 (88%)	208/219 (95%)	106/130 (82%)	6/14 (43%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	71/79 (90%)	37/41 (90%)	31/31 (100%)	3/7 (43%)
Overall	713/782 (91%)	375/395 (95%)	266/301 (88%)	72/86 (84%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 91%, i.e. 937 atoms were assigned a chemical shift out of a possible 1034. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	425/456 (93%)	173/181 (96%)	170/188 (90%)	82/87 (94%)
Sidechain	424/482 (88%)	271/289 (94%)	146/175 (83%)	7/18 (39%)
Aromatic	88/96 (92%)	46/50 (92%)	39/39 (100%)	3/7 (43%)
Overall	937/1034 (91%)	490/520 (94%)	355/402 (88%)	92/112 (82%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	156	SER	HA	1.68	6.53 – 2.43	-6.8
1	A	156	SER	HB3	2.20	5.25 – 2.45	-5.9

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

