



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 6, 2020 – 11:51 pm BST

PDB ID : 3LG2  
Title : A Ykr043C/ fructose-1,6-bisphosphate product complex following ligand soaking  
Authors : Singer, A.; Xu, X.; Cui, H.; Dong, A.; Edwards, A.M.; Joachimiak, A.; Yakunin, A.F.; Savchenko, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2010-01-19  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>%</span> <span>68%</span> <span>22%</span> <span>• 8%</span> </div> </div>
1	B	292	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>3%</span> <span>68%</span> <span>21%</span> <span>• 8%</span> </div> </div>
1	C	292	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>5%</span> <span>69%</span> <span>23%</span> <span>• 8%</span> </div> </div>
1	D	292	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span></span> <span>71%</span> <span>19%</span> <span>• 8%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	300	-	-	X	-
2	PO4	B	302	-	-	X	-
2	PO4	C	272	-	-	X	-
2	PO4	C	302	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein YKR043C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	Se	0	0	0
			2172	1362	391	412	4	3			
1	B	269	Total	C	N	O	S	Se	0	0	0
			2172	1362	391	412	4	3			
1	C	269	Total	C	N	O	S	Se	0	0	0
			2172	1362	391	412	4	3			
1	D	269	Total	C	N	O	S	Se	0	0	0
			2172	1362	391	412	4	3			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MSE	-	expression tag	UNP P36136
A	-19	GLY	-	expression tag	UNP P36136
A	-18	SER	-	expression tag	UNP P36136
A	-17	SER	-	expression tag	UNP P36136
A	-16	HIS	-	expression tag	UNP P36136
A	-15	HIS	-	expression tag	UNP P36136
A	-14	HIS	-	expression tag	UNP P36136
A	-13	HIS	-	expression tag	UNP P36136
A	-12	HIS	-	expression tag	UNP P36136
A	-11	HIS	-	expression tag	UNP P36136
A	-10	SER	-	expression tag	UNP P36136
A	-9	SER	-	expression tag	UNP P36136
A	-8	GLY	-	expression tag	UNP P36136
A	-7	ARG	-	expression tag	UNP P36136
A	-6	GLU	-	expression tag	UNP P36136
A	-5	ASN	-	expression tag	UNP P36136
A	-4	LEU	-	expression tag	UNP P36136
A	-3	TYR	-	expression tag	UNP P36136
A	-2	PHE	-	expression tag	UNP P36136
A	-1	GLN	-	expression tag	UNP P36136
A	0	GLY	-	expression tag	UNP P36136

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MSE	-	expression tag	UNP P36136
B	-19	GLY	-	expression tag	UNP P36136
B	-18	SER	-	expression tag	UNP P36136
B	-17	SER	-	expression tag	UNP P36136
B	-16	HIS	-	expression tag	UNP P36136
B	-15	HIS	-	expression tag	UNP P36136
B	-14	HIS	-	expression tag	UNP P36136
B	-13	HIS	-	expression tag	UNP P36136
B	-12	HIS	-	expression tag	UNP P36136
B	-11	HIS	-	expression tag	UNP P36136
B	-10	SER	-	expression tag	UNP P36136
B	-9	SER	-	expression tag	UNP P36136
B	-8	GLY	-	expression tag	UNP P36136
B	-7	ARG	-	expression tag	UNP P36136
B	-6	GLU	-	expression tag	UNP P36136
B	-5	ASN	-	expression tag	UNP P36136
B	-4	LEU	-	expression tag	UNP P36136
B	-3	TYR	-	expression tag	UNP P36136
B	-2	PHE	-	expression tag	UNP P36136
B	-1	GLN	-	expression tag	UNP P36136
B	0	GLY	-	expression tag	UNP P36136
C	-20	MSE	-	expression tag	UNP P36136
C	-19	GLY	-	expression tag	UNP P36136
C	-18	SER	-	expression tag	UNP P36136
C	-17	SER	-	expression tag	UNP P36136
C	-16	HIS	-	expression tag	UNP P36136
C	-15	HIS	-	expression tag	UNP P36136
C	-14	HIS	-	expression tag	UNP P36136
C	-13	HIS	-	expression tag	UNP P36136
C	-12	HIS	-	expression tag	UNP P36136
C	-11	HIS	-	expression tag	UNP P36136
C	-10	SER	-	expression tag	UNP P36136
C	-9	SER	-	expression tag	UNP P36136
C	-8	GLY	-	expression tag	UNP P36136
C	-7	ARG	-	expression tag	UNP P36136
C	-6	GLU	-	expression tag	UNP P36136
C	-5	ASN	-	expression tag	UNP P36136
C	-4	LEU	-	expression tag	UNP P36136
C	-3	TYR	-	expression tag	UNP P36136
C	-2	PHE	-	expression tag	UNP P36136
C	-1	GLN	-	expression tag	UNP P36136
C	0	GLY	-	expression tag	UNP P36136

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MSE	-	expression tag	UNP P36136
D	-19	GLY	-	expression tag	UNP P36136
D	-18	SER	-	expression tag	UNP P36136
D	-17	SER	-	expression tag	UNP P36136
D	-16	HIS	-	expression tag	UNP P36136
D	-15	HIS	-	expression tag	UNP P36136
D	-14	HIS	-	expression tag	UNP P36136
D	-13	HIS	-	expression tag	UNP P36136
D	-12	HIS	-	expression tag	UNP P36136
D	-11	HIS	-	expression tag	UNP P36136
D	-10	SER	-	expression tag	UNP P36136
D	-9	SER	-	expression tag	UNP P36136
D	-8	GLY	-	expression tag	UNP P36136
D	-7	ARG	-	expression tag	UNP P36136
D	-6	GLU	-	expression tag	UNP P36136
D	-5	ASN	-	expression tag	UNP P36136
D	-4	LEU	-	expression tag	UNP P36136
D	-3	TYR	-	expression tag	UNP P36136
D	-2	PHE	-	expression tag	UNP P36136
D	-1	GLN	-	expression tag	UNP P36136
D	0	GLY	-	expression tag	UNP P36136

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		
3	B	2	Total	O	0	0
			2	2		
3	C	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Uncharacterized protein YKR043C



#### • Molecule 1: Uncharacterized protein YKR043C



#### • Molecule 1: Uncharacterized protein YKR043C







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.30 Å 83.39 Å 104.69 Å 90.00° 104.19° 90.00°	Depositor
Resolution (Å)	43.35 – 2.60 43.35 – 2.60	Depositor EDS
% Data completeness (in resolution range)	91.2 (43.35-2.60) 91.2 (43.35-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.45 (at 2.61 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.220 , 0.270 0.242 , 0.289	Depositor DCC
$R_{free}$ test set	2008 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.4	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 30.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8732	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 62.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1303e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	0/2218	0.78	1/2999 (0.0%)
1	B	0.71	0/2218	0.80	0/2999
1	C	0.81	1/2218 (0.0%)	0.84	0/2999
1	D	0.81	1/2218 (0.0%)	0.87	1/2999 (0.0%)
All	All	0.77	2/8872 (0.0%)	0.82	2/11996 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	61	TYR	CD1-CE1	5.56	1.47	1.39
1	C	12	ARG	CB-CG	5.32	1.66	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	147	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	12	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2172	0	2114	45	0
1	B	2172	0	2114	55	0
1	C	2172	0	2114	48	0
1	D	2172	0	2114	44	0
2	A	5	0	0	0	0
2	B	15	0	0	7	0
2	C	15	0	0	4	0
2	D	5	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
All	All	8732	0	8456	186	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:247:ILE:HD12	1:C:247:ILE:O	1.47	1.15
1:C:50:ASN:HB2	1:C:52:GLN:NE2	1.63	1.13
1:D:108:MSE:HE2	1:D:113:ILE:HG13	1.28	1.10
1:D:108:MSE:HE1	1:D:113:ILE:HA	1.41	1.01
1:D:199:ILE:O	1:D:203:GLN:HG2	1.63	0.99

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/292 (91%)	257 (96%)	10 (4%)	0	100	100
1	B	267/292 (91%)	257 (96%)	8 (3%)	2 (1%)	22	43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	267/292 (91%)	255 (96%)	11 (4%)	1 (0%)	34	57
1	D	267/292 (91%)	258 (97%)	7 (3%)	2 (1%)	22	43
All	All	1068/1168 (91%)	1027 (96%)	36 (3%)	5 (0%)	29	52

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	206	LYS
1	B	175	ALA
1	B	241	SER
1	D	175	ALA
1	C	241	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/250 (94%)	218 (93%)	17 (7%)	14	29
1	B	235/250 (94%)	216 (92%)	19 (8%)	11	23
1	C	235/250 (94%)	219 (93%)	16 (7%)	16	32
1	D	235/250 (94%)	220 (94%)	15 (6%)	17	35
All	All	940/1000 (94%)	873 (93%)	67 (7%)	14	29

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	188	PHE
1	C	10	ILE
1	D	164	SER
1	B	194	LYS
1	B	231	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	52	GLN
1	C	58	ASN
1	D	137	ASN
1	C	23	GLN
1	D	142	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	B	272	-	4,4,4	0.84	0	6,6,6	1.36	2 (33%)
2	PO4	D	302	-	4,4,4	0.97	0	6,6,6	1.88	2 (33%)
2	PO4	C	300	-	4,4,4	0.68	0	6,6,6	2.51	3 (50%)
2	PO4	B	302	-	4,4,4	0.64	0	6,6,6	1.49	2 (33%)
2	PO4	C	302	-	4,4,4	1.53	0	6,6,6	0.66	0
2	PO4	C	272	-	4,4,4	1.46	1 (25%)	6,6,6	2.01	2 (33%)
2	PO4	A	302	-	4,4,4	0.85	0	6,6,6	1.30	1 (16%)
2	PO4	B	300	-	4,4,4	0.88	0	6,6,6	1.86	2 (33%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	272	PO4	P-O2	-2.25	1.47	1.54

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	300	PO4	O4-P-O3	4.23	121.53	107.97
2	C	272	PO4	O2-P-O1	-3.87	96.74	110.89
2	C	300	PO4	O4-P-O1	-3.08	99.61	110.89
2	D	302	PO4	O3-P-O1	-2.97	100.04	110.89
2	B	300	PO4	O2-P-O1	-2.74	100.88	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	PO4	2	0
2	C	302	PO4	2	0
2	C	272	PO4	2	0
2	B	300	PO4	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/292 (91%)	-0.11	3 (1%) 80 78	14, 24, 41, 56	0
1	B	266/292 (91%)	0.02	9 (3%) 45 38	14, 24, 41, 56	0
1	C	266/292 (91%)	0.02	14 (5%) 26 20	14, 24, 41, 56	0
1	D	266/292 (91%)	-0.07	1 (0%) 92 91	14, 24, 41, 55	0
All	All	1064/1168 (91%)	-0.03	27 (2%) 57 51	14, 24, 43, 56	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	125	GLU	3.9
1	A	197	GLU	3.3
1	C	200	GLU	3.0
1	B	198	THR	2.8
1	C	203	GLN	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	B	300	5/5	0.89	0.19	32,37,42,43	0
2	PO4	B	302	5/5	0.96	0.20	16,24,25,27	0
2	PO4	D	302	5/5	0.96	0.16	4,9,15,18	0
2	PO4	B	272	5/5	0.97	0.15	27,27,29,29	0
2	PO4	C	272	5/5	0.97	0.12	22,22,22,22	0
2	PO4	A	302	5/5	0.97	0.18	15,22,23,24	0
2	PO4	C	300	5/5	0.97	0.15	14,15,19,21	0
2	PO4	C	302	5/5	0.99	0.13	2,5,9,9	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.