



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 09:18 pm BST

PDB ID : 4LLU
Title : Structure of Pertuzumab Fab with light chain Clambda at 2.16Å
Authors : Pustilnik, A.; Lewis, S.M.; Wu, X.; Sereno, A.; Huang, F.; Guntas, G.; Leaver-Fay, A.; Smith, E.M.; Ho, C.; Hansen-Estruch, C.; Chamberlain, A.K.; Truhlar, S.M.; Kuhlman, B.; Demarest, S.J.; Atwell, S.
Deposited on : 2013-07-09
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 87%, yellow 87%, yellow 96%, grey 96%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 87% 9% .. </div> </div>
1	C	227	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 84%, yellow 84%, yellow 96%, grey 96%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 84% 12% .. </div> </div>
2	B	212	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, green 3%, green 87%, yellow 87%, yellow 99%, grey 99%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 87% 12% . </div> </div>
2	D	212	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 17%, green 17%, green 84%, yellow 84%, yellow 95%, grey 95%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 17% 84% 11% .. </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PERTUZUMAB FAB Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1653	1045	278	324	6			
1	C	220	Total	C	N	O	S	0	2	0
			1653	1044	277	324	8			

- Molecule 2 is a protein called Light chain CLAMBDA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	210	Total	C	N	O	S	0	0	0
			1596	1012	258	321	5			
2	D	206	Total	C	N	O	S	0	0	0
			1540	974	247	314	5			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	76	Total O 76 76	0	0
5	B	53	Total O 53 53	0	0
5	C	71	Total O 71 71	0	0
5	D	45	Total O 45 45	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: PERTUZUMAB FAB Heavy chain





- Molecule 1: PERTUZUMAB FAB Heavy chain



- Molecule 2: Light chain CLAMBDA



- Molecule 2: Light chain CLAMBDA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	210.02Å 69.74Å 70.83Å 90.00° 98.75° 90.00°	Depositor
Resolution (Å)	19.90 – 2.16 19.90 – 2.16	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.90-2.16) 95.6 (19.90-2.16)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.17Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.222 , 0.273 0.220 , 0.267	Depositor DCC
R_{free} test set	2654 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6706	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/1694	0.64	0/2308
1	C	0.56	0/1702	0.63	0/2320
2	B	0.54	0/1638	0.64	0/2237
2	D	0.52	0/1581	0.59	0/2163
All	All	0.55	0/6615	0.62	0/9028

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1653	0	1614	11	0
1	C	1653	0	1603	18	0
2	B	1596	0	1542	15	0
2	D	1540	0	1439	15	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	B	4	0	3	1	0
5	A	76	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	53	0	0	3	0
5	C	71	0	0	1	0
5	D	45	0	0	0	0
All	All	6706	0	6201	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:79:GLN:HG3	2:D:80:PRO:HD2	1.64	0.78
1:C:83:ARG:HD3	1:C:85:GLU:HG2	1.71	0.73
2:D:121:PRO:HD3	2:D:133:LEU:CD2	2.20	0.71
1:C:214:LYS:NZ	5:C:410:HOH:O	2.25	0.68
1:A:52:ASN:HD21	1:A:53:ASN:HD22	1.43	0.66
2:B:193:SER:OG	2:B:206:THR:HG22	1.98	0.62
1:A:34:MET:HB3	1:A:78:LEU:HD22	1.81	0.61
2:B:1:ASP:OD1	2:B:2:ILE:HD12	2.01	0.61
2:D:111:LYS:H	2:D:111:LYS:HE3	1.67	0.60
2:B:83:PHE:CE1	2:B:106:ILE:HD12	2.38	0.59
2:B:121:PRO:HD3	2:B:133:LEU:CD2	2.34	0.58
2:B:111:LYS:HG2	2:B:142:PRO:HD3	1.85	0.58
2:B:150:LYS:HE2	2:B:195:GLN:OE1	2.04	0.57
1:A:126:PRO:HB2	1:A:187:SER:HA	1.87	0.56
2:D:121:PRO:HD3	2:D:133:LEU:HD23	1.88	0.55
2:B:42:LYS:HD3	5:B:421:HOH:O	2.08	0.53
2:D:121:PRO:HD3	2:D:133:LEU:HD21	1.91	0.53
1:C:159:LEU:HD21	1:C:182:VAL:HG21	1.89	0.53
1:C:35:ASP:OD1	1:C:95:ASN:ND2	2.42	0.52
1:C:61:GLN:HG2	2:D:94:TYR:OH	2.11	0.51
1:C:49:ALA:HB1	1:C:69:LEU:HD12	1.92	0.50
2:D:33:VAL:HA	2:D:89:GLN:O	2.13	0.49
2:B:195:GLN:NE2	5:B:435:HOH:O	2.39	0.49
2:B:109:GLN:HB3	2:B:110:PRO:HD2	1.94	0.48
2:D:111:LYS:N	2:D:111:LYS:HE3	2.28	0.48
2:B:198:HIS:HD2	4:B:301:ACT:O	1.97	0.47
1:C:61:GLN:C	1:C:63:PHE:H	2.18	0.47
1:A:181:VAL:HG21	2:B:136:LEU:HD23	1.95	0.47
2:D:145:VAL:HG12	2:D:198:HIS:HB2	1.96	0.47
2:D:35:TRP:CZ3	2:D:88:CYS:HB3	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:16:GLY:H	2:D:78:LEU:HB3	1.80	0.46
2:D:80:PRO:O	2:D:83:PHE:CD1	2.68	0.46
2:D:18:ARG:HA	2:D:75:ILE:O	2.16	0.46
1:A:61:GLN:NE2	1:A:61:GLN:HA	2.31	0.46
1:A:11:LEU:HD11	1:A:112:SER:HB3	1.99	0.45
1:A:199:ASN:HD22	1:A:200:HIS:N	2.14	0.45
2:B:121:PRO:HD3	2:B:133:LEU:HD23	2.00	0.44
1:C:192:GLN:HG2	1:C:194:TYR:CZ	2.52	0.44
2:B:206:THR:HG23	5:B:432:HOH:O	2.17	0.44
1:C:40:ALA:HB3	1:C:43:LYS:HB2	1.99	0.43
1:C:150:VAL:HG22	1:C:178:LEU:HD21	2.01	0.42
1:C:1:GLU:HG2	1:C:2:VAL:H	1.84	0.42
2:B:142:PRO:O	2:B:198:HIS:HE1	2.03	0.42
1:C:52:ASN:HA	1:C:52(A):PRO:HD3	1.91	0.42
2:D:95:PRO:HG2	2:D:97:THR:HG23	2.01	0.42
1:A:181:VAL:CG2	2:B:136:LEU:HD23	2.49	0.41
1:A:61:GLN:HE22	1:A:64:LYS:HG3	1.85	0.41
1:C:11:LEU:HD11	1:C:112:SER:HB3	2.01	0.41
1:C:127:SER:HB3	1:C:214:LYS:HD2	2.01	0.41
1:A:2:VAL:HG13	1:A:27:PHE:CD1	2.55	0.41
1:A:126:PRO:HA	1:A:137:ALA:O	2.21	0.41
1:C:39:GLN:HB2	1:C:45:LEU:HD23	2.02	0.41
1:C:200:HIS:CD2	1:C:202:PRO:HD2	2.56	0.40
1:C:83:ARG:HG2	1:C:85:GLU:H	1.86	0.40
2:D:58:VAL:HA	2:D:59:PRO:HD3	1.87	0.40
1:C:61:GLN:C	1:C:63:PHE:N	2.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/227 (96%)	213 (98%)	5 (2%)	0	100	100
1	C	220/227 (97%)	211 (96%)	9 (4%)	0	100	100
2	B	208/212 (98%)	201 (97%)	7 (3%)	0	100	100
2	D	202/212 (95%)	189 (94%)	12 (6%)	1 (0%)	29	22
All	All	848/878 (97%)	814 (96%)	33 (4%)	1 (0%)	51	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	15	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/192 (96%)	176 (95%)	9 (5%)	25	21
1	C	186/192 (97%)	182 (98%)	4 (2%)	52	55
2	B	177/182 (97%)	167 (94%)	10 (6%)	21	16
2	D	166/182 (91%)	158 (95%)	8 (5%)	25	22
All	All	714/748 (96%)	683 (96%)	31 (4%)	28	27

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	50	ASP
1	A	62	ARG
1	A	64	LYS
1	A	113	SER
1	A	161	SER
1	A	172	SER
1	A	197	ASN
1	A	199	ASN
2	B	3	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	9	SER
2	B	54	ARG
2	B	70	ASP
2	B	97	THR
2	B	105	GLU
2	B	107	LYS
2	B	136	LEU
2	B	139	ASP
2	B	156	VAL
1	C	13	GLN
1	C	69	LEU
1	C	83	ARG
1	C	164	HIS
2	D	2	ILE
2	D	3	GLN
2	D	79	GLN
2	D	97	THR
2	D	111	LYS
2	D	136	LEU
2	D	169	SER
2	D	176	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	61	GLN
1	A	76	ASN
1	A	199	ASN
1	A	204	ASN
2	B	3	GLN
2	B	198	HIS
1	C	76	ASN
1	C	204	ASN
2	D	79	GLN
2	D	189	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	B	301	-	1,3,3	1.95	0	0,3,3	0.00	-
3	SO4	A	301	-	4,4,4	0.15	0	6,6,6	0.09	0
3	SO4	D	301	-	4,4,4	0.12	0	6,6,6	0.16	0
3	SO4	C	301	-	4,4,4	0.17	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/227 (96%)	0.01	2 (0%) 84 88	4, 20, 46, 56	0
1	C	220/227 (96%)	0.27	8 (3%) 42 51	5, 26, 54, 65	0
2	B	210/212 (99%)	0.13	6 (2%) 51 61	4, 24, 40, 50	0
2	D	206/212 (97%)	0.77	36 (17%) 1 1	5, 33, 80, 87	0
All	All	856/878 (97%)	0.29	52 (6%) 21 28	4, 24, 60, 87	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	77	SER	5.3
2	D	78	LEU	4.9
2	D	8	PRO	4.7
2	B	169	SER	4.5
2	D	104	VAL	4.4
2	D	15	VAL	4.1
1	C	54	SER	4.1
2	D	11	LEU	4.0
2	D	73	LEU	3.8
1	C	61	GLN	3.8
2	D	80	PRO	3.7
2	D	16	GLY	3.5
1	C	56	GLY	3.5
2	D	106	ILE	3.4
1	C	42	GLY	3.4
2	D	10	SER	3.4
2	D	65	SER	3.4
2	D	20	THR	3.3
2	D	76	SER	3.3
2	B	56	THR	3.2
2	D	53	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	65	GLY	3.0
2	D	9	SER	3.0
2	D	74	THR	3.0
2	B	210	THR	3.0
2	D	23	CYS	2.9
2	D	169	SER	2.8
2	B	1	ASP	2.7
2	D	60	SER	2.6
2	D	35	TRP	2.6
2	D	22	THR	2.5
2	B	106	ILE	2.5
2	D	86	TYR	2.4
2	D	79	GLN	2.4
1	C	78	LEU	2.4
2	D	21	ILE	2.4
1	C	41	PRO	2.4
1	C	26	GLY	2.4
2	D	17	ASP	2.3
1	C	1	GLU	2.3
2	D	33	VAL	2.3
2	D	99	GLY	2.3
2	D	57	GLY	2.3
2	D	49	TYR	2.2
2	D	12	SER	2.2
2	D	40	PRO	2.2
2	D	70	ASP	2.2
1	A	64	LYS	2.1
2	D	4	MET	2.1
2	B	35	TRP	2.1
2	D	63	SER	2.0
2	D	100	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ACT	B	301	4/4	0.91	0.17	40,41,41,42	0
3	SO4	A	301	5/5	0.94	0.32	59,64,65,66	0
3	SO4	C	301	5/5	0.94	0.17	34,40,46,48	0
3	SO4	D	301	5/5	0.96	0.23	60,61,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.