



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:21 am BST

PDB ID : 4LO4
Title : Apo HA-70
Authors : Lee, K.; Gu, S.; Jin, L.; Le, T.T.; Cheng, L.W.; Strotmeier, J.; Kruel, A.M.;
Yao, G.; Perry, K.; Rummel, A.; Jin, R.
Deposited on : 2013-07-12
Resolution : 2.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>•</div> <div>11%</div> </div> </div>
2	B	433	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA-70.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	0	0	0
			1389	898	224	265	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP Q8KHU9
A	1	SER	-	EXPRESSION TAG	UNP Q8KHU9

- Molecule 2 is a protein called HA-70.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	421	Total	C	N	O	S	18	0	0
			3371	2124	565	677	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	627	PRO	-	EXPRESSION TAG	UNP Q8KHU9
B	628	GLY	-	EXPRESSION TAG	UNP Q8KHU9
B	629	SER	-	EXPRESSION TAG	UNP Q8KHU9
B	630	ALA	-	EXPRESSION TAG	UNP Q8KHU9
B	631	TRP	-	EXPRESSION TAG	UNP Q8KHU9
B	632	SER	-	EXPRESSION TAG	UNP Q8KHU9
B	633	HIS	-	EXPRESSION TAG	UNP Q8KHU9
B	634	PRO	-	EXPRESSION TAG	UNP Q8KHU9
B	635	GLN	-	EXPRESSION TAG	UNP Q8KHU9
B	636	PHE	-	EXPRESSION TAG	UNP Q8KHU9
B	637	GLU	-	EXPRESSION TAG	UNP Q8KHU9
B	638	LYS	-	EXPRESSION TAG	UNP Q8KHU9

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total 7	O 7	0	0
4	B	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HA-70



• Molecule 2: HA-70



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	261.75Å 261.75Å 261.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.89 – 2.87 44.89 – 2.87	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.89-2.87) 99.7 (44.89-2.87)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.86Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.186 , 0.206 0.184 , 0.201	Depositor DCC
R_{free} test set	1757 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	77.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4787	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1421	0.50	0/1929
2	B	0.29	0/3434	0.49	0/4672
All	All	0.29	0/4855	0.50	0/6601

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1389	0	1364	10	0
2	B	3371	0	3281	30	0
3	B	1	0	0	0	0
4	A	7	0	0	1	0
4	B	19	0	0	1	0
All	All	4787	0	4645	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:ILE:HG13	1:A:177:LEU:HD13	1.80	0.64
2:B:529:GLN:HB3	4:B:810:HOH:O	1.97	0.63
1:A:165:LYS:NZ	4:A:203:HOH:O	2.28	0.62
2:B:473:LEU:H	2:B:509:ASN:HD21	1.48	0.61
2:B:473:LEU:H	2:B:509:ASN:ND2	2.01	0.58
2:B:557:LEU:HD13	2:B:606:LEU:HD22	1.88	0.54
2:B:227:ARG:HD3	2:B:232:ASP:OD2	2.08	0.53
2:B:219:ILE:HG12	2:B:372:VAL:HG22	1.91	0.53
2:B:302:ASP:OD1	2:B:302:ASP:N	2.42	0.52
1:A:31:VAL:HG21	1:A:91:VAL:HG21	1.91	0.51
1:A:43:GLN:HG2	1:A:46:ILE:HD13	1.92	0.51
2:B:568:LEU:HD13	2:B:583:ILE:HD11	1.91	0.51
2:B:264:ASN:OD1	2:B:266:THR:HG22	2.12	0.50
2:B:503:ASP:HB3	2:B:580:LEU:HB2	1.94	0.49
2:B:552:THR:HG23	2:B:597:ARG:HH21	1.77	0.49
1:A:31:VAL:HG13	1:A:186:ILE:HD11	1.95	0.48
1:A:52:ILE:HG13	1:A:111:ALA:HB2	1.96	0.47
1:A:112:PRO:HG2	1:A:115:LYS:HB2	1.96	0.47
2:B:625:THR:O	2:B:626:ASN:HB2	2.15	0.47
2:B:508:MET:HB2	2:B:509:ASN:ND2	2.30	0.46
2:B:276:ARG:NH1	2:B:279:ASP:OD2	2.47	0.46
2:B:348:PRO:HA	2:B:349:GLU:HA	1.58	0.46
2:B:554:ASN:HD22	2:B:597:ARG:HG2	1.80	0.46
2:B:276:ARG:HA	2:B:276:ARG:HD3	1.63	0.45
1:A:89:GLN:HG2	1:A:94:ASN:HA	1.98	0.45
1:A:172:ASN:OD1	2:B:478:LYS:NZ	2.47	0.45
2:B:320:ASP:HB3	2:B:328:ASN:HB2	1.99	0.44
1:A:45:GLN:NE2	2:B:369:GLU:OE2	2.37	0.43
2:B:376:ILE:HG23	2:B:407:TYR:CE2	2.53	0.43
2:B:536:TYR:CE1	2:B:609:GLU:HB2	2.54	0.43
2:B:554:ASN:HB3	2:B:625:THR:HG23	2.00	0.43
2:B:472:VAL:HA	2:B:509:ASN:HD22	1.84	0.43
2:B:562:ASN:OD1	2:B:614:ASN:N	2.50	0.42
2:B:468:SER:HA	2:B:512:VAL:O	2.20	0.42
2:B:508:MET:HA	2:B:509:ASN:HA	1.74	0.41
2:B:283:VAL:HG23	2:B:286:PHE:CZ	2.55	0.41
2:B:214:ASN:HB3	2:B:322:GLN:NE2	2.35	0.41
2:B:345:ALA:HA	2:B:346:PRO:HD3	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/190 (88%)	159 (95%)	9 (5%)	0	100	100
2	B	419/433 (97%)	402 (96%)	17 (4%)	0	100	100
All	All	587/623 (94%)	561 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/173 (89%)	146 (95%)	8 (5%)	23	53
2	B	383/393 (98%)	364 (95%)	19 (5%)	24	54
All	All	537/566 (95%)	510 (95%)	27 (5%)	24	54

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	31	VAL
1	A	43	GLN
1	A	47	LEU
1	A	65	VAL
1	A	93	THR
1	A	177	LEU
1	A	184	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	240	GLU
2	B	276	ARG
2	B	284	GLN
2	B	302	ASP
2	B	331	GLU
2	B	337	LEU
2	B	359	GLU
2	B	370	GLU
2	B	451	ASP
2	B	470	LEU
2	B	484	VAL
2	B	487	THR
2	B	508	MET
2	B	546	ASN
2	B	567	THR
2	B	568	LEU
2	B	587	LEU
2	B	589	LEU
2	B	625	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	GLN
1	A	180	GLN
2	B	287	ASN
2	B	423	GLN
2	B	509	ASN
2	B	626	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	170/190 (89%)	-0.11	4 (2%) 59 57	58, 86, 131, 171	0
2	B	421/433 (97%)	-0.29	6 (1%) 75 75	57, 79, 126, 214	7 (1%)
All	All	591/623 (94%)	-0.24	10 (1%) 70 70	57, 80, 128, 214	7 (1%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	399	GLY	8.9
2	B	398	PRO	8.8
2	B	400	ILE	8.4
2	B	401	VAL	6.9
1	A	189	THR	4.6
2	B	206	THR	4.1
1	A	23	ILE	3.5
1	A	181	PHE	2.4
2	B	402	ASN	2.2
1	A	30	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CL	B	701	1/1	0.96	0.10	69,69,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.