



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 06:39 pm BST

PDB ID : 4LO8
Title : HA70(D3)-HA17
Authors : Lee, K.; Gu, S.; Jin, L.; Le, T.T.; Cheng, L.W.; Strotmeier, J.; Kruel, A.M.;
Yao, G.; Perry, K.; Rummel, A.; Jin, R.
Deposited on : 2013-07-12
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>8%</div> </div> </div>
1	C	254	<div> <div>11%</div> <div> <div></div> <div>73%</div> <div>16%</div> <div>8%</div> </div> </div>
1	E	254	<div> <div>6%</div> <div> <div></div> <div>73%</div> <div>17%</div> <div>8%</div> </div> </div>
1	G	254	<div> <div>14%</div> <div> <div></div> <div>74%</div> <div>15%</div> <div>8%</div> </div> </div>
2	B	147	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>•</div> </div> </div>
2	D	147	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>•</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	147	
2	H	147	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA-70.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	15	0	0
			1889	1195	321	372	1			
1	C	233	Total	C	N	O	S	22	0	0
			1889	1195	321	372	1			
1	E	233	Total	C	N	O	S	22	0	0
			1889	1195	321	372	1			
1	G	233	Total	C	N	O	S	18	0	0
			1889	1195	321	372	1			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	373	GLY	-	EXPRESSION TAG	UNP Q8KHU9
A	374	PRO	-	EXPRESSION TAG	UNP Q8KHU9
A	375	LEU	-	EXPRESSION TAG	UNP Q8KHU9
A	376	GLY	-	EXPRESSION TAG	UNP Q8KHU9
A	377	SER	-	EXPRESSION TAG	UNP Q8KHU9
C	373	GLY	-	EXPRESSION TAG	UNP Q8KHU9
C	374	PRO	-	EXPRESSION TAG	UNP Q8KHU9
C	375	LEU	-	EXPRESSION TAG	UNP Q8KHU9
C	376	GLY	-	EXPRESSION TAG	UNP Q8KHU9
C	377	SER	-	EXPRESSION TAG	UNP Q8KHU9
E	373	GLY	-	EXPRESSION TAG	UNP Q8KHU9
E	374	PRO	-	EXPRESSION TAG	UNP Q8KHU9
E	375	LEU	-	EXPRESSION TAG	UNP Q8KHU9
E	376	GLY	-	EXPRESSION TAG	UNP Q8KHU9
E	377	SER	-	EXPRESSION TAG	UNP Q8KHU9
G	373	GLY	-	EXPRESSION TAG	UNP Q8KHU9
G	374	PRO	-	EXPRESSION TAG	UNP Q8KHU9
G	375	LEU	-	EXPRESSION TAG	UNP Q8KHU9
G	376	GLY	-	EXPRESSION TAG	UNP Q8KHU9
G	377	SER	-	EXPRESSION TAG	UNP Q8KHU9

- Molecule 2 is a protein called HA-17.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	143	Total	C	N	O	S	0	0	0
			1180	762	188	225	5			
2	D	143	Total	C	N	O	S	0	0	0
			1180	762	188	225	5			
2	F	142	Total	C	N	O	S	0	0	0
			1171	757	187	222	5			
2	H	143	Total	C	N	O	S	0	0	0
			1180	762	188	225	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	EXPRESSION TAG	UNP Q45878
B	1	PRO	-	EXPRESSION TAG	UNP Q45878
D	0	GLY	-	EXPRESSION TAG	UNP Q45878
D	1	PRO	-	EXPRESSION TAG	UNP Q45878
F	0	GLY	-	EXPRESSION TAG	UNP Q45878
F	1	PRO	-	EXPRESSION TAG	UNP Q45878
H	0	GLY	-	EXPRESSION TAG	UNP Q45878
H	1	PRO	-	EXPRESSION TAG	UNP Q45878

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	75	Total	O	0	0
			75	75		
3	B	43	Total	O	0	0
			43	43		
3	C	45	Total	O	0	0
			45	45		
3	D	51	Total	O	0	0
			51	51		
3	E	66	Total	O	0	0
			66	66		
3	F	47	Total	O	0	0
			47	47		
3	G	40	Total	O	0	0
			40	40		
3	H	46	Total	O	0	0
			46	46		




• Molecule 2: HA-17




• Molecule 2: HA-17



• Molecule 2: HA-17



• Molecule 2: HA-17



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.81Å 228.86Å 78.72Å 90.00° 92.05° 90.00°	Depositor
Resolution (Å)	45.78 – 2.40 46.27 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.4 (45.78-2.40) 99.4 (46.27-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.197 , 0.249 0.194 , 0.243	Depositor DCC
R_{free} test set	3181 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	1.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12680	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1923	0.45	0/2619
1	C	0.23	0/1923	0.45	0/2619
1	E	0.23	0/1923	0.45	0/2619
1	G	0.23	0/1923	0.44	0/2619
2	B	0.24	0/1210	0.42	0/1643
2	D	0.24	0/1210	0.41	0/1643
2	F	0.24	0/1201	0.42	0/1631
2	H	0.24	0/1210	0.41	0/1643
All	All	0.24	0/12523	0.43	0/17036

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1889	0	1849	23	0
1	C	1889	0	1849	26	0
1	E	1889	0	1849	27	0
1	G	1889	0	1849	23	0
2	B	1180	0	1141	9	0
2	D	1180	0	1141	12	0
2	F	1171	0	1135	12	0
2	H	1180	0	1141	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	75	0	0	2	0
3	B	43	0	0	1	0
3	C	45	0	0	0	0
3	D	51	0	0	1	0
3	E	66	0	0	3	0
3	F	47	0	0	0	0
3	G	40	0	0	0	0
3	H	46	0	0	3	0
All	All	12680	0	11954	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:505:ARG:NH2	2:H:57:LYS:HE2	1.38	1.36
1:E:505:ARG:HH22	2:H:57:LYS:CE	1.65	1.07
1:E:505:ARG:NH2	2:H:57:LYS:CE	2.32	0.80
1:A:550:LYS:NZ	3:A:755:HOH:O	2.20	0.75
1:G:546:ASN:OD1	2:H:142:LYS:NZ	2.19	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/254 (90%)	217 (95%)	11 (5%)	1 (0%)	34	48
1	C	229/254 (90%)	215 (94%)	13 (6%)	1 (0%)	34	48
1	E	229/254 (90%)	216 (94%)	12 (5%)	1 (0%)	34	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	229/254 (90%)	217 (95%)	11 (5%)	1 (0%)	34	48
2	B	141/147 (96%)	133 (94%)	8 (6%)	0	100	100
2	D	141/147 (96%)	135 (96%)	5 (4%)	1 (1%)	22	32
2	F	140/147 (95%)	133 (95%)	7 (5%)	0	100	100
2	H	141/147 (96%)	135 (96%)	6 (4%)	0	100	100
All	All	1479/1604 (92%)	1401 (95%)	73 (5%)	5 (0%)	41	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	94	VAL
1	E	391	ASP
1	G	391	ASP
1	C	391	ASP
1	A	488	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/232 (93%)	200 (93%)	15 (7%)	15	24
1	C	215/232 (93%)	200 (93%)	15 (7%)	15	24
1	E	215/232 (93%)	199 (93%)	16 (7%)	13	22
1	G	215/232 (93%)	200 (93%)	15 (7%)	15	24
2	B	135/138 (98%)	131 (97%)	4 (3%)	41	61
2	D	135/138 (98%)	133 (98%)	2 (2%)	65	80
2	F	134/138 (97%)	130 (97%)	4 (3%)	41	61
2	H	135/138 (98%)	128 (95%)	7 (5%)	23	38
All	All	1399/1480 (94%)	1321 (94%)	78 (6%)	21	34

5 of 78 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	126	LEU
1	E	465	ASN
2	H	65	ASN
1	E	406	PHE
1	E	414	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	118	ASN
1	E	392	ASN
1	G	392	ASN
2	D	26	ASN
2	F	26	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/254 (91%)	-0.23	6 (2%) 56 54	17, 39, 106, 140	4 (1%)
1	C	233/254 (91%)	0.60	28 (12%) 4 3	28, 66, 144, 173	6 (2%)
1	E	233/254 (91%)	0.08	16 (6%) 16 15	17, 53, 116, 149	6 (2%)
1	G	233/254 (91%)	0.62	35 (15%) 2 1	26, 68, 151, 175	5 (2%)
2	B	143/147 (97%)	-0.25	1 (0%) 87 86	26, 42, 78, 109	0
2	D	143/147 (97%)	-0.34	1 (0%) 87 86	28, 41, 68, 92	0
2	F	142/147 (96%)	-0.29	3 (2%) 63 61	26, 43, 78, 108	0
2	H	143/147 (97%)	-0.33	0 100 100	29, 44, 70, 112	0
All	All	1503/1604 (93%)	0.05	90 (5%) 21 20	17, 49, 118, 175	21 (1%)

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	380	GLU	11.8
1	C	400	ILE	9.9
1	C	397	ILE	7.9
1	C	481	TYR	7.8
1	G	397	ILE	7.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.