



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 08:42 am BST

PDB ID : 5LPF
Title : Kallikrein-related peptidase 10
Authors : Goettig, P.; Debela, M.; Magdolen, V.; Bode, W.; Brandstetter, H.
Deposited on : 2016-08-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	<div> <div>4%</div> <div> <div></div> <div>60%</div> <div>24%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	234	<div> <div>2%</div> <div> <div></div> <div>58%</div> <div>26%</div> <div>•</div> <div>14%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	301	-	-	X	-
2	SO4	A	304	-	-	X	-
2	SO4	B	303	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3277 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kallikrein-10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	35	0	0
			1553	974	280	284	15			
1	B	202	Total	C	N	O	S	40	0	0
			1554	974	282	283	15			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	68	Total	O	0	0
			68	68		
3	B	67	Total	O	0	0
			67	67		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kallikrein-10



• Molecule 1: Kallikrein-10



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.60Å 66.84Å 57.90Å 90.00° 104.29° 90.00°	Depositor
Resolution (Å)	36.69 – 2.70 51.94 – 2.70	Depositor EDS
% Data completeness (in resolution range)	94.4 (36.69-2.70) 94.6 (51.94-2.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.00 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.223 , 0.285 0.226 , 0.288	Depositor DCC
R_{free} test set	593 reflections (5.66%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3277	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7669e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	1/1589 (0.1%)	0.61	1/2159 (0.0%)
1	B	0.32	0/1591	0.65	1/2163 (0.0%)
All	All	0.35	1/3180 (0.0%)	0.63	2/4322 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	100	GLU	CB-CG	-6.26	1.40	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	82	LEU	CA-CB-CG	6.62	130.53	115.30
1	A	163	LEU	C-N-CA	-5.33	108.39	121.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	143	THR	Peptide
1	A	177	THR	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	59	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1553	0	1517	41	1
1	B	1554	0	1514	49	1
2	A	20	0	0	7	0
2	B	15	0	0	3	0
3	A	68	0	0	1	0
3	B	67	0	0	2	0
All	All	3277	0	3031	89	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:GLN:NE2	2:A:303:SO4:O1	1.82	1.11
1:B:83:ARG:NH2	2:B:303:SO4:O1	2.05	0.89
1:B:24:ARG:HD3	1:B:24:ARG:H	1.48	0.77
1:B:239:ASN:ND2	3:B:401:HOH:O	2.17	0.77
1:B:185:LEU:HD11	1:B:192:GLN:NE2	2.02	0.74
1:A:81:GLN:NE2	1:A:117:ARG:HH11	1.85	0.73
1:A:69:GLY:HA3	1:A:118:VAL:HG22	1.72	0.70
1:A:67:VAL:HG21	1:A:112:VAL:HG11	1.74	0.70
1:A:48:ASP:HB3	1:A:51:TRP:HB2	1.74	0.70
1:A:113:VAL:O	1:A:117:ARG:NH1	2.25	0.69
1:B:160:ILE:HD12	1:B:185:LEU:HD23	1.74	0.68
1:A:53:LEU:HD11	1:A:103:LEU:HG	1.76	0.68
1:B:24:ARG:HD3	1:B:24:ARG:N	2.08	0.67
1:B:195:SER:HA	1:B:213:LEU:HD12	1.77	0.65
1:A:21:PRO:HB2	1:A:157:CYS:H	1.61	0.65
1:B:31:VAL:HG22	1:B:67:VAL:HG23	1.78	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:GLN:HE22	1:B:118:VAL:HG21	1.62	0.64
1:B:81:GLN:NE2	1:B:118:VAL:HG21	2.13	0.63
1:B:243:ARG:NH2	3:B:402:HOH:O	2.32	0.63
1:A:98:THR:N	2:A:304:SO4:S	2.71	0.63
1:A:218:TYR:N	2:A:301:SO4:O4	2.31	0.61
1:B:81:GLN:HG2	2:B:303:SO4:O1	2.00	0.61
1:B:165:PRO:HB3	1:B:177:THR:HG21	1.82	0.61
1:B:213:LEU:HD21	1:B:226:ALA:HB1	1.83	0.61
1:A:115:GLY:O	1:A:119:ARG:HG3	2.01	0.61
1:B:185:LEU:CD1	1:B:188:GLN:HB3	2.32	0.59
1:A:179:ASN:HB3	1:A:233:LYS:HG3	1.83	0.58
2:A:301:SO4:O2	3:A:401:HOH:O	2.13	0.58
1:A:45:VAL:HG22	1:A:198:PRO:HB3	1.85	0.57
1:B:186:ASP:C	1:B:223(A):GLN:HG2	2.27	0.55
1:A:81:GLN:HE22	1:A:117:ARG:HH11	1.54	0.54
1:A:217:VAL:N	2:A:301:SO4:O4	2.40	0.54
1:A:160:ILE:HG12	1:A:183:ALA:HB1	1.92	0.52
1:A:48:ASP:OD1	1:A:49:GLN:N	2.43	0.52
1:B:233:LYS:NZ	2:B:302:SO4:O3	2.30	0.52
1:B:239:ASN:O	1:B:243:ARG:N	2.42	0.52
1:B:196:GLY:HA2	1:B:212:ILE:HG23	1.92	0.51
1:B:116:PRO:O	1:B:119:ARG:NH1	2.44	0.51
1:B:201:CYS:HB2	1:B:210:GLN:HG3	1.94	0.50
1:A:136:CYS:HB2	1:A:160:ILE:HG22	1.94	0.50
1:B:97:ARG:HB2	1:B:97:ARG:HH21	1.77	0.50
1:A:121:LEU:HD11	1:A:200:VAL:HG22	1.94	0.49
1:A:179:ASN:O	1:A:180:MET:HG2	2.12	0.49
1:B:180:MET:HG2	1:B:229:THR:HA	1.95	0.49
1:A:121:LEU:HD13	1:A:209:LEU:HB2	1.97	0.47
1:B:21:PRO:HA	1:B:156:THR:HA	1.97	0.47
1:A:220:CYS:HB2	1:B:170:VAL:O	2.15	0.47
1:A:194:ASP:O	1:A:213:LEU:HD23	2.13	0.47
1:A:81:GLN:NE2	1:A:118:VAL:HG21	2.30	0.47
1:B:213:LEU:HD11	1:B:217:VAL:HG23	1.97	0.47
1:B:136:CYS:HB2	1:B:160:ILE:HG22	1.96	0.46
1:A:165:PRO:HB2	1:A:177:THR:HB	1.96	0.46
1:A:211:GLY:HA2	1:A:231:ILE:HG12	1.96	0.46
1:B:160:ILE:HG13	1:B:184:GLY:O	2.16	0.46
1:B:121:LEU:HD23	1:B:209:LEU:HB2	1.97	0.46
1:B:63:LEU:HD12	1:B:106:LEU:HD11	1.96	0.46
1:A:123:LEU:HB3	1:A:235:MET:HE1	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:VAL:CG2	1:B:198:PRO:HB3	2.46	0.46
1:A:164:SER:C	1:A:166:LYS:H	2.19	0.45
1:A:61:LYS:HE3	1:A:61:LYS:HB3	1.72	0.45
1:B:80:GLU:HG2	1:B:81:GLN:N	2.31	0.45
1:A:49:GLN:O	1:A:112:VAL:HG22	2.17	0.45
1:B:128:ARG:HD2	1:B:129:CYS:O	2.17	0.44
1:A:87:ARG:HH12	1:A:243:ARG:NH1	2.15	0.44
1:B:185:LEU:HD12	1:B:188:GLN:HB3	1.97	0.44
1:B:31:VAL:O	1:B:44:GLY:N	2.45	0.44
1:B:48:ASP:HB3	1:B:51:TRP:HB2	1.99	0.44
1:A:216:GLY:N	2:A:301:SO4:O3	2.47	0.44
1:B:185:LEU:HD11	1:B:192:GLN:HE22	1.78	0.44
1:B:45:VAL:HG22	1:B:198:PRO:HB3	1.99	0.44
1:B:32:SER:HB3	1:B:141:TRP:CH2	2.53	0.43
1:A:173:PRO:HD2	1:B:218:TYR:HE1	1.84	0.43
1:B:160:ILE:HG21	1:B:199:LEU:HD11	2.00	0.43
1:A:98:THR:N	2:A:304:SO4:O2	2.52	0.43
1:B:124:PRO:HA	1:B:208:THR:HG21	1.99	0.43
1:B:234:TYR:N	1:B:234:TYR:CD1	2.87	0.43
1:B:199:LEU:HB2	1:B:228:TYR:CE2	2.54	0.42
1:A:165:PRO:O	1:A:177:THR:HG21	2.19	0.42
1:A:45:VAL:CG2	1:A:198:PRO:HB3	2.48	0.42
1:A:78:GLN:HG3	1:A:79:GLY:N	2.34	0.42
1:A:190:PRO:O	1:A:193:SER:OG	2.37	0.42
1:B:114:LEU:N	1:B:114:LEU:HD12	2.35	0.42
1:B:67:VAL:HG12	1:B:81:GLN:O	2.20	0.41
1:B:24:ARG:CD	1:B:24:ARG:H	2.26	0.41
1:A:22:CYS:HB2	1:A:155:LEU:O	2.20	0.41
1:B:56:ALA:HA	1:B:104:MET:HB2	2.03	0.40
1:A:240:LYS:HA	1:A:245:ASN:HA	2.03	0.40
1:A:22:CYS:SG	1:A:157:CYS:N	2.93	0.40
1:B:81:GLN:OE1	1:B:112:VAL:HG13	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:ASN:OXT	1:B:49:GLN:NE2[1_656]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/234 (83%)	184 (94%)	11 (6%)	0	100	100
1	B	194/234 (83%)	179 (92%)	15 (8%)	0	100	100
All	All	389/468 (83%)	363 (93%)	26 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/197 (87%)	164 (96%)	7 (4%)	30	59
1	B	172/197 (87%)	161 (94%)	11 (6%)	17	39
All	All	343/394 (87%)	325 (95%)	18 (5%)	23	49

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	ARG
1	A	119	ARG
1	A	121	LEU
1	A	159	SER
1	A	182	CYS
1	A	195	SER
1	A	243	ARG
1	B	24	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	64	TRP
1	B	95(A)	GLN
1	B	97	ARG
1	B	157	CYS
1	B	178	ASN
1	B	182	CYS
1	B	185	LEU
1	B	199	LEU
1	B	208	THR
1	B	223(A)	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	178	ASN
1	A	179	ASN
1	B	101	HIS
1	B	122	GLN
1	B	188	GLN
1	B	192	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	302	-	4,4,4	0.15	0	6,6,6	0.06	0
2	SO4	A	304	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	A	303	-	4,4,4	0.36	0	6,6,6	0.42	0
2	SO4	B	301	-	4,4,4	0.16	0	6,6,6	0.09	0
2	SO4	A	301	-	4,4,4	0.21	0	6,6,6	0.13	0
2	SO4	A	302	-	4,4,4	0.16	0	6,6,6	0.06	0
2	SO4	B	303	-	4,4,4	0.14	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	SO4	1	0
2	A	304	SO4	2	0
2	A	303	SO4	1	0
2	A	301	SO4	4	0
2	B	303	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/234 (86%)	0.21	9 (4%) 34 33	11, 26, 54, 81	8 (3%)
1	B	202/234 (86%)	0.32	5 (2%) 57 59	15, 31, 58, 72	10 (4%)
All	All	405/468 (86%)	0.26	14 (3%) 44 44	11, 27, 57, 81	18 (4%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	245	ASN	5.7
1	B	70	ASP	4.5
1	B	79	GLY	4.0
1	A	152	ASN	3.4
1	A	79	GLY	3.3
1	B	223	ALA	3.1
1	A	180	MET	3.0
1	A	153	LYS	3.0
1	B	245	ASN	2.8
1	B	185	LEU	2.3
1	A	178	ASN	2.3
1	A	244	SER	2.2
1	A	98	THR	2.2
1	A	60	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	303	5/5	0.43	0.39	50,50,71,96	0
2	SO4	A	303	5/5	0.75	0.33	58,68,85,85	0
2	SO4	B	301	5/5	0.83	0.27	35,38,62,81	0
2	SO4	A	304	5/5	0.85	0.36	65,70,81,92	0
2	SO4	B	302	5/5	0.87	0.23	59,64,68,71	0
2	SO4	A	302	5/5	0.90	0.20	31,34,39,51	0
2	SO4	A	301	5/5	0.90	0.33	38,42,65,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.