



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 11:07 am BST

PDB ID : 1LQU
Title : Mycobacterium tuberculosis FprA in complex with NADPH
Authors : Bossi, R.T.; Aliverti, A.; Raimondi, D.; Fischer, F.; Zanetti, G.; Ferrari, D.; Tahallah, N.; Maier, C.S.; Heck, A.J.R.; Rizzi, M.; Mattevi, A.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2002-05-13
Resolution : 1.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1060 (1.28-1.24)
Ramachandran outliers	138981	1029 (1.28-1.24)
Sidechain outliers	138945	1028 (1.28-1.24)
RSRZ outliers	127900	1004 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	
1	B	456	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	1423	-	-	X	-
2	ACT	A	1427	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	1424	-	-	X	-
2	ACT	B	1430	-	-	X	-
2	ACT	B	1431	-	-	X	-
2	ACT	B	1432	-	-	X	-
2	ACT	B	1433	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 18737 atoms, of which 10063 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FprA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	453	Total	C	H	N	O	S	186	10	0
			7047	2212	3535	630	662	8			
1	B	454	Total	C	H	N	O	S	180	6	0
			7021	2205	3527	625	656	8			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	A	1	Total	C	H	O	0	0
			7	2	3	2		

Continued on next page...

[illegible]

-
- The chemical structure of FAD (Flavin Adenine Dinucleotide) is shown. It consists of a flavin mononucleotide (FMN) part and an adenosine diphosphate (ADP) part. The FMN part includes a ribityl chain (C1-C3) and an isoalloxazine ring system (C4-C10). The ADP part includes an adenine base (C11-C17) and a ribose sugar (C18-C22). The structure is labeled with various atoms and bonds, including nitrogen (N), oxygen (O), carbon (C), and phosphorus (P). The overall structure is a complex molecule with multiple functional groups and a high degree of structural complexity.

- Molecule 4 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 77	C 21	H 29	N 7	O 17	P 3	10	0
4	B	1	Total 77	C 21	H 29	N 7	O 17	P 3	10	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	752	Total 2256	H 1504	O 752	1504	0
5	B	670	Total 2010	H 1340	O 670	1340	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FprA



• Molecule 1: FprA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.51Å 89.47Å 161.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.25 20.10 – 1.24	Depositor EDS
% Data completeness (in resolution range)	98.5 (40.00-1.25) 97.9 (20.10-1.24)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.24Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.126 , 0.142 0.127 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	10.5	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.49 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	18737	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, FAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3606	0.83	4/4896 (0.1%)
1	B	0.64	1/3588 (0.0%)	0.86	10/4874 (0.2%)
All	All	0.64	1/7194 (0.0%)	0.85	14/9770 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	327	VAL	CB-CG2	-6.31	1.39	1.52

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	441	ARG	NE-CZ-NH1	10.52	125.56	120.30
1	A	33	ASP	CB-CG-OD2	7.88	125.39	118.30
1	B	265	ARG	NE-CZ-NH2	-7.48	116.56	120.30
1	B	441	ARG	NE-CZ-NH2	-7.04	116.78	120.30
1	B	401	ASP	CB-CG-OD2	7.02	124.62	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3512	3535	3524	63	1
1	B	3494	3527	3523	42	1
2	A	20	15	15	30	0
2	B	24	18	18	21	0
3	A	53	33	31	0	0
3	B	53	33	31	0	0
4	A	48	29	26	4	0
4	B	48	29	26	6	0
5	A	752	1504	0	13	2
5	B	670	1340	0	22	2
All	All	8674	10063	7194	116	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ILE:CD1	2:A:1427:ACT:H1	1.08	1.53
2:B:1433:ACT:H3	4:B:2458:NDP:O1A	1.18	1.32
1:A:349[A]:ASN:ND2	1:B:342:PRO:HG3	1.51	1.25
1:A:60:ILE:HD12	2:A:1427:ACT:CH3	1.53	1.19
1:A:60:ILE:HD11	2:A:1427:ACT:H1	1.27	1.10

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:GLU:OE2	1:B:416:LYS:HZ1[3_545]	1.42	0.18
5:A:2019:HOH:O	5:B:3122:HOH:O[3_645]	2.12	0.08
5:A:1830:HOH:O	5:B:2885:HOH:O[4_455]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/456 (100%)	448 (98%)	9 (2%)	1 (0%)	47	18
1	B	458/456 (100%)	447 (98%)	9 (2%)	2 (0%)	34	10
All	All	916/912 (100%)	895 (98%)	18 (2%)	3 (0%)	41	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	TRP
1	B	46	TRP
1	B	262	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/365 (102%)	372 (100%)	1 (0%)	92	79
1	B	371/365 (102%)	369 (100%)	2 (0%)	88	66
All	All	744/730 (102%)	741 (100%)	3 (0%)	91	77

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	TYR
1	B	399	PRO
1	B	437	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	349	ASN
1	B	437	HIS
1	B	388	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	402	HIS
1	B	385	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	B	1431	-	1,3,3	1.68	0	0,3,3	0.00	-
3	FAD	B	2457	-	51,58,58	1.14	4 (7%)	60,89,89	2.03	9 (15%)
2	ACT	B	1425	-	1,3,3	0.64	0	0,3,3	0.00	-
2	ACT	A	1427	1	1,3,3	4.01	1 (100%)	0,3,3	0.00	-
2	ACT	B	1430	-	1,3,3	0.56	0	0,3,3	0.00	-
2	ACT	A	1423	-	1,3,3	1.20	0	0,3,3	0.00	-
4	NDP	B	2458	2	45,52,52	1.21	4 (8%)	53,80,80	1.57	8 (15%)
2	ACT	B	1424	-	1,3,3	1.14	0	0,3,3	0.00	-
3	FAD	A	1457	-	51,58,58	1.20	6 (11%)	60,89,89	1.81	7 (11%)
2	ACT	A	1429	-	1,3,3	1.35	0	0,3,3	0.00	-
2	ACT	B	1433	4	1,3,3	1.20	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NDP	A	1458	-	45,52,52	1.13	4 (8%)	53,80,80	1.34	6 (11%)
2	ACT	A	1428	-	1,3,3	1.20	0	0,3,3	0.00	-
2	ACT	A	1426	-	1,3,3	1.17	0	0,3,3	0.00	-
2	ACT	B	1432	-	1,3,3	0.11	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	1457	-	-	5/30/50/50	0/6/6/6
3	FAD	B	2457	-	-	6/30/50/50	0/6/6/6
4	NDP	B	2458	2	-	3/30/77/77	0/5/5/5
4	NDP	A	1458	-	-	4/30/77/77	0/5/5/5

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1427	ACT	CH3-C	-4.01	1.43	1.48
4	B	2458	NDP	O7N-C7N	3.59	1.33	1.24
3	A	1457	FAD	C4X-N5	3.48	1.38	1.33
3	B	2457	FAD	C4X-N5	3.43	1.38	1.33
3	B	2457	FAD	C10-N1	3.03	1.37	1.33

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2457	FAD	C4-N3-C2	10.50	124.01	115.14
3	A	1457	FAD	C4-N3-C2	9.39	123.07	115.14
4	B	2458	NDP	O4D-C1D-N1N	5.03	117.89	108.06
4	B	2458	NDP	C1D-N1N-C6N	-4.92	110.24	120.83
3	B	2457	FAD	C4X-C4-N3	-4.69	117.02	123.43

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2458	NDP	C2B-O2B-P2B-O2X
4	B	2458	NDP	O4D-C1D-N1N-C2N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1458	NDP	C2B-O2B-P2B-O2X
4	A	1458	NDP	O4D-C1D-N1N-C2N
3	B	2457	FAD	C2'-C3'-C4'-O4'


There are no ring outliers.


10 monomers are involved in 51 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1431	ACT	4	0
2	A	1427	ACT	22	0
2	B	1430	ACT	2	0
2	A	1423	ACT	7	0
4	B	2458	NDP	6	0
2	B	1424	ACT	3	0
2	A	1429	ACT	1	0
2	B	1433	ACT	6	0
4	A	1458	NDP	4	0
2	B	1432	ACT	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	453/456 (99%)	-0.48	10 (2%) 62 52	7, 12, 22, 36	0
1	B	454/456 (99%)	-0.41	18 (3%) 38 31	7, 13, 25, 37	0
All	All	907/912 (99%)	-0.45	28 (3%) 49 41	7, 13, 24, 37	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	393	ALA	5.6
1	B	262	PRO	5.4
1	B	282	ARG	5.2
1	B	231	ASP	4.8
1	B	281	LYS	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	A	1426	4/4	0.40	0.31	39,40,40,41	0
2	ACT	B	1430	4/4	0.44	0.26	49,49,49,49	0
2	ACT	A	1429	4/4	0.67	0.18	27,29,30,30	0
2	ACT	B	1432	4/4	0.74	0.33	30,32,32,33	0
2	ACT	A	1423	4/4	0.79	0.30	25,28,29,29	0
2	ACT	B	1433	4/4	0.83	0.30	24,29,30,30	0
2	ACT	A	1427	4/4	0.84	0.25	19,25,25,28	0
2	ACT	B	1425	4/4	0.87	0.21	27,27,27,27	0
2	ACT	B	1424	4/4	0.91	0.22	18,19,24,25	0
2	ACT	B	1431	4/4	0.96	0.05	16,17,19,20	0
2	ACT	A	1428	4/4	0.97	0.04	14,15,16,17	0
3	FAD	A	1457	53/53	0.99	0.05	6,7,10,10	9
4	NDP	B	2458	48/48	0.99	0.04	7,9,12,14	10
3	FAD	B	2457	53/53	0.99	0.05	5,7,9,10	9
4	NDP	A	1458	48/48	1.00	0.04	7,8,12,13	10

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NDP B 2458:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 2457:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NDP A 1458:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.