



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:22 am BST

PDB ID : 4LQ0
Title : Crystal structure of the I-LtrWI LAGLIDADG homing endonuclease bound to target DNA.
Authors : Chik, J.; Shen, B.; Stoddard, B.
Deposited on : 2013-07-16
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	<div> <div style="width: 61%; background-color: green;"></div> <div style="width: 31%; background-color: yellow;"></div> <div style="width: 6%; background-color: orange;"></div> <div style="width: 2%; background-color: red;"></div> <div style="width: 2%; background-color: grey;"></div> </div> <div>61% 31% 6% .</div>
2	B	26	<div> <div style="width: 42%; background-color: green;"></div> <div style="width: 58%; background-color: yellow;"></div> </div> <div>42% 58%</div>
3	C	26	<div> <div style="width: 58%; background-color: green;"></div> <div style="width: 42%; background-color: yellow;"></div> </div> <div>58% 42%</div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LAGLIDADG homing endonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2421	1572	396	440	13			

- Molecule 2 is a DNA chain called top strand DNA target.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	26	Total	C	N	O	P	0	0	0
			531	256	95	155	25			

- Molecule 3 is a DNA chain called bottom strand DNA target.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	26	Total	C	N	O	P	0	0	0
			529	255	96	153	25			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Ca	0	0
			3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	12	Total 12	O 12	0	0
7	B	4	Total 4	O 4	0	0
7	C	2	Total 2	O 2	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LAGLIDADG homing endonuclease



- Molecule 2: top strand DNA target



- Molecule 3: bottom strand DNA target



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.55Å 71.96Å 168.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.34 – 2.68 44.30 – 2.67	Depositor EDS
% Data completeness (in resolution range)	94.6 (44.34-2.68) 94.2 (44.30-2.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.08 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.7.0025	Depositor
R, R_{free}	0.197 , 0.271 0.195 , 0.270	Depositor DCC
R_{free} test set	688 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	53.6	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3517	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, PEG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2466	0.82	1/3313 (0.0%)
2	B	0.50	0/595	0.97	0/917
3	C	0.43	0/593	0.86	0/913
All	All	0.60	0/3654	0.85	1/5143 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	273	LEU	CA-CB-CG	6.00	129.09	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2421	0	2495	90	0
2	B	531	0	297	16	0
3	C	529	0	296	11	0
4	A	3	0	0	0	0
5	B	7	0	10	0	0
6	C	8	0	12	1	0
7	A	12	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	4	0	0	0	0
7	C	2	0	0	0	0
All	All	3517	0	3110	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:17:DT:H2''	2:B:18:DA:H5''	1.34	1.08
1:A:50:ASN:ND2	1:A:52:LYS:HB2	1.86	0.90
1:A:102:LEU:O	1:A:107:ARG:HD2	1.76	0.85
3:C:24:DT:H4'	3:C:25:DG:OP1	1.75	0.85
1:A:28:ASN:HD22	1:A:29:VAL:H	1.23	0.85
1:A:196:LEU:H	1:A:196:LEU:HD12	1.41	0.83
3:C:12:DA:H2''	3:C:13:DT:OP2	1.79	0.82
2:B:23:DA:H4'	2:B:24:DC:OP1	1.80	0.80
1:A:178:ILE:HG23	1:A:182:ALA:CB	2.13	0.79
2:B:17:DT:H2''	2:B:18:DA:C5'	2.11	0.79
1:A:178:ILE:HG23	1:A:182:ALA:HB2	1.64	0.78
1:A:201:GLN:HE21	1:A:242:THR:CG2	1.98	0.76
1:A:61:LYS:HD2	1:A:70:ILE:CD1	2.16	0.75
1:A:21:ALA:HB1	1:A:181:GLU:OE1	1.88	0.74
1:A:247:ILE:HG22	1:A:252:ILE:HD13	1.68	0.74
2:B:25:DC:H2''	2:B:26:DC:C5	2.24	0.73
1:A:28:ASN:ND2	1:A:29:VAL:H	1.86	0.73
1:A:129:ASN:O	1:A:132:MET:HB2	1.90	0.71
1:A:14:TYR:OH	1:A:179:GLU:HG2	1.92	0.70
1:A:61:LYS:HD2	1:A:70:ILE:HD11	1.75	0.69
1:A:267:ASN:HD21	1:A:300:ASN:H	1.40	0.68
3:C:10:DA:H2''	3:C:11:DC:OP2	1.93	0.68
2:B:17:DT:C2'	2:B:18:DA:H5''	2.20	0.66
1:A:14:TYR:OH	1:A:179:GLU:CG	2.44	0.65
1:A:216:ILE:HG21	1:A:228:VAL:HG13	1.79	0.65
1:A:201:GLN:HE21	1:A:242:THR:HG23	1.62	0.64
1:A:242:THR:HB	3:C:6:DA:OP2	1.98	0.64
1:A:303:ARG:CG	1:A:304:ILE:H	2.11	0.64
3:C:25:DG:H4'	3:C:26:DG:OP1	1.97	0.64
1:A:15:ILE:HD11	1:A:59:LEU:HG	1.81	0.63
1:A:186:VAL:HG21	1:A:296:LYS:HB2	1.79	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:VAL:HG23	1:A:186:VAL:O	1.99	0.62
1:A:50:ASN:HD22	1:A:52:LYS:HB2	1.63	0.62
1:A:118:GLU:O	1:A:122:ASN:ND2	2.34	0.60
1:A:171:LYS:HG3	1:A:222:LEU:HD22	1.82	0.60
1:A:52:LYS:HD2	1:A:304:ILE:HG12	1.81	0.60
1:A:267:ASN:ND2	1:A:300:ASN:H	1.98	0.60
1:A:217:GLU:HG3	1:A:228:VAL:HG21	1.84	0.60
1:A:15:ILE:HD13	1:A:63:THR:OG1	2.02	0.59
1:A:242:THR:HG22	3:C:5:DA:H2'	1.84	0.59
1:A:50:ASN:HD21	1:A:52:LYS:HB2	1.68	0.59
1:A:28:ASN:HD22	1:A:29:VAL:N	1.96	0.58
1:A:247:ILE:CG2	1:A:252:ILE:HD13	2.34	0.58
1:A:216:ILE:CG2	1:A:228:VAL:HG13	2.34	0.57
1:A:186:VAL:CG2	1:A:296:LYS:HB2	2.33	0.57
1:A:44:TYR:HB3	1:A:84:SER:HB3	1.87	0.56
1:A:303:ARG:HG2	1:A:304:ILE:N	2.21	0.56
1:A:303:ARG:HG2	1:A:304:ILE:H	1.71	0.56
1:A:124:GLU:O	1:A:130:GLY:HA3	2.06	0.55
2:B:21:DT:H2''	2:B:22:DA:O5'	2.05	0.55
3:C:24:DT:C4'	3:C:25:DG:OP1	2.50	0.54
1:A:186:VAL:O	1:A:296:LYS:HE3	2.07	0.54
2:B:13:DA:H1'	2:B:14:DT:O4'	2.08	0.54
1:A:201:GLN:HE21	1:A:242:THR:HG21	1.72	0.53
1:A:303:ARG:CG	1:A:304:ILE:N	2.71	0.53
1:A:201:GLN:NE2	1:A:242:THR:CG2	2.70	0.53
1:A:186:VAL:HG23	1:A:296:LYS:HD2	1.91	0.53
2:B:23:DA:H1'	2:B:24:DC:H5'	1.92	0.52
3:C:23:DC:H2'	3:C:24:DT:C5	2.45	0.52
1:A:91:ASN:O	1:A:95:PRO:HG2	2.11	0.51
2:B:22:DA:H4'	2:B:23:DA:OP1	2.11	0.51
1:A:190:ASN:HD22	1:A:198:LYS:HD3	1.75	0.51
1:A:73:ASN:HD22	1:A:73:ASN:C	2.14	0.50
1:A:201:GLN:NE2	1:A:242:THR:HG23	2.25	0.50
1:A:28:ASN:ND2	1:A:29:VAL:N	2.58	0.49
1:A:268:TYR:OH	1:A:272:LYS:HD2	2.13	0.49
2:B:2:DA:H2''	2:B:3:DG:OP2	2.13	0.48
1:A:179:GLU:OE2	1:A:262:GLY:N	2.38	0.48
1:A:181:GLU:HG3	2:B:14:DT:H3'	1.96	0.48
1:A:120:ILE:HA	1:A:125:HIS:CD2	2.50	0.47
1:A:247:ILE:HA	1:A:251:ILE:HB	1.96	0.47
1:A:106:LYS:NZ	7:A:508:HOH:O	2.48	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:LEU:HD12	1:A:131:LEU:HA	1.53	0.47
3:C:6:DA:H2''	3:C:7:DA:O5'	2.15	0.46
1:A:243:SER:OG	1:A:246:GLU:HG3	2.15	0.46
1:A:134:ILE:HG13	1:A:135:LEU:N	2.30	0.46
1:A:110:PHE:O	1:A:113:PHE:HB3	2.16	0.45
1:A:209:HIS:HB3	1:A:211:ARG:HG2	1.98	0.45
1:A:247:ILE:CG2	1:A:252:ILE:CD1	2.95	0.45
1:A:8:ILE:HD12	1:A:8:ILE:N	2.32	0.45
3:C:23:DC:H2'	3:C:24:DT:C6	2.52	0.45
1:A:94:ILE:O	1:A:95:PRO:C	2.55	0.45
1:A:186:VAL:CG2	1:A:186:VAL:O	2.65	0.44
2:B:3:DG:H2''	2:B:4:DT:OP2	2.17	0.44
2:B:3:DG:H1'	2:B:4:DT:H5'	1.99	0.44
2:B:3:DG:H4'	6:C:101:TRS:H22	1.99	0.44
1:A:41:VAL:CG2	1:A:120:ILE:HD11	2.48	0.44
1:A:184:PHE:HE2	1:A:264:LYS:HG2	1.83	0.44
1:A:56:LEU:HD12	1:A:56:LEU:HA	1.53	0.43
1:A:102:LEU:HB2	1:A:107:ARG:HB2	2.01	0.43
2:B:22:DA:C2	2:B:23:DA:C4	3.06	0.43
1:A:216:ILE:HG22	1:A:228:VAL:HG22	2.00	0.43
1:A:242:THR:CG2	3:C:5:DA:H2'	2.46	0.43
1:A:264:LYS:O	1:A:265:LEU:C	2.57	0.42
1:A:71:TYR:O	1:A:79:LEU:HA	2.19	0.42
1:A:14:TYR:CE1	1:A:179:GLU:HG3	2.54	0.42
1:A:12:ASN:HD22	1:A:63:THR:CG2	2.32	0.42
1:A:73:ASN:HD22	1:A:74:PRO:N	2.17	0.42
2:B:7:DT:H2''	2:B:8:DG:H5'	2.01	0.42
1:A:201:GLN:NE2	1:A:242:THR:HG21	2.33	0.42
1:A:190:ASN:ND2	1:A:198:LYS:HD3	2.34	0.42
1:A:15:ILE:O	1:A:19:VAL:HG13	2.20	0.42
1:A:58:GLU:HA	1:A:58:GLU:OE1	2.20	0.42
1:A:14:TYR:O	1:A:176:GLY:HA3	2.21	0.41
1:A:296:LYS:NZ	7:A:512:HOH:O	2.53	0.41
1:A:12:ASN:ND2	1:A:63:THR:HG23	2.36	0.41
1:A:14:TYR:HH	1:A:179:GLU:HG2	1.84	0.41
1:A:43:THR:HG22	1:A:89:ILE:HD13	2.03	0.40
1:A:278:MET:HB3	1:A:283:HIS:ND1	2.36	0.40
1:A:11:LEU:HD23	1:A:59:LEU:HD11	2.03	0.40
1:A:288:GLY:O	1:A:289:LEU:C	2.59	0.40
1:A:178:ILE:O	1:A:179:GLU:C	2.59	0.40
1:A:89:ILE:HA	1:A:93:ILE:HG12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/304 (97%)	279 (94%)	17 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/279 (98%)	229 (84%)	44 (16%)	2	5

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	VAL
1	A	20	ASP
1	A	28	ASN
1	A	41	VAL
1	A	43	THR
1	A	52	LYS
1	A	53	ASP
1	A	60	ILE
1	A	62	LYS
1	A	73	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	76	ASP
1	A	88	GLN
1	A	104	SER
1	A	106	LYS
1	A	116	VAL
1	A	117	VAL
1	A	124	GLU
1	A	126	LEU
1	A	127	THR
1	A	128	LEU
1	A	129	ASN
1	A	132	MET
1	A	134	ILE
1	A	136	SER
1	A	150	LYS
1	A	151	LYS
1	A	163	GLU
1	A	168	ASN
1	A	181	GLU
1	A	196	LEU
1	A	218	SER
1	A	220	VAL
1	A	222	LEU
1	A	224	ASN
1	A	234	ASN
1	A	244	ILE
1	A	261	ILE
1	A	273	LEU
1	A	278	MET
1	A	286	GLU
1	A	287	GLU
1	A	290	SER
1	A	294	GLU
1	A	296	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	ASN
1	A	28	ASN
1	A	50	ASN
1	A	73	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	122	ASN
1	A	125	HIS
1	A	148	ASN
1	A	168	ASN
1	A	190	ASN
1	A	201	GLN
1	A	234	ASN
1	A	249	ASN
1	A	267	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	TRS	C	101	-	7,7,7	0.44	0	9,9,9	0.41	0
5	PEG	B	101	-	6,6,6	0.59	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TRS	C	101	-	-	3/9/9/9	-
5	PEG	B	101	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	101	TRS	C3-C-C2-O2
6	C	101	TRS	N-C-C2-O2
5	B	101	PEG	O2-C3-C4-O4
6	C	101	TRS	C1-C-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	101	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/304 (98%)	-0.36	1 (0%) 94 95	28, 52, 78, 105	0
2	B	26/26 (100%)	-0.71	0 100 100	31, 59, 101, 117	0
3	C	26/26 (100%)	-0.76	0 100 100	35, 64, 99, 110	0
All	All	350/356 (98%)	-0.42	1 (0%) 94 95	28, 53, 82, 117	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	304	ILE	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	B	101	7/7	0.74	0.26	79,83,87,93	0
6	TRS	C	101	8/8	0.88	0.10	69,71,76,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CA	A	403	1/1	0.96	0.08	79,79,79,79	0
4	CA	A	401	1/1	0.99	0.14	42,42,42,42	0
4	CA	A	402	1/1	0.99	0.13	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.