



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2021 – 02:04 PM EDT

PDB ID : 7LR8
Title : Crystal structure of GH5_18-E153A from *Streptomyces cattleya* in complex with Manb1-4GlcNAc
Authors : Higgins, M.A.; Ryan, K.S.
Deposited on : 2021-02-16
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.18
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div>5%</div> </div> </div>
1	B	433	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>...</div> </div> </div>
2	C	2	<div> <div></div> <div>50%</div> <div>50%</div> </div>
2	E	2	<div> <div></div> <div>50%</div> <div>50%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	B	603	-	-	X	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein ScGH5_18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	413	Total	C	N	O	S	0	7	0
			3279	2068	611	593	7			
1	B	415	Total	C	N	O	S	0	5	0
			3253	2056	597	593	7			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	153	ALA	GLU	engineered mutation	UNP F8JJ04
A	426	LEU	-	expression tag	UNP F8JJ04
A	427	GLU	-	expression tag	UNP F8JJ04
A	428	HIS	-	expression tag	UNP F8JJ04
A	429	HIS	-	expression tag	UNP F8JJ04
A	430	HIS	-	expression tag	UNP F8JJ04
A	431	HIS	-	expression tag	UNP F8JJ04
A	432	HIS	-	expression tag	UNP F8JJ04
A	433	HIS	-	expression tag	UNP F8JJ04
B	153	ALA	GLU	engineered mutation	UNP F8JJ04
B	426	LEU	-	expression tag	UNP F8JJ04
B	427	GLU	-	expression tag	UNP F8JJ04
B	428	HIS	-	expression tag	UNP F8JJ04
B	429	HIS	-	expression tag	UNP F8JJ04
B	430	HIS	-	expression tag	UNP F8JJ04
B	431	HIS	-	expression tag	UNP F8JJ04
B	432	HIS	-	expression tag	UNP F8JJ04
B	433	HIS	-	expression tag	UNP F8JJ04

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			26	14	1	11			
2	C	2	Total	C	N	O	0	0	0
			26	14	1	11			

- Molecule 3 is 2-(2-ETHOXYETHOXY)ETHANOL (three-letter code: AE3) (formula: $C_6H_{14}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	6	3		
3	B	1	Total	C	O	0	0
			9	6	3		
3	B	1	Total	C	O	0	0
			9	6	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	402	Total 402	O 402	0	0
6	B	427	Total 427	O 427	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein ScGH5_18



- Molecule 1: Uncharacterized protein ScGH5_18



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.87Å 76.45Å 79.92Å 90.00° 117.24° 90.00°	Depositor
Resolution (Å)	38.98 – 1.60 38.98 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.3 (38.98-1.60) 98.3 (38.98-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.162 , 0.191 0.162 , 0.191	Depositor DCC
R_{free} test set	5393 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.879	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7459	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AE3, EDO, BMA, NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3381	0.54	0/4638
1	B	0.35	0/3356	0.53	0/4609
All	All	0.35	0/6737	0.53	0/9247

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3279	0	3116	27	0
1	B	3253	0	3072	21	1
2	C	26	0	24	0	0
2	E	26	0	24	0	0
3	A	9	0	14	1	0
3	B	18	0	28	1	0
4	B	4	0	6	3	1
5	B	15	0	0	0	0
6	A	402	0	0	9	2
6	B	427	0	0	7	0
All	All	7459	0	6284	49	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:VAL:HG22	1:A:373:ARG:HA	1.59	0.84
1:B:152:ASN:ND2	6:B:702:HOH:O	2.13	0.80
4:B:603:EDO:O1	6:B:701:HOH:O	2.04	0.76
1:A:178[A]:ARG:NH1	6:A:604:HOH:O	2.23	0.71
1:B:67:THR:HB	4:B:603:EDO:O2	1.91	0.70
1:A:167:ARG:NH2	6:A:602:HOH:O	2.13	0.69
1:A:206:ASP:OD2	6:A:601:HOH:O	2.09	0.69
1:B:119:PHE:HB2	1:B:168[A]:VAL:HG11	1.76	0.65
1:A:119:PHE:HB2	1:A:168:VAL:HG11	1.77	0.65
1:A:364:VAL:HG13	1:A:370:ALA:HA	1.80	0.62
1:A:312[B]:ARG:NH2	6:A:610:HOH:O	2.32	0.61
1:B:141[B]:ASP:OD2	6:B:704:HOH:O	2.17	0.59
1:A:238:ARG:NH1	6:A:614:HOH:O	2.37	0.57
1:A:160[B]:HIS:HE2	1:A:206:ASP:CG	2.06	0.57
1:A:175:ASP:OD1	6:A:603:HOH:O	2.18	0.56
1:B:57:PRO:HD2	1:B:94:ASP:O	2.07	0.55
1:A:341:GLU:C	1:A:343:VAL:H	2.13	0.52
1:B:238:ARG:NH1	6:B:715:HOH:O	2.42	0.52
1:B:221:THR:HG21	1:B:253:ALA:HB1	1.92	0.51
1:B:67:THR:HB	4:B:603:EDO:C2	2.41	0.51
1:B:14:PRO:HD2	1:B:346:TRP:CZ2	2.47	0.50
1:B:316:PHE:HB2	1:B:321:TYR:CZ	2.47	0.50
1:A:114:HIS:CE1	3:A:501:AE3:H5C1	2.46	0.50
1:B:318[B]:GLU:HG3	6:B:799:HOH:O	2.12	0.49
1:A:221:THR:HG21	1:A:253:ALA:HB1	1.95	0.48
1:A:57:PRO:HD2	1:A:94:ASP:O	2.14	0.47
1:A:316:PHE:HB2	1:A:321:TYR:CZ	2.50	0.47
1:B:119:PHE:CB	1:B:168[A]:VAL:HG11	2.44	0.47
1:B:329:ASP:OD1	1:B:331:ARG:HG3	2.15	0.46
1:B:422:HIS:O	1:B:425:HIS:HB3	2.15	0.46
1:A:178[B]:ARG:NH2	6:A:604:HOH:O	2.41	0.45
1:A:202:TRP:CH2	1:A:223:VAL:HG12	2.52	0.45
1:A:424:VAL:O	1:A:425:HIS:HB2	2.17	0.44
1:A:266:LEU:HD22	1:A:292[A]:VAL:HG21	2.00	0.44
1:A:72[C]:ARG:HD3	6:B:919:HOH:O	2.18	0.44
1:B:27:PHE:CD2	1:B:100:LEU:HD13	2.54	0.43
1:A:311:ARG:HD3	6:A:955:HOH:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:VAL:C	1:A:341:GLU:H	2.22	0.43
1:B:422:HIS:O	1:B:422:HIS:HD2	2.02	0.43
1:B:142:ARG:O	1:B:191[A]:ARG:NH2	2.52	0.43
1:A:97:GLN:HA	1:A:98:GLY:HA3	1.80	0.42
1:A:72[B]:ARG:HD2	6:B:919:HOH:O	2.19	0.42
1:B:304:TRP:HA	1:B:305:CYS:HA	1.82	0.42
1:A:110:THR:O	1:A:115:ARG:HA	2.20	0.42
1:A:160[B]:HIS:HD2	6:A:602:HOH:O	2.02	0.41
1:B:13:SER:HB2	1:B:346:TRP:CH2	2.55	0.41
1:B:113:TRP:HB3	3:B:602:AE3:H4C1	2.02	0.41
1:A:90:ASP:HB3	1:A:146:LEU:HG	2.02	0.41
1:B:207:HIS:HA	1:B:208:PRO:HD3	1.95	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:603:EDO:O2	4:B:603:EDO:O2[2_555]	1.86	0.34
6:A:610:HOH:O	6:A:610:HOH:O[2_555]	1.95	0.25
1:B:86:GLU:OE2	1:B:354:ARG:NH1[4_445]	2.10	0.10
6:A:736:HOH:O	6:A:953:HOH:O[2_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/433 (97%)	405 (97%)	13 (3%)	1 (0%)	47	26
1	B	418/433 (96%)	409 (98%)	8 (2%)	1 (0%)	47	26
All	All	837/866 (97%)	814 (97%)	21 (2%)	2 (0%)	47	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	ALA
1	B	112	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/340 (96%)	322 (98%)	6 (2%)	59	36
1	B	325/340 (96%)	320 (98%)	5 (2%)	65	44
All	All	653/680 (96%)	642 (98%)	11 (2%)	60	38

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	94	ASP
1	A	191	ARG
1	A	194	LEU
1	A	228	PHE
1	A	364	VAL
1	B	29	HIS
1	B	94	ASP
1	B	194	LEU
1	B	329	ASP
1	B	425	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	152	ASN
1	B	422	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2	15,15,15	0.20	0	21,21,21	0.51	0
2	BMA	C	2	2	11,11,12	1.27	2 (18%)	15,15,17	1.12	1 (6%)
2	NAG	E	1	2	15,15,15	0.33	0	21,21,21	0.56	0
2	BMA	E	2	2	11,11,12	1.16	1 (9%)	15,15,17	1.03	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	0/6/26/26	0/1/1/1
2	BMA	C	2	2	-	0/2/19/22	1/1/1/1
2	NAG	E	1	2	-	0/6/26/26	0/1/1/1
2	BMA	E	2	2	-	0/2/19/22	1/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	BMA	C4-C3	2.82	1.59	1.52
2	E	2	BMA	C4-C3	2.39	1.58	1.52
2	C	2	BMA	C1-C2	2.00	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BMA	C1-O5-C5	3.10	116.39	112.19
2	E	2	BMA	C1-O5-C5	2.97	116.22	112.19

There are no chirality outliers.


There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	2	BMA	C1-C2-C3-C4-C5-O5
2	C	2	BMA	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	B	605	-	4,4,4	0.14	0	6,6,6	0.06	0
3	AE3	B	601	-	8,8,8	0.55	0	7,7,7	0.22	0
5	SO4	B	606	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	B	604	-	4,4,4	0.14	0	6,6,6	0.05	0
3	AE3	B	602	-	8,8,8	0.55	0	7,7,7	0.41	0
4	EDO	B	603	-	3,3,3	0.24	0	2,2,2	1.37	0
3	AE3	A	501	-	8,8,8	0.55	0	7,7,7	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AE3	B	601	-	-	5/6/6/6	-
4	EDO	B	603	-	-	1/1/1/1	-
3	AE3	A	501	-	-	3/6/6/6	-
3	AE3	B	602	-	-	0/6/6/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	AE3	O2-C3-C4-O3
3	B	601	AE3	O2-C3-C4-O3
3	A	501	AE3	C1-C2-O2-C3
3	A	501	AE3	C3-C4-O3-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	601	AE3	C4-C3-O2-C2
3	B	601	AE3	C1-C2-O2-C3
4	B	603	EDO	O1-C1-C2-O2
3	B	601	AE3	C6-C5-O3-C4
3	B	601	AE3	O3-C5-C6-O4

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	AE3	1	0
4	B	603	EDO	3	1
3	A	501	AE3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	413/433 (95%)	-0.20	9 (2%) 62 60	15, 24, 45, 75	0
1	B	415/433 (95%)	-0.23	6 (1%) 75 75	15, 23, 45, 75	0
All	All	828/866 (95%)	-0.22	15 (1%) 68 67	15, 24, 45, 75	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	425	HIS	5.8
1	A	425	HIS	4.0
1	B	426	LEU	3.8
1	A	342	ALA	3.3
1	B	422	HIS	3.2
1	A	408	ASP	2.9
1	A	404	ALA	2.8
1	A	407	ALA	2.7
1	B	12	ARG	2.5
1	B	404	ALA	2.3
1	A	348	THR	2.2
1	A	409	HIS	2.1
1	B	419	VAL	2.1
1	A	403	HIS	2.0
1	A	366	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	1	15/15	0.88	0.11	21,32,40,45	0
2	BMA	C	2	11/12	0.91	0.09	17,24,29,34	0
2	NAG	E	1	15/15	0.93	0.09	17,24,36,37	0
2	BMA	E	2	11/12	0.95	0.07	17,20,23,23	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	603	4/4	0.28	0.80	109,110,110,111	0
3	AE3	B	602	9/9	0.82	0.13	38,39,48,51	0
3	AE3	B	601	9/9	0.85	0.12	50,58,67,68	0
5	SO4	B	606	5/5	0.87	0.23	97,99,102,105	0
3	AE3	A	501	9/9	0.89	0.12	44,51,54,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	B	605	5/5	0.94	0.12	91,91,92,92	0
5	SO4	B	604	5/5	0.96	0.11	62,65,69,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.