



wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 05:56 am BST

PDB ID : 3LS3
Title : Padron0.9-ON (fluorescent state)
Authors : Brakemann, T.; Weber, G.; Trowitzsch, S.; Wahl, M.C.; Jakobs, S.
Deposited on : 2010-02-12
Resolution : 1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	<div> <div>7%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>8%</div> </div> </div>
1	B	235	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>9%</div> </div> </div>
1	C	235	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>9%</div> </div> </div>
1	D	235	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div></div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SPD	B	225	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Padron0.9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	16	0
			1846	1172	311	348	15			
1	B	215	Total	C	N	O	S	0	19	0
			1857	1188	312	344	13			
1	C	214	Total	C	N	O	S	0	16	0
			1816	1159	309	335	13			
1	D	225	Total	C	N	O	S	0	18	0
			1921	1223	329	355	14			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SPERMIDINE (three-letter code: SPD) (formula: $C_7H_{19}N_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			10	7	3		
3	B	1	Total	C	N	0	0
			10	7	3		
3	B	1	Total	C	N	0	0
			10	7	3		
3	C	1	Total	C	N	0	0
			10	7	3		
3	C	1	Total	C	N	0	0
			10	7	3		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	D	1	Total	C	N	O	S	0	0
			15	8	2	4	1		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	156	Total	O	0	10
			166	166		
5	B	169	Total	O	0	1
			170	170		
5	C	153	Total	O	0	2
			155	155		
5	D	202	Total	O	0	3
			205	205		


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
-
- Sequence logo for Chain A. The y-axis represents information content in bits (0.00 to 0.25). The x-axis lists amino acids. A color scale at the top indicates conservation levels: 7% (red), 83% (green), 9% (yellow), and 8% (grey).
- | Position | Amino Acid | Information Content (bits) |
|----------|------------|----------------------------|
| 1 | MET | 0.00 |
| 2 | ARG | 0.00 |
| 3 | GLY | 0.00 |
| 4 | SER | 0.00 |
| 5 | HIS | 0.00 |
| 6 | HIS | 0.00 |
| 7 | HIS | 0.00 |
| 8 | HIS | 0.00 |
| 9 | HIS | 0.00 |
| 10 | HIS | 0.00 |
| 11 | THR | 0.00 |
| 12 | ASP | 0.00 |
| 13 | PRO | 0.00 |
| 14 | M1 | 0.00 |
| 15 | S2 | 0.00 |
| 16 | V4 | 0.00 |
| 17 | I4 | 0.00 |
| 18 | K5 | 0.00 |
| 19 | I10 | 0.00 |
| 20 | K11 | 0.00 |
| 21 | L12 | 0.00 |
| 22 | G31 | 0.00 |
| 23 | M40 | 0.00 |
| 24 | K45 | 0.00 |
| 25 | I56 | 0.00 |
| 26 | CYG62 | 0.00 |
| 27 | V67 | 0.00 |
| 28 | F68 | 0.00 |
| 29 | E73 | 0.00 |
| 30 | N74 | 0.00 |
| 31 | I94 | 0.00 |
| 32 | D112 | 0.00 |
| 33 | V123 | 0.00 |
| 34 | N128 | 0.00 |
| 35 | R135 | 0.00 |
| 36 | S142 | 0.00 |
| 37 | D156 | 0.00 |
| 38 | Y159 | 0.00 |
| 39 | E164 | 0.00 |
| 40 | G165 | 0.00 |
| 41 | K174 | 0.00 |
| 42 | I195 | 0.00 |
| 43 | I198 | 0.00 |
| 44 | K202 | 0.00 |
| 45 | L209 | 0.00 |
| 46 | H210 | 0.00 |
| 47 | E211 | 0.00 |
| 48 | E218 | 0.00 |
| 49 | LEU | 0.00 |
| 50 | PRO | 0.00 |
| 51 | ARG | 0.00 |
| 52 | GLN | 0.00 |
| 53 | ALA | 0.00 |
| 54 | LYS | 0.00 |

- Chain B:
-
- 4% 83% 8% 9%
- MET ARG GLY SER HIS HIS HIS HIS HIS THR ASP PRO MET S2 V3 I4 K5 I56 M59 C1G62 N65 R66 V67 F68 E73 F79 E96 I107 D112 N128 S142 K145 L146 D156 Y159 R170 F173 K174 S190 H193 H194 L195
- H216 S217 E218 LEU PRO ARG GLN ALA LYS

- Chain C: 
- 

- Chain D:  5% 88% 8% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.86Å 103.81Å 121.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.65 19.97 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.3 (20.00-1.65) 99.3 (19.97-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.5.0063	Depositor
R, R_{free}	0.168 , 0.200 0.183 , 0.217	Depositor DCC
R_{free} test set	5479 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8231	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EPE, GYC, SPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1888	0.64	0/2544
1	B	0.66	0/1911	0.69	0/2570
1	C	0.65	0/1877	0.73	0/2528
1	D	0.71	0/1992	0.74	0/2685
All	All	0.65	0/7668	0.70	0/10327

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1846	0	1771	29	0
1	B	1857	0	1823	17	0
1	C	1816	0	1771	16	0
1	D	1921	0	1854	24	0
2	A	12	0	16	0	0
2	C	6	0	8	0	0
2	D	12	0	16	1	0
3	A	10	0	19	1	0
3	B	20	0	38	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	20	0	38	2	0
4	D	15	0	17	0	0
5	A	166	0	0	4	0
5	B	170	0	0	2	0
5	C	155	0	0	2	0
5	D	205	0	0	9	0
All	All	8231	0	7371	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:158[A]:ASN:OD1	5:D:508:HOH:O	1.55	1.24
1:C:158[A]:ASN:OD1	5:C:637:HOH:O	1.54	1.19
1:B:59[B]:MET:HG2	5:B:250:HOH:O	1.39	1.19
1:A:135[B]:ARG:HG2	1:A:135[B]:ARG:HH11	1.08	1.13
1:A:135[B]:ARG:CG	1:A:135[B]:ARG:HH11	1.76	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/235 (97%)	228 (100%)	1 (0%)	0	100	100
1	B	232/235 (99%)	230 (99%)	2 (1%)	0	100	100
1	C	227/235 (97%)	226 (100%)	1 (0%)	0	100	100
1	D	240/235 (102%)	239 (100%)	1 (0%)	0	100	100
All	All	928/940 (99%)	923 (100%)	5 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/202 (99%)	196 (98%)	4 (2%)	55	32
1	B	204/202 (101%)	202 (99%)	2 (1%)	76	62
1	C	199/202 (98%)	197 (99%)	2 (1%)	76	62
1	D	212/202 (105%)	209 (99%)	3 (1%)	67	46
All	All	815/808 (101%)	804 (99%)	11 (1%)	65	50

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	193	HIS
1	B	218	GLU
1	D	66	ARG
1	A	211	GLU
1	C	193	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	216	HIS
1	D	-8	HIS
1	D	21	HIS
1	D	38	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	GYC	C	62	1	22,22,23	2.75	6 (27%)	26,30,32	2.42	7 (26%)
1	GYC	B	62	1	22,22,23	3.06	6 (27%)	26,30,32	2.91	9 (34%)
1	GYC	D	62	1	22,22,23	2.95	5 (22%)	26,30,32	2.49	8 (30%)
1	GYC	A	62	1	22,22,23	2.73	5 (22%)	26,30,32	2.67	7 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GYC	C	62	1	-	2/9/29/30	0/2/2/2
1	GYC	B	62	1	-	2/9/29/30	0/2/2/2
1	GYC	D	62	1	-	1/9/29/30	0/2/2/2
1	GYC	A	62	1	-	2/9/29/30	0/2/2/2

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	62	GYC	CA2-C2	-9.09	1.39	1.48
1	D	62	GYC	CB2-CA2	8.94	1.42	1.35
1	A	62	GYC	CA2-C2	-7.97	1.40	1.48
1	C	62	GYC	CB2-CA2	7.61	1.41	1.35
1	B	62	GYC	CB2-CA2	7.51	1.41	1.35

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	GYC	CA2-C2-N3	9.73	107.97	103.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	62	GYC	CA2-C2-N3	9.71	107.96	103.37
1	C	62	GYC	CA2-C2-N3	9.30	107.77	103.37
1	D	62	GYC	CA2-C2-N3	8.45	107.37	103.37
1	B	62	GYC	O2-C2-CA2	-5.83	127.68	130.96

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	62	GYC	C-CA3-N3-C2
1	B	62	GYC	C-CA3-N3-C2
1	D	62	GYC	C-CA3-N3-C2
1	A	62	GYC	C-CA3-N3-C2
1	B	62	GYC	N2-CA2-CB2-CG2

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	62	GYC	1	0
1	B	62	GYC	2	0
1	D	62	GYC	1	0
1	A	62	GYC	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	225	-	5,5,5	0.34	0	5,5,5	0.42	0
2	GOL	D	227	-	5,5,5	0.44	0	5,5,5	0.32	0
2	GOL	C	225	-	5,5,5	0.31	0	5,5,5	0.31	0
3	SPD	B	226	-	9,9,9	0.28	0	8,8,8	0.77	0
3	SPD	C	226	-	9,9,9	0.29	0	8,8,8	0.67	0
3	SPD	B	225	-	9,9,9	0.28	0	8,8,8	0.93	0
4	EPE	D	225	-	15,15,15	1.03	1 (6%)	18,20,20	2.02	5 (27%)
3	SPD	C	227	-	9,9,9	0.28	0	8,8,8	0.61	0
3	SPD	A	227	-	9,9,9	0.33	0	8,8,8	0.69	0
2	GOL	D	226	-	5,5,5	0.47	0	5,5,5	0.37	0
2	GOL	A	226	-	5,5,5	0.38	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	225	-	-	4/4/4/4	-
2	GOL	D	227	-	-	4/4/4/4	-
2	GOL	C	225	-	-	2/4/4/4	-
3	SPD	B	226	-	-	4/7/7/7	-
3	SPD	C	226	-	-	5/7/7/7	-
3	SPD	B	225	-	-	5/7/7/7	-
4	EPE	D	225	-	-	6/9/19/19	0/1/1/1
3	SPD	C	227	-	-	3/7/7/7	-
3	SPD	A	227	-	-	5/7/7/7	-
2	GOL	D	226	-	-	2/4/4/4	-
2	GOL	A	226	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	225	EPE	C10-S	3.43	1.82	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	225	EPE	C5-N4-C3	4.68	119.36	108.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	225	EPE	C7-N4-C3	3.71	120.72	111.23
4	D	225	EPE	C7-N4-C5	3.39	119.92	111.23
4	D	225	EPE	O2S-S-C10	3.35	110.95	106.92
4	D	225	EPE	C6-N1-C2	2.90	115.35	108.83

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	227	GOL	O1-C1-C2-C3
2	D	227	GOL	C1-C2-C3-O3
2	C	225	GOL	O1-C1-C2-C3
4	D	225	EPE	S-C10-C9-N1
2	D	226	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	227	GOL	1	0
3	C	226	SPD	1	0
3	B	225	SPD	6	0
3	C	227	SPD	1	0
3	A	227	SPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/235 (91%)	0.34	16 (7%) 14 14	15, 20, 27, 47	1 (0%)
1	B	214/235 (91%)	0.18	10 (4%) 31 30	14, 19, 26, 40	0
1	C	213/235 (90%)	0.17	8 (3%) 40 40	11, 16, 24, 32	1 (0%)
1	D	224/235 (95%)	0.16	11 (4%) 29 28	11, 16, 25, 36	1 (0%)
All	All	866/940 (92%)	0.21	45 (5%) 27 26	11, 18, 26, 47	3 (0%)

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	9.3
1	D	0	PRO	6.6
1	B	2	SER	6.4
1	A	2	SER	5.5
1	D	165	GLY	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	GYC	C	62	21/22	0.94	0.09	15,19,22,27	0
1	GYC	A	62	21/22	0.95	0.08	19,22,27,31	0
1	GYC	D	62	21/22	0.96	0.07	15,17,22,25	0
1	GYC	B	62	21/22	0.96	0.07	17,20,24,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	C	225	6/6	0.64	0.17	52,55,56,58	0
3	SPD	C	227	10/10	0.72	0.24	39,55,57,57	0
2	GOL	A	226	6/6	0.74	0.21	64,64,64,64	0
2	GOL	D	227	6/6	0.75	0.19	55,58,59,59	0
3	SPD	B	226	10/10	0.78	0.14	50,55,56,56	0
3	SPD	B	225	10/10	0.78	0.16	50,56,60,61	0
3	SPD	A	227	10/10	0.79	0.16	47,51,57,57	0
3	SPD	C	226	10/10	0.79	0.12	59,61,63,63	0
2	GOL	D	226	6/6	0.80	0.18	59,61,62,63	0
2	GOL	A	225	6/6	0.81	0.19	79,79,79,80	0
4	EPE	D	225	15/15	0.89	0.36	55,67,69,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.