



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 11:58 PM BST

PDB ID : 6LS9
Title : Crystal structure of bovine herpesvirus 1 glycoprotein D
Authors : Yue, D.; Chen, Z.J.; Yang, F.L.; Ye, F.; Lin, S.; Cheng, Y.W.; Wang, J.C.;
Chen, Z.M.; Lin, X.; Yang, J.; Chen, H.; Zhang, Z.L.; You, Y.; Sun, H.L.;
Wen, A.; Wang, L.L.; Zheng, Y.; Cao, Y.; Li, Y.H.; Lu, G.W.
Deposited on : 2020-01-17
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	<div> <div>2%</div> <div> <div></div> <div>68%</div> <div>9%</div> <div>23%</div> </div> </div>
1	B	311	<div> <div>2%</div> <div> <div></div> <div>69%</div> <div>7%</div> <div>23%</div> </div> </div>
1	C	311	<div> <div>3%</div> <div> <div></div> <div>64%</div> <div>12%</div> <div>23%</div> </div> </div>
2	D	3	<div> <div></div> <div>100%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	241	Total	C	N	O	S	0	0	0
			1964	1278	329	345	12			
1	B	240	Total	C	N	O	S	0	0	0
			1960	1274	328	346	12			
1	C	240	Total	C	N	O	S	0	0	0
			1959	1273	328	346	12			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ALA	-	expression tag	UNP Q76PF1
A	-2	ASP	-	expression tag	UNP Q76PF1
A	-1	GLU	-	expression tag	UNP Q76PF1
A	0	PHE	-	expression tag	UNP Q76PF1
A	302	HIS	-	expression tag	UNP Q76PF1
A	303	HIS	-	expression tag	UNP Q76PF1
A	304	HIS	-	expression tag	UNP Q76PF1
A	305	HIS	-	expression tag	UNP Q76PF1
A	306	HIS	-	expression tag	UNP Q76PF1
A	307	HIS	-	expression tag	UNP Q76PF1
B	-3	ALA	-	expression tag	UNP Q76PF1
B	-2	ASP	-	expression tag	UNP Q76PF1
B	-1	GLU	-	expression tag	UNP Q76PF1
B	0	PHE	-	expression tag	UNP Q76PF1
B	302	HIS	-	expression tag	UNP Q76PF1
B	303	HIS	-	expression tag	UNP Q76PF1
B	304	HIS	-	expression tag	UNP Q76PF1
B	305	HIS	-	expression tag	UNP Q76PF1
B	306	HIS	-	expression tag	UNP Q76PF1
B	307	HIS	-	expression tag	UNP Q76PF1
C	-3	ALA	-	expression tag	UNP Q76PF1
C	-2	ASP	-	expression tag	UNP Q76PF1
C	-1	GLU	-	expression tag	UNP Q76PF1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	PHE	-	expression tag	UNP Q76PF1
C	302	HIS	-	expression tag	UNP Q76PF1
C	303	HIS	-	expression tag	UNP Q76PF1
C	304	HIS	-	expression tag	UNP Q76PF1
C	305	HIS	-	expression tag	UNP Q76PF1
C	306	HIS	-	expression tag	UNP Q76PF1
C	307	HIS	-	expression tag	UNP Q76PF1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			42	24	3	15			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		
4	B	23	Total	O	0	0
			23	23		
4	C	14	Total	O	0	0
			14	14		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

29% 68% 9% 23%

ALA ASP GLU PHE LEU PRO THR P4 A5 A39 ASP GLY ARG GLU Q44 P45 R49 R70 T71 L72 W73 V76 R77 R81 Y101 Y105 T106 E107 C108 C117 GLU F124 D138 L142 Q154 Y160 V165 D169 K183 T191 F192 R199 L207 T213 E219 W228 E242 Y246 E247 F248 PRO PRO ALA ALA ASP GLY GLY SER PRO ALA PRO PRO GLY ASP ASP GLU ALA ARG GLU GLU ASP ASP GLY ALA ALA GLY ARG GLU ASN GLY PRO GLY PRO GLU GLY ASP THR GLY ASP GLY ALA GLY ARG GLU ASN GLY GLY PRO GLY PRO GLU GLY SER

[illegible]

Chain C:

64% 12% 23%

ALA ASP GLU PHE LEU PRO THR A5 P6 M23 A39 ASP GLY ARG E43 M59 L60 A61 L62 L63 R70 W73 W74 A75 V76 R77 R78 R79 A80 A81 E93 E98 M103 E104 Y105 T106 E107 C108 C117 R118 L142 E152 Y160 I161 T164 V165 S173 G177 K183 R188 G189 Y190 T191 F192 G193 A198 R199 E219 A220 I224 Y246 E247 PRO PRO PRO ALA ALA ASP GLY GLY SER R78 R79 ALA PRO PRO GLY ASP ASP GLU ALA ARG ARG ASP GLU GLY THR GLU ASP GLY ALA ALA GLY ARG GLU ASN

GLY
PRO
PRO
GLY
PRO
GLU
GLY
ASP
GLY
GLU
SER
GLN
THR
PRO
GLU
ALA
ASN
GLY
HIS
HIS
HIS
HIS
HIS

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

100%

MAP1
MAP2
MAP3

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.27Å 104.39Å 98.77Å 90.00° 112.50° 90.00°	Depositor
Resolution (Å)	37.43 – 2.50 37.43 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.43-2.50) 99.7 (37.43-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.10.1 _2155	Depositor
R, R_{free}	0.205 , 0.243 0.206 , 0.243	Depositor DCC
R_{free} test set	1848 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	44.0	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6036	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2037	0.63	1/2782 (0.0%)
1	B	0.59	0/2032	0.68	2/2774 (0.1%)
1	C	0.83	6/2030 (0.3%)	0.71	3/2771 (0.1%)
All	All	0.70	6/6099 (0.1%)	0.67	6/8327 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	219	GLU	CD-OE1	-13.32	1.10	1.25
1	C	219	GLU	CD-OE2	-11.39	1.13	1.25
1	C	81	ARG	CZ-NH2	-7.14	1.23	1.33
1	C	81	ARG	NE-CZ	-6.06	1.25	1.33
1	C	219	GLU	CG-CD	-5.86	1.43	1.51
1	C	81	ARG	CZ-NH1	-5.24	1.26	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	78	ARG	N-CA-C	-9.84	84.43	111.00
1	B	199	ARG	NE-CZ-NH1	-7.64	116.48	120.30
1	B	21	ARG	NE-CZ-NH1	-7.05	116.77	120.30
1	A	138	ASP	CB-CG-OD1	6.69	124.32	118.30
1	C	199	ARG	NE-CZ-NH2	6.05	123.33	120.30
1	C	60	LEU	CA-CB-CG	5.62	128.23	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	1863	18	0
1	B	1960	0	1856	13	1
1	C	1959	0	1855	33	1
2	D	42	0	37	0	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	14	0	13	1	0
4	A	32	0	0	0	0
4	B	23	0	0	4	0
4	C	14	0	0	0	0
All	All	6036	0	5650	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:93:GLU:OE2	1:C:98:ARG:NE	1.61	1.32
1:B:92:ILE:O	4:B:1101:HOH:O	1.90	0.89
1:C:191:THR:HG22	1:C:192:PHE:H	1.38	0.87
1:C:191:THR:HG22	1:C:192:PHE:N	2.02	0.73
1:A:81:ARG:NH2	1:A:107:GLU:OE1	2.22	0.73
1:C:246:TYR:CD1	1:C:247:GLU:HG2	2.23	0.73
1:B:231:ARG:NH1	4:B:1103:HOH:O	2.15	0.73
1:B:241:GLU:O	4:B:1102:HOH:O	2.11	0.69
1:B:121:THR:HG23	1:B:122:PRO:O	1.92	0.69
1:C:61:ALA:HB1	1:C:63:ILE:HD13	1.76	0.67
1:A:70:ARG:NE	1:C:70:ARG:HD2	2.10	0.67
1:C:81:ARG:HD3	1:C:81:ARG:H	1.60	0.65
1:A:154:GLN:NE2	1:A:169:ASP:OD2	2.30	0.64
1:A:191:THR:HG23	1:A:192:PHE:H	1.62	0.63
1:C:246:TYR:HD1	1:C:247:GLU:HG2	1.64	0.63
1:C:98:ARG:NH1	1:C:193:GLY:O	2.31	0.62
1:C:75:ALA:O	1:C:78:ARG:O	2.18	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:ILE:O	1:C:164:THR:HG22	2.05	0.57
1:B:107:GLU:HB2	1:B:118:ARG:HB3	1.86	0.56
1:B:191:THR:HG23	1:B:192:PHE:H	1.71	0.56
1:A:105:TYR:HB3	1:A:117:CYS:SG	2.48	0.54
1:A:73:TRP:O	1:A:77:ARG:HG3	2.08	0.53
1:C:160:TYR:CE2	1:C:165:VAL:HB	2.43	0.53
1:C:246:TYR:CE1	1:C:247:GLU:HG2	2.45	0.52
1:C:93:GLU:OE2	1:C:98:ARG:CD	2.54	0.52
1:C:23:ASN:OD1	3:C:1001:NAG:O5	2.28	0.51
1:C:220:ALA:O	1:C:224:ILE:HD13	2.10	0.51
1:A:72:LEU:O	1:A:76:VAL:HG23	2.11	0.51
1:C:81:ARG:HD3	1:C:81:ARG:N	2.24	0.51
1:C:105:TYR:HB3	1:C:117:CYS:SG	2.51	0.51
1:A:191:THR:HG23	1:A:192:PHE:N	2.26	0.50
1:C:61:ALA:HB1	1:C:63:ILE:CD1	2.42	0.50
1:C:152:GLU:HG2	1:C:173:SER:HA	1.94	0.49
1:A:191:THR:CG2	1:A:192:PHE:N	2.76	0.48
1:C:5:ALA:HB3	1:C:6:PRO:HD3	1.94	0.48
1:B:191:THR:CG2	1:B:192:PHE:N	2.77	0.47
1:B:15:PRO:HD3	1:B:227:TYR:CG	2.49	0.47
1:A:108:CYS:HA	1:A:117:CYS:HA	1.95	0.47
1:C:108:CYS:HA	1:C:117:CYS:HA	1.98	0.45
1:B:191:THR:HG23	1:B:192:PHE:N	2.32	0.44
1:C:79:HIS:N	1:C:79:HIS:ND1	2.64	0.44
1:C:188:ARG:HD2	1:C:190:TYR:HE2	1.80	0.44
1:A:160:TYR:CE2	1:A:165:VAL:HB	2.52	0.44
1:A:101:TYR:HA	1:A:124:PHE:O	2.19	0.43
1:C:191:THR:CG2	1:C:192:PHE:N	2.72	0.43
1:C:103:MET:HG3	1:C:104:GLU:N	2.33	0.43
1:C:188:ARG:HB2	1:C:190:TYR:CD2	2.54	0.42
1:B:39:ALA:O	1:B:167:TYR:OH	2.34	0.42
1:C:220:ALA:O	1:C:224:ILE:CD1	2.67	0.42
1:C:107:GLU:HG2	1:C:118:ARG:HH11	1.85	0.42
1:A:199:ARG:HB3	1:A:199:ARG:CZ	2.49	0.42
1:B:185:GLY:O	4:B:1107:HOH:O	2.21	0.42
1:C:73:TRP:CZ2	1:C:77:ARG:HD2	2.55	0.41
1:C:160:TYR:CD2	1:C:165:VAL:HB	2.55	0.41
1:B:118:ARG:HD3	1:B:119:TYR:CE2	2.56	0.41
1:A:183:LYS:HE3	1:A:228:TRP:CE2	2.56	0.41
1:B:136:THR:HG22	1:B:143:ILE:HD11	2.03	0.41
1:A:73:TRP:CH2	1:A:77:ARG:HD2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:ARG:HH12	1:A:242:GLU:HG3	1.85	0.40
1:A:70:ARG:CD	1:C:70:ARG:HD2	2.51	0.40
1:A:142:LEU:HA	1:A:142:LEU:HD23	1.97	0.40
1:C:142:LEU:HA	1:C:142:LEU:HD12	1.90	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:TYR:OH	1:C:183:LYS:NZ[2_656]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/311 (76%)	228 (96%)	9 (4%)	0	100	100
1	B	236/311 (76%)	228 (97%)	8 (3%)	0	100	100
1	C	236/311 (76%)	226 (96%)	10 (4%)	0	100	100
All	All	709/933 (76%)	682 (96%)	27 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/246 (80%)	193 (98%)	5 (2%)	47	73
1	B	198/246 (80%)	190 (96%)	8 (4%)	31	56
1	C	197/246 (80%)	193 (98%)	4 (2%)	55	79
All	All	593/738 (80%)	576 (97%)	17 (3%)	42	69

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	THR
1	A	191	THR
1	A	207	LEU
1	A	213	THR
1	A	219	GLU
1	B	72	LEU
1	B	78	ARG
1	B	81	ARG
1	B	121	THR
1	B	140	LEU
1	B	191	THR
1	B	237	PRO
1	B	247	GLU
1	C	59	MET
1	C	79	HIS
1	C	81	ARG
1	C	107	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	1,2	14,14,15	0.68	1 (7%)	17,19,21	0.67	0
2	NAG	D	2	2	14,14,15	0.56	0	17,19,21	1.47	3 (17%)
2	NAG	D	3	2	14,14,15	2.87	2 (14%)	17,19,21	2.62	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	3	2	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	NAG	O5-C1	8.99	1.58	1.43
2	D	3	NAG	C1-C2	5.48	1.60	1.52
2	D	1	NAG	O5-C1	-2.40	1.39	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	NAG	C1-O5-C5	9.84	125.52	112.19
2	D	2	NAG	O4-C4-C3	3.54	118.52	110.35
2	D	2	NAG	C3-C4-C5	-3.27	104.41	110.24
2	D	3	NAG	C1-C2-N2	-3.16	105.09	110.49
2	D	3	NAG	C3-C4-C5	-2.61	105.58	110.24
2	D	2	NAG	O3-C3-C4	2.36	115.80	110.35

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	D	3	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	1001	1	14,14,15	0.47	0	17,19,21	0.49	0
3	NAG	A	1001	1	14,14,15	0.91	2 (14%)	17,19,21	0.71	0
3	NAG	C	1001	1	14,14,15	2.02	2 (14%)	17,19,21	1.18	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1001	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1001	1	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1001	NAG	O5-C1	-5.63	1.34	1.43
3	C	1001	NAG	C1-C2	4.54	1.59	1.52
3	A	1001	NAG	O5-C1	-2.48	1.39	1.43
3	A	1001	NAG	C1-C2	2.10	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1001	NAG	C1-O5-C5	-2.00	109.48	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1001	NAG	O5-C5-C6-O6
3	A	1001	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1001	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/311 (77%)	-0.20	6 (2%) 57 61	20, 30, 54, 112	0
1	B	240/311 (77%)	-0.16	6 (2%) 57 61	21, 34, 65, 105	0
1	C	240/311 (77%)	0.07	9 (3%) 40 43	23, 38, 65, 102	0
All	All	721/933 (77%)	-0.10	21 (2%) 51 55	20, 33, 65, 112	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	248	PRO	6.2
1	C	43	GLU	6.1
1	A	45	PRO	4.2
1	B	39	ALA	3.4
1	A	44	GLN	3.4
1	B	40	ASP	3.2
1	A	5	ALA	3.1
1	C	246	TYR	3.1
1	C	81	ARG	3.0
1	C	247	GLU	2.9
1	A	247	GLU	2.9
1	B	21	ARG	2.8
1	C	177	GLY	2.8
1	A	246	TYR	2.8
1	B	81	ARG	2.8
1	C	39	ALA	2.7
1	C	198	ALA	2.5
1	C	75	ALA	2.3
1	B	6	PRO	2.2
1	B	44	GLN	2.2
1	C	78	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	D	3	14/15	0.49	0.34	64,76,87,89	0
2	NAG	D	2	14/15	0.88	0.31	53,66,74,75	0
2	NAG	D	1	14/15	0.90	0.24	51,62,67,70	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	C	1001	14/15	0.62	0.33	51,60,65,72	0
3	NAG	A	1001	14/15	0.70	0.25	52,70,78,80	0
3	NAG	B	1001	14/15	0.79	0.34	50,57,61,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.