



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 6, 2021 – 02:33 PM EDT

PDB ID : 7LSU  
Title : Ruminococcus bromii Amy12-D392A with 63-a-D-glucosyl-maltotriose  
Authors : Koropatkin, N.M.; Cockburn, D.W.; Brown, H.A.; Kibler, R.D.  
Deposited on : 2021-02-18  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.22  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.22


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	799	
2	C	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	815	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6832 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pullulanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	765	Total	C	N	O	S	0	1	0
			5887	3685	959	1216	27			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	392	ALA	ASP	engineered mutation	UNP A0A2N0UU23

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Na	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	800	Total	O	0	0
			800	800		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pullulanase



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.09Å 99.20Å 167.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.91 – 1.95 41.88 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.91-1.95) 99.2 (41.88-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, $R_{free}$	0.188 , 0.212 0.197 , 0.219	Depositor DCC
$R_{free}$ test set	2913 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.0	Xtriage
Anisotropy	1.192	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 36.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6832	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CA, GOL, NA, ACT, BGC, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/6016	0.47	0/8179

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5887	0	5556	11	0
2	C	45	0	39	0	0
3	A	42	0	56	2	0
4	A	28	0	21	2	0
5	A	28	0	40	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	800	0	0	0	0
All	All	6832	0	5712	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:ARG:HH12	3:A:802:GOL:H31	1.58	0.68
1:A:419:TRP:HB3	1:A:456:ALA:HB3	1.92	0.51
1:A:738[B]:THR:HG23	1:A:740:GLU:H	1.79	0.47
1:A:276:ASN:HA	1:A:580:TYR:O	2.17	0.45
1:A:735:TYR:HD2	1:A:738[B]:THR:HG22	1.83	0.43
1:A:233:THR:HG21	1:A:238:GLY:HA2	2.01	0.43
1:A:463:ARG:HH12	3:A:802:GOL:C3	2.29	0.43
1:A:253:THR:HG22	4:A:817:ACT:H3	2.01	0.42
1:A:68:TYR:CE1	1:A:130:LYS:HB3	2.56	0.41
1:A:400:GLU:HG3	4:A:818:ACT:H3	2.03	0.41
1:A:615:SER:N	1:A:616:PRO:CD	2.84	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	764/799 (96%)	743 (97%)	19 (2%)	2 (0%)	41 30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	798	GLU
1	A	797	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	633/663 (96%)	627 (99%)	6 (1%)	78 77

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	201	TRP
1	A	259	PRO
1	A	471	MET
1	A	514	TYR
1	A	783	SER
1	A	784	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	C	1	2	12,12,12	0.43	0	17,17,17	0.50	0
2	GLC	C	2	2	11,11,12	0.28	0	15,15,17	0.63	0
2	GLC	C	3	2	11,11,12	0.24	0	15,15,17	0.60	0
2	GLC	C	4	2	11,11,12	0.22	0	15,15,17	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	C	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 2 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACT	A	817	-	1,3,3	1.37	0	0,3,3	0.00	-
5	PEG	A	812	-	6,6,6	0.46	0	5,5,5	0.21	0
3	GOL	A	802	-	5,5,5	0.32	0	5,5,5	0.17	0
5	PEG	A	814	-	6,6,6	0.47	0	5,5,5	0.17	0
3	GOL	A	805	-	5,5,5	0.32	0	5,5,5	0.18	0
5	PEG	A	813	-	6,6,6	0.46	0	5,5,5	0.22	0
3	GOL	A	804	-	5,5,5	0.28	0	5,5,5	0.22	0
4	ACT	A	808	-	1,3,3	1.29	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ACT	A	811	-	1,3,3	1.35	0	0,3,3	0.00	-
3	GOL	A	803	-	5,5,5	0.31	0	5,5,5	0.22	0
5	PEG	A	815	-	6,6,6	0.46	0	5,5,5	0.24	0
3	GOL	A	816	-	5,5,5	0.33	0	5,5,5	0.25	0
4	ACT	A	809	-	1,3,3	1.34	0	0,3,3	0.00	-
3	GOL	A	801	-	5,5,5	0.31	0	5,5,5	0.13	0
4	ACT	A	818	-	1,3,3	1.38	0	0,3,3	0.00	-
4	ACT	A	810	-	1,3,3	1.38	0	0,3,3	0.00	-
3	GOL	A	806	-	5,5,5	0.32	0	5,5,5	0.26	0
4	ACT	A	807	-	1,3,3	1.48	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	A	812	-	-	2/4/4/4	-
3	GOL	A	802	-	-	2/4/4/4	-
5	PEG	A	814	-	-	3/4/4/4	-
3	GOL	A	805	-	-	2/4/4/4	-
3	GOL	A	804	-	-	4/4/4/4	-
5	PEG	A	815	-	-	3/4/4/4	-
3	GOL	A	803	-	-	2/4/4/4	-
3	GOL	A	816	-	-	2/4/4/4	-
3	GOL	A	801	-	-	2/4/4/4	-
5	PEG	A	813	-	-	1/4/4/4	-
3	GOL	A	806	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	GOL	O1-C1-C2-O2
3	A	801	GOL	O1-C1-C2-C3
3	A	804	GOL	C1-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	805	GOL	O1-C1-C2-C3
3	A	816	GOL	C1-C2-C3-O3
3	A	802	GOL	O1-C1-C2-C3
3	A	803	GOL	O1-C1-C2-C3
3	A	804	GOL	O1-C1-C2-C3
3	A	806	GOL	O1-C1-C2-C3
3	A	806	GOL	C1-C2-C3-O3
3	A	802	GOL	O1-C1-C2-O2
3	A	803	GOL	O1-C1-C2-O2
3	A	804	GOL	O2-C2-C3-O3
3	A	805	GOL	O1-C1-C2-O2
3	A	816	GOL	O2-C2-C3-O3
5	A	812	PEG	O1-C1-C2-O2
5	A	815	PEG	O1-C1-C2-O2
5	A	815	PEG	O2-C3-C4-O4
5	A	815	PEG	C4-C3-O2-C2
5	A	813	PEG	O1-C1-C2-O2
5	A	814	PEG	O1-C1-C2-O2
3	A	804	GOL	O1-C1-C2-O2
3	A	806	GOL	O2-C2-C3-O3
5	A	814	PEG	C4-C3-O2-C2
3	A	806	GOL	O1-C1-C2-O2
5	A	814	PEG	O2-C3-C4-O4
5	A	812	PEG	C1-C2-O2-C3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	817	ACT	1	0
3	A	802	GOL	2	0
4	A	818	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	765/799 (95%)	0.11	17 (2%) 62 70	7, 14, 27, 38	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	GLY	4.0
1	A	112	LEU	3.5
1	A	798	GLU	3.1
1	A	114	ASP	2.8
1	A	113	VAL	2.8
1	A	115	GLY	2.8
1	A	69	THR	2.7
1	A	36	VAL	2.6
1	A	35	ALA	2.6
1	A	71	ASP	2.5
1	A	784	ASP	2.5
1	A	149	ASP	2.3
1	A	799	SER	2.3
1	A	797	PRO	2.1
1	A	126	VAL	2.1
1	A	776	ASN	2.0
1	A	150	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	C	2	11/12	0.93	0.12	14,15,16,17	0
2	GLC	C	4	11/12	0.93	0.11	21,22,22,22	0
2	GLC	C	3	11/12	0.95	0.10	17,18,20,20	0
2	BGC	C	1	12/12	0.95	0.15	13,14,15,15	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ACT	A	809	4/4	0.45	0.31	48,48,48,49	0
3	GOL	A	801	6/6	0.62	0.34	44,45,45,46	0
4	ACT	A	817	4/4	0.62	0.36	26,27,27,28	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	803	6/6	0.67	0.23	32,32,33,33	0
4	ACT	A	808	4/4	0.70	0.20	40,41,41,42	0
5	PEG	A	813	7/7	0.75	0.30	19,19,20,21	0
4	ACT	A	811	4/4	0.76	0.19	32,32,32,33	0
3	GOL	A	802	6/6	0.77	0.20	22,22,23,23	0
5	PEG	A	815	7/7	0.78	0.41	29,29,30,30	0
3	GOL	A	805	6/6	0.79	0.22	24,27,28,28	0
3	GOL	A	804	6/6	0.81	0.23	16,17,17,17	0
3	GOL	A	806	6/6	0.83	0.30	22,24,25,26	0
5	PEG	A	812	7/7	0.83	0.21	37,39,40,40	0
4	ACT	A	818	4/4	0.84	0.44	26,26,27,27	0
3	GOL	A	816	6/6	0.87	0.25	18,19,19,19	0
5	PEG	A	814	7/7	0.87	0.22	20,20,22,23	0
4	ACT	A	807	4/4	0.87	0.35	37,37,38,38	0
4	ACT	A	810	4/4	0.89	0.23	33,33,34,34	0
7	NA	A	820	1/1	0.90	0.28	20,20,20,20	0
6	CA	A	819	1/1	0.99	0.07	13,13,13,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.