



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 09:05 AM BST

PDB ID : 6LV5
Title : Ni- Carbonic Anhydrase II pH 7.8 0 atm CO2
Authors : Kim, C.U.; Kim, J.K.
Deposited on : 2020-02-02
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	301	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4554 atoms, of which 2117 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	257	4259	1378	2109	364	406	2	42	15	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	14	3	8	3	2	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ni	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	279	Total 279	O 279	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.40 Å 41.28 Å 71.92 Å 90.00° 104.03° 90.00°	Depositor
Resolution (Å)	30.00 – 1.20 29.14 – 1.20	Depositor EDS
% Data completeness (in resolution range)	96.6 (30.00-1.20) 96.6 (29.14-1.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.13 (at 1.20 Å)	Xtriage
Refinement program	REFMAC 5.8.0124 2015/06/02	Depositor
R, R_{free}	0.116 , 0.141 0.123 , 0.145	Depositor DCC
R_{free} test set	3734 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	8.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 59.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4554	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.59	34/2230 (1.5%)	1.37	32/3026 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14[A]	GLU	CD-OE2	18.23	1.45	1.25
1	A	14[B]	GLU	CD-OE2	18.23	1.45	1.25
1	A	234	GLU	CD-OE1	11.25	1.38	1.25
1	A	100	LEU	CA-CB	10.09	1.76	1.53
1	A	14[A]	GLU	CD-OE1	9.11	1.35	1.25
1	A	14[B]	GLU	CD-OE1	9.11	1.35	1.25
1	A	239[A]	GLU	CG-CD	8.07	1.64	1.51
1	A	239[B]	GLU	CG-CD	8.07	1.64	1.51
1	A	46	PRO	CA-C	7.81	1.68	1.52
1	A	89	ARG	CD-NE	-7.46	1.33	1.46
1	A	221	GLU	CD-OE1	-7.42	1.17	1.25
1	A	89	ARG	NE-CZ	-7.20	1.23	1.33
1	A	161	VAL	CA-CB	7.04	1.69	1.54
1	A	24	LYS	CE-NZ	-6.92	1.31	1.49
1	A	39	LYS	CE-NZ	-6.56	1.32	1.49
1	A	52	ASP	CG-OD1	6.53	1.40	1.25
1	A	45	LYS	C-O	6.20	1.35	1.23
1	A	46	PRO	CA-CB	6.04	1.65	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	114	TYR	CE2-CZ	-5.98	1.30	1.38
1	A	225	LYS	CD-CE	5.97	1.66	1.51
1	A	239[A]	GLU	CD-OE1	5.89	1.32	1.25
1	A	239[B]	GLU	CD-OE1	5.89	1.32	1.25
1	A	39	LYS	CA-CB	-5.75	1.41	1.53
1	A	75	ASP	C-N	-5.68	1.21	1.34
1	A	45	LYS	N-CA	5.61	1.57	1.46
1	A	27	ARG	CZ-NH2	-5.54	1.25	1.33
1	A	192	TRP	CZ3-CH2	-5.37	1.31	1.40
1	A	14[A]	GLU	CG-CD	5.23	1.59	1.51
1	A	14[B]	GLU	CG-CD	5.23	1.59	1.51
1	A	127	LYS	CB-CG	-5.22	1.38	1.52
1	A	159	LYS	CD-CE	5.21	1.64	1.51
1	A	236	GLU	CD-OE1	5.19	1.31	1.25
1	A	40	TYR	CE1-CZ	5.13	1.45	1.38
1	A	113	LYS	CE-NZ	-5.01	1.36	1.49

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ASP	CB-CG-OD2	-12.83	106.75	118.30
1	A	141	LEU	CB-CG-CD2	9.12	126.50	111.00
1	A	226	PHE	CB-CG-CD1	9.10	127.17	120.80
1	A	52	ASP	CB-CG-OD2	-8.92	110.28	118.30
1	A	175	ASP	CB-CG-OD1	8.69	126.12	118.30
1	A	14[A]	GLU	OE1-CD-OE2	8.47	133.46	123.30
1	A	14[B]	GLU	OE1-CD-OE2	8.47	133.46	123.30
1	A	52	ASP	CB-CG-OD1	8.12	125.61	118.30
1	A	47[A]	LEU	CA-CB-CG	7.92	133.52	115.30
1	A	47[B]	LEU	CA-CB-CG	7.92	133.52	115.30
1	A	37[A]	THR	OG1-CB-CG2	-7.63	92.46	110.00
1	A	37[B]	THR	OG1-CB-CG2	-7.63	92.46	110.00
1	A	239[A]	GLU	CG-CD-OE1	7.27	132.84	118.30
1	A	239[B]	GLU	CG-CD-OE1	7.27	132.84	118.30
1	A	89	ARG	CG-CD-NE	-7.18	96.72	111.80
1	A	226	PHE	CB-CG-CD2	-6.95	115.93	120.80
1	A	27	ARG	NE-CZ-NH2	-6.86	116.87	120.30
1	A	75	ASP	CB-CG-OD1	6.76	124.38	118.30
1	A	172	LYS	CD-CE-NZ	6.61	126.89	111.70
1	A	159	LYS	CG-CD-CE	6.36	130.99	111.90
1	A	27	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	47[A]	LEU	CB-CG-CD1	5.90	121.03	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47[B]	LEU	CB-CG-CD1	5.90	121.03	111.00
1	A	239[A]	GLU	CG-CD-OE2	-5.69	106.92	118.30
1	A	239[B]	GLU	CG-CD-OE2	-5.69	106.92	118.30
1	A	213	LYS	CD-CE-NZ	5.32	123.95	111.70
1	A	165	ASP	CB-CG-OD2	-5.30	113.53	118.30
1	A	114	TYR	CB-CG-CD1	5.29	124.17	121.00
1	A	130	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	14[A]	GLU	CG-CD-OE2	-5.13	108.03	118.30
1	A	14[B]	GLU	CG-CD-OE2	-5.13	108.03	118.30
1	A	80	LYS	CD-CE-NZ	5.09	123.41	111.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	234	GLU	Sidechain
1	A	75	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2150	2109	2093	12	0
2	A	6	8	7	4	0
3	A	2	0	0	0	0
4	A	279	0	0	2	0
All	All	2437	2117	2100	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:LEU:CB	1:A:100:LEU:CA	1.77	1.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301:GOL:C2	2:A:301:GOL:O2	1.64	1.44
1:A:100:LEU:CG	1:A:100:LEU:CA	2.58	0.80
1:A:100:LEU:CB	1:A:100:LEU:C	2.53	0.76
1:A:34[B]:ASP:OD2	1:A:37[B]:THR:HG22	1.96	0.65
2:A:301:GOL:O2	2:A:301:GOL:C3	2.47	0.62
2:A:301:GOL:C2	2:A:301:GOL:HO2	2.07	0.61
1:A:100:LEU:N	1:A:100:LEU:CB	2.60	0.59
2:A:301:GOL:H11	4:A:438:HOH:O	2.04	0.56
1:A:34[B]:ASP:HB3	1:A:37[B]:THR:CG2	2.45	0.47
1:A:161:VAL:CG1	1:A:225:LYS:HD2	2.44	0.47
1:A:89:ARG:HG3	1:A:125:THR:CG2	2.45	0.47
1:A:34[B]:ASP:HB3	1:A:37[B]:THR:HG22	1.98	0.45
1:A:134:ALA:O	1:A:140:GLY:HA3	2.17	0.43
1:A:47[A]:LEU:HG	1:A:191:TYR:CE2	2.56	0.40
1:A:64[B]:HIS:HD2	4:A:568:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/260 (104%)	261 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/225 (105%)	235 (99%)	2 (1%)	81	55

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	52	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	ASN
1	A	137	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	301	-	5,5,5	3.69	3 (60%)	5,5,5	2.26	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	GOL	O2-C2	6.93	1.64	1.43
2	A	301	GOL	C3-C2	2.94	1.63	1.51
2	A	301	GOL	C1-C2	-2.71	1.40	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	GOL	O1-C1-C2	-3.86	91.68	110.20
2	A	301	GOL	O2-C2-C3	-2.49	98.17	109.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/260 (98%)	0.25	8 (3%) 49 48	5, 9, 20, 50	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	LYS	4.5
1	A	40	TYR	3.4
1	A	46	PRO	3.4
1	A	42	PRO	3.3
1	A	261	LYS	3.2
1	A	4	HIS	2.7
1	A	43	SER	2.2
1	A	45	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	301	6/6	0.95	0.17	11,19,22,29	2
3	NI	A	303	1/1	0.98	0.18	30,30,30,30	0
3	NI	A	302	1/1	1.00	0.08	5,5,5,5	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.