



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 13, 2020 – 09:37 PM BST

PDB ID : 5LW0
Title : Oryza sativa APL macrodomain in complex with ADP-ribose
Authors : Ariza, A.
Deposited on : 2016-09-14
Resolution : 1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1931 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	231	 79% 8% 13%
1	B	231	 77% 11% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	B	302	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Basic helix-loop-helix, putative, expressed.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	12	0
			1609	1016	286	301	6			
1	B	204	Total	C	N	O	S	0	7	0
			1592	1003	282	301	6			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	66	MET	-	initiating methionine	UNP Q10MW4
A	67	GLY	-	expression tag	UNP Q10MW4
A	68	HIS	-	expression tag	UNP Q10MW4
A	69	HIS	-	expression tag	UNP Q10MW4
A	70	HIS	-	expression tag	UNP Q10MW4
A	71	HIS	-	expression tag	UNP Q10MW4
A	72	HIS	-	expression tag	UNP Q10MW4
A	73	HIS	-	expression tag	UNP Q10MW4
A	74	GLY	-	expression tag	UNP Q10MW4
A	75	GLY	-	expression tag	UNP Q10MW4
A	76	VAL	-	expression tag	UNP Q10MW4
A	77	LYS	-	expression tag	UNP Q10MW4
B	66	MET	-	initiating methionine	UNP Q10MW4
B	67	GLY	-	expression tag	UNP Q10MW4
B	68	HIS	-	expression tag	UNP Q10MW4
B	69	HIS	-	expression tag	UNP Q10MW4
B	70	HIS	-	expression tag	UNP Q10MW4
B	71	HIS	-	expression tag	UNP Q10MW4
B	72	HIS	-	expression tag	UNP Q10MW4
B	73	HIS	-	expression tag	UNP Q10MW4
B	74	GLY	-	expression tag	UNP Q10MW4
B	75	GLY	-	expression tag	UNP Q10MW4
B	76	VAL	-	expression tag	UNP Q10MW4
B	77	LYS	-	expression tag	UNP Q10MW4

- Molecule 2 is [(2R,3S,4R,5R)-5-(6-AMINOPURIN-9-YL)-3,4-DIHYDROXY-OXOLAN-2-YL]METHYL [HYDROXY-[(2R,3S,4R,5S)-3,4,5-TRIHYDROXYOXOLAN-2-YL]METHOXY]PHOSPHORYL] HYDROGEN PHOSPHATE (three-letter code: AR6) (formula: C₁₅H₂₃N₅O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is GLYCINE (three-letter code: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C N O 5 2 1 2	0	0
4	B	1	Total C N O 5 2 1 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	168	Total O 169 169	0	1
5	B	197	Total O 197 197	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Basic helix-loop-helix, putative, expressed



- Molecule 1: Basic helix-loop-helix, putative, expressed



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	73.50Å 91.32Å 62.88Å 90.00° 103.32° 90.00°	Depositor
Resolution (Å)	56.31 – 1.65	Depositor
% Data completeness (in resolution range)	97.2 (56.31-1.65)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.155 , 0.187	Depositor
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.504	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3664	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, AR6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	6/1673 (0.4%)	0.80	2/2265 (0.1%)
1	B	0.98	0/1640	0.73	2/2222 (0.1%)
All	All	1.03	6/3313 (0.2%)	0.77	4/4487 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	TYR	CZ-OH	7.45	1.50	1.37
1	A	135	TYR	CD1-CE1	6.91	1.49	1.39
1	A	135	TYR	CB-CG	6.45	1.61	1.51
1	A	135	TYR	CG-CD2	-5.69	1.31	1.39
1	A	135	TYR	CG-CD1	5.59	1.46	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	MET	CG-SD-CE	-10.06	84.11	100.20
1	A	168	ARG	NE-CZ-NH2	-7.81	116.40	120.30
1	B	168	ARG	NE-CZ-NH2	-7.28	116.66	120.30
1	B	261	MET	CG-SD-CE	-6.89	89.17	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1609	0	1659	17	0
1	B	1592	0	1629	23	0
2	A	36	0	21	1	0
2	B	36	0	21	3	0
3	A	10	0	0	1	0
3	B	5	0	0	2	0
4	B	10	0	4	4	0
5	A	169	0	0	14	0
5	B	197	0	0	7	0
All	All	3664	0	3334	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 42 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301:AR6:H1D	5:A:431:HOH:O	1.43	1.19
2:B:301:AR6:H1D	5:B:450:HOH:O	1.50	1.08
1:B:165:ASN:HD22	4:B:303:GLY:HA2	1.29	0.97
1:B:114:GLN:OE1	5:B:401:HOH:O	1.96	0.82
1:A:289[B]:GLU:OE2	5:A:401:HOH:O	2.05	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	303	-	4,4,4	0.64	0	6,6,6	0.58	0
2	AR6	A	301	-	34,39,39	1.33	4 (11%)	40,60,60	1.36	7 (17%)
3	PO4	B	302	-	4,4,4	0.82	0	6,6,6	0.49	0
4	GLY	B	304	-	1,4,4	0.11	0	0,4,4	0.00	-
2	AR6	B	301	-	34,39,39	1.54	4 (11%)	40,60,60	1.41	9 (22%)
4	GLY	B	303	-	1,4,4	0.10	0	0,4,4	0.00	-
3	PO4	A	302	-	4,4,4	1.90	1 (25%)	6,6,6	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AR6	A	301	-	-	3/18/54/54	0/4/4/4
4	GLY	B	304	-	-	0/0/2/2	-
2	AR6	B	301	-	-	2/18/54/54	0/4/4/4
4	GLY	B	303	-	-	0/0/2/2	-

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	AR6	O4'-C1'	6.04	1.49	1.41
2	A	301	AR6	O4'-C1'	3.62	1.46	1.41
3	A	302	PO4	P-O1	3.56	1.59	1.50
2	A	301	AR6	O4D-C1D	3.25	1.47	1.43
2	B	301	AR6	O4D-C1D	3.21	1.47	1.43

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	AR6	O4D-C1D-C2D	3.46	108.72	104.46
2	B	301	AR6	C4-C5-N7	-3.12	106.14	109.40
2	A	301	AR6	O4'-C1'-C2'	-3.03	102.50	106.93
2	A	301	AR6	N3-C2-N1	-2.90	124.15	128.68
2	B	301	AR6	C1'-N9-C4	-2.77	121.77	126.64

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	AR6	C5'-O5'-PA-O2A
2	B	301	AR6	C5'-O5'-PA-O2A
2	A	301	AR6	C5'-O5'-PA-O3A
2	B	301	AR6	C5'-O5'-PA-O3A
2	A	301	AR6	PB-O3A-PA-O1A


There are no ring outliers.


6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	PO4	1	0
2	A	301	AR6	1	0
3	B	302	PO4	2	0
4	B	304	GLY	2	0
2	B	301	AR6	3	0
4	B	303	GLY	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS failed to run properly - this section is therefore empty.


6.3 Carbohydrates ⓘ


EDS failed to run properly - this section is therefore empty.

6.4 Ligands ⓘ

EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.