



# Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 11:18 pm BST

PDB ID : 3LY5  
Title : DDX18 dead-domain  
Authors : Schutz, P.; Karlberg, T.; Arrowsmith, C.H.; Berglund, H.; Bountra, C.; Collins, R.; Edwards, A.M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Johansson, A.; Johansson, I.; Kallas, A.; Kraulis, P.; Kotenyova, T.; Kotzsch, A.; Markova, N.; Moche, M.; Nielsen, T.K.; Nordlund, P.; Nyman, T.; Persson, C.; Roos, A.K.; Siponen, M.I.; Svensson, L.; Thorsell, A.G.; Tresaugues, L.; Van Den Berg, S.; Wahlberg, E.; Weigelt, J.; Welin, M.; Wisniewska, M.; Schuler, H.M.; Structural Genomics Consortium (SGC)  
Deposited on : 2010-02-26  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	
1	B	262	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase DDX18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1518	958	264	287	9			
1	B	213	Total	C	N	O	S	0	0	0
			1542	977	262	294	9			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	126	MET	-	EXPRESSION TAG	UNP Q9NVP1
A	127	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	128	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	129	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	130	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	131	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	132	HIS	-	EXPRESSION TAG	UNP Q9NVP1
A	133	SER	-	EXPRESSION TAG	UNP Q9NVP1
A	134	SER	-	EXPRESSION TAG	UNP Q9NVP1
A	135	GLY	-	EXPRESSION TAG	UNP Q9NVP1
A	136	VAL	-	EXPRESSION TAG	UNP Q9NVP1
A	137	ASP	-	EXPRESSION TAG	UNP Q9NVP1
A	138	LEU	-	EXPRESSION TAG	UNP Q9NVP1
A	139	GLY	-	EXPRESSION TAG	UNP Q9NVP1
A	140	THR	-	EXPRESSION TAG	UNP Q9NVP1
A	141	GLU	-	EXPRESSION TAG	UNP Q9NVP1
A	142	ASN	-	EXPRESSION TAG	UNP Q9NVP1
A	143	LEU	-	EXPRESSION TAG	UNP Q9NVP1
A	144	TYR	-	EXPRESSION TAG	UNP Q9NVP1
A	145	PHE	-	EXPRESSION TAG	UNP Q9NVP1
A	146	GLN	-	EXPRESSION TAG	UNP Q9NVP1
A	147	SER	-	EXPRESSION TAG	UNP Q9NVP1
A	148	MET	-	EXPRESSION TAG	UNP Q9NVP1
B	126	MET	-	EXPRESSION TAG	UNP Q9NVP1
B	127	HIS	-	EXPRESSION TAG	UNP Q9NVP1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	128	HIS	-	EXPRESSION TAG	UNP Q9NVP1
B	129	HIS	-	EXPRESSION TAG	UNP Q9NVP1
B	130	HIS	-	EXPRESSION TAG	UNP Q9NVP1
B	131	HIS	-	EXPRESSION TAG	UNP Q9NVP1
B	132	HIS	-	EXPRESSION TAG	UNP Q9NVP1
B	133	SER	-	EXPRESSION TAG	UNP Q9NVP1
B	134	SER	-	EXPRESSION TAG	UNP Q9NVP1
B	135	GLY	-	EXPRESSION TAG	UNP Q9NVP1
B	136	VAL	-	EXPRESSION TAG	UNP Q9NVP1
B	137	ASP	-	EXPRESSION TAG	UNP Q9NVP1
B	138	LEU	-	EXPRESSION TAG	UNP Q9NVP1
B	139	GLY	-	EXPRESSION TAG	UNP Q9NVP1
B	140	THR	-	EXPRESSION TAG	UNP Q9NVP1
B	141	GLU	-	EXPRESSION TAG	UNP Q9NVP1
B	142	ASN	-	EXPRESSION TAG	UNP Q9NVP1
B	143	LEU	-	EXPRESSION TAG	UNP Q9NVP1
B	144	TYR	-	EXPRESSION TAG	UNP Q9NVP1
B	145	PHE	-	EXPRESSION TAG	UNP Q9NVP1
B	146	GLN	-	EXPRESSION TAG	UNP Q9NVP1
B	147	SER	-	EXPRESSION TAG	UNP Q9NVP1
B	148	MET	-	EXPRESSION TAG	UNP Q9NVP1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	S	0	0
			3	1	1	1		
4	B	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	25	Total	O	0	0
			25	25		
5	B	20	Total	O	0	0
			20	20		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-dependent RNA helicase DDX18

Chain A: 



- Molecule 1: ATP-dependent RNA helicase DDX18

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.34Å 41.34Å 230.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.42 – 2.80 38.42 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.42-2.80) 100.0 (38.42-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.21	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.246 , 0.274 0.244 , 0.273	Depositor DCC
$R_{free}$ test set	491 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.2	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 32.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.55$ , $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.330 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	3118	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, SCN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/1537	0.49	0/2079
1	B	0.35	0/1563	0.52	0/2120
All	All	0.35	0/3100	0.50	0/4199

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1518	0	1451	18	0
1	B	1542	0	1470	24	0
2	A	5	0	0	0	0
3	A	2	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	A	25	0	0	0	0
5	B	20	0	0	0	0
All	All	3118	0	2921	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.



All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ASP:O	1:A:383:PRO:HD2	1.79	0.82
1:A:205:ILE:O	1:A:209:SER:HB3	1.88	0.74
1:B:346:GLU:O	1:B:350:ILE:HG23	1.93	0.67
1:B:261:ARG:O	1:B:265:MET:HB2	1.97	0.65
1:A:252:THR:HG22	1:A:324:TYR:HA	1.79	0.64
1:B:217:ARG:HH21	1:B:383:PRO:HB3	1.61	0.64
1:B:202:MET:HE3	1:B:210:ILE:HD12	1.79	0.64
1:B:179:PHE:CB	1:B:185:LEU:HA	2.28	0.64
1:B:235:ILE:HB	1:B:236:PRO:HD3	1.80	0.64
1:A:337:ARG:O	1:A:341:VAL:HG23	1.98	0.63
1:B:263:LEU:CD2	1:B:334:GLU:HG3	2.31	0.61
1:A:327:LEU:HD22	1:A:354:LEU:HD22	1.82	0.60
1:B:255:LEU:HD21	1:B:312:LEU:HD21	1.84	0.59
1:B:327:LEU:HD22	1:B:354:LEU:HD22	1.87	0.56
1:B:218:ASP:O	1:B:383:PRO:HD2	2.08	0.54
1:B:222:ALA:HB3	1:B:386:VAL:HA	1.90	0.54
1:B:217:ARG:NH2	1:B:383:PRO:HB3	2.24	0.53
1:A:347:LEU:HD13	1:A:374:LEU:HD13	1.90	0.53
1:B:287:GLY:HA2	1:B:311:ARG:NH1	2.25	0.51
1:A:263:LEU:HD22	1:A:334:GLU:HG3	1.94	0.50
1:A:335:ALA:HB3	1:A:364:SER:HB2	1.94	0.50
1:A:268:PHE:HB2	1:A:284:LEU:HD13	1.93	0.50
1:A:202:MET:HE2	1:A:206:GLN:HB3	1.94	0.48
1:A:229:LYS:NZ	1:A:334:GLU:OE2	2.44	0.48
1:B:205:ILE:HG21	1:B:227:SER:HB3	1.96	0.46
1:A:254:VAL:HB	1:A:304:ILE:HG23	1.98	0.46
1:B:230:THR:HG23	1:B:267:THR:HG23	1.98	0.46
1:B:285:ILE:O	1:B:307:ALA:HA	2.16	0.46
1:B:217:ARG:HB3	1:B:383:PRO:HG3	1.98	0.46
1:A:263:LEU:CD2	1:A:334:GLU:HG3	2.48	0.44
1:A:230:THR:HA	1:A:233:PHE:CE1	2.53	0.43
1:B:230:THR:HA	1:B:233:PHE:CE1	2.54	0.43
1:B:237:ALA:HB1	1:B:254:VAL:HG11	2.01	0.43
1:A:173:ALA:HB3	1:A:176:ASP:OD1	2.18	0.42
1:A:179:PHE:N	1:A:185:LEU:N	2.66	0.42
1:A:202:MET:HE1	1:A:207:HIS:HA	2.01	0.42
1:B:350:ILE:CG1	1:B:351:ILE:N	2.82	0.42
1:B:202:MET:HE2	1:B:206:GLN:HB3	2.02	0.41
1:A:259:PRO:HD2	1:A:263:LEU:HD23	2.02	0.41
1:B:285:ILE:HB	1:B:307:ALA:HB2	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:MET:HB2	1:B:199:PHE:HD1	1.86	0.41
1:B:259:PRO:HD2	1:B:263:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/262 (79%)	198 (96%)	9 (4%)	0	100	100
1	B	209/262 (80%)	199 (95%)	10 (5%)	0	100	100
All	All	416/524 (79%)	397 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/227 (65%)	139 (95%)	8 (5%)	22	53
1	B	149/227 (66%)	145 (97%)	4 (3%)	44	78
All	All	296/454 (65%)	284 (96%)	12 (4%)	30	64

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	176	ASP
1	A	186	VAL
1	A	209	SER
1	A	224	LYS
1	A	252	THR
1	A	341	VAL
1	A	356	THR
1	A	374	LEU
1	B	177	THR
1	B	311	ARG
1	B	334	GLU
1	B	357	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SCN	B	2	-	1,2,2	1.02	0	0,1,1	0.00	-
4	SCN	A	390	-	1,2,2	1.06	0	0,1,1	0.00	-
2	PO4	A	1	-	4,4,4	0.91	0	6,6,6	0.62	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/262 (80%)	-0.49	0 100 100	37, 46, 55, 57	0
1	B	213/262 (81%)	-0.44	0 100 100	38, 48, 55, 61	0
All	All	424/524 (80%)	-0.46	0 100 100	37, 46, 55, 61	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SCN	B	2	3/3	0.82	0.25	55,55,55,56	0
3	MG	A	388	1/1	0.85	0.18	55,55,55,55	0
4	SCN	A	390	3/3	0.92	0.19	48,48,48,48	0
3	MG	A	389	1/1	0.97	0.11	48,48,48,48	0
2	PO4	A	1	5/5	0.99	0.17	14,15,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.