



# Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 11:24 pm BST

PDB ID : 3LZ7  
Title : Crystal Structure of thioesterase HI1161 EC3.1.2.- from Haemophilus influenzae. Orthorombic crystal form. Northeast Structural Genomics Consortium Target IR63  
Authors : Kuzin, A.; Min, S.; Seetharaman, J.; Janjua, J.; Xiao, R.; Foote, E.L.; Ciccosanti, C.; Cooper, B.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2010-03-01  
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	<div> <div>86%</div> <div>12%</div> <div>.</div> </div>
1	B	146	<div> <div>2%</div> <div>79%</div> <div>14%</div> <div>.</div> <div>.</div> </div>
1	C	146	<div> <div>3%</div> <div>81%</div> <div>16%</div> <div>.</div> <div>.</div> </div>
1	D	146	<div> <div>7%</div> <div>77%</div> <div>19%</div> <div>.</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative esterase HI1161.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	S	Se	0	1	0
			1111	696	203	205	4	3			
1	B	140	Total	C	N	O	S	Se	0	1	0
			1079	677	191	204	4	3			
1	C	143	Total	C	N	O	S	Se	0	1	0
			1106	693	200	205	5	3			
1	D	141	Total	C	N	O	S	Se	0	0	0
			1080	678	193	202	4	3			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	139	LEU	-	expression tag	UNP P45083
A	140	GLU	-	expression tag	UNP P45083
A	141	HIS	-	expression tag	UNP P45083
A	142	HIS	-	expression tag	UNP P45083
A	143	HIS	-	expression tag	UNP P45083
A	144	HIS	-	expression tag	UNP P45083
A	145	HIS	-	expression tag	UNP P45083
A	146	HIS	-	expression tag	UNP P45083
B	139	LEU	-	expression tag	UNP P45083
B	140	GLU	-	expression tag	UNP P45083
B	141	HIS	-	expression tag	UNP P45083
B	142	HIS	-	expression tag	UNP P45083
B	143	HIS	-	expression tag	UNP P45083
B	144	HIS	-	expression tag	UNP P45083
B	145	HIS	-	expression tag	UNP P45083
B	146	HIS	-	expression tag	UNP P45083
C	139	LEU	-	expression tag	UNP P45083
C	140	GLU	-	expression tag	UNP P45083
C	141	HIS	-	expression tag	UNP P45083
C	142	HIS	-	expression tag	UNP P45083
C	143	HIS	-	expression tag	UNP P45083

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	144	HIS	-	expression tag	UNP P45083
C	145	HIS	-	expression tag	UNP P45083
C	146	HIS	-	expression tag	UNP P45083
D	139	LEU	-	expression tag	UNP P45083
D	140	GLU	-	expression tag	UNP P45083
D	141	HIS	-	expression tag	UNP P45083
D	142	HIS	-	expression tag	UNP P45083
D	143	HIS	-	expression tag	UNP P45083
D	144	HIS	-	expression tag	UNP P45083
D	145	HIS	-	expression tag	UNP P45083
D	146	HIS	-	expression tag	UNP P45083

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total	O	0	0
			88	88		
3	B	62	Total	O	0	0
			62	62		
3	C	54	Total	O	0	0
			54	54		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	35	Total	O	0	0
			35	35		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative esterase HI1161

Chain A: 




- Molecule 1: Putative esterase HI1161

Chain B: 




- Molecule 1: Putative esterase HI1161

Chain C: 



- Molecule 1: Putative esterase HI1161

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.53Å 71.19Å 112.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.04 – 2.19 37.04 – 2.19	Depositor EDS
% Data completeness (in resolution range)	98.7 (37.04-2.19) 98.7 (37.04-2.19)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.27 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.5_2, REFMAC	Depositor
R, $R_{free}$	0.168 , 0.231 0.163 , 0.221	Depositor DCC
$R_{free}$ test set	1439 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4622	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/1128	0.55	0/1527
1	B	0.37	0/1093	0.54	0/1480
1	C	0.40	0/1123	0.54	0/1521
1	D	0.33	0/1095	0.53	0/1483
All	All	0.37	0/4439	0.54	0/6011

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1111	0	1127	15	0
1	B	1079	0	1099	16	0
1	C	1106	0	1119	20	0
1	D	1080	0	1101	25	0
2	A	7	0	10	2	0
3	A	88	0	0	2	0
3	B	62	0	0	2	0
3	C	54	0	0	2	0
3	D	35	0	0	1	0
All	All	4622	0	4456	72	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MSE:HE2	1:B:33:GLU:HG2	1.45	0.96
1:C:142:HIS:N	1:C:143:HIS:HA	1.89	0.88
1:A:52:VAL:HG12	1:A:94:ARG:HG2	1.66	0.76
1:A:44:HIS:HA	1:A:47:MSE:HE2	1.68	0.75
1:D:112:GLN:NE2	1:D:114:TRP:HE1	1.83	0.75
1:D:44:HIS:HA	1:D:47:MSE:HE2	1.69	0.75
1:D:112:GLN:HE21	1:D:114:TRP:HE1	1.34	0.73
1:B:84:ASP:OD1	1:B:132:THR:HB	1.88	0.72
1:D:1:MSE:HE2	1:D:3:TRP:HD1	1.54	0.72
1:C:141:HIS:HB3	1:C:142:HIS:HB3	1.73	0.71
1:D:137:ASN:H	1:D:141:HIS:CE1	2.09	0.70
1:B:139:LEU:O	1:B:140:GLU:HB2	1.91	0.70
1:C:140:GLU:CD	1:C:143:HIS:CE1	2.69	0.66
1:D:1:MSE:HG2	1:D:2:LEU:H	1.61	0.65
1:A:44:HIS:HD2	3:B:183:HOH:O	1.79	0.65
1:C:43:ASP:O	1:C:47:MSE:HG3	1.96	0.64
1:A:111:ILE:HB	2:A:147:PEG:H31	1.80	0.64
1:A:112:GLN:NE2	1:A:114:TRP:HE1	1.98	0.62
1:B:120:THR:OG1	1:B:122[B]:GLU:HG2	2.00	0.61
1:C:142:HIS:CD2	1:C:142:HIS:O	2.55	0.60
1:B:75:GLU:HG3	3:B:158:HOH:O	2.03	0.58
1:B:112:GLN:NE2	1:B:114:TRP:HE1	2.01	0.58
1:D:70:GLY:HA2	1:D:112:GLN:HE22	1.68	0.58
1:B:112:GLN:HE21	1:B:114:TRP:HE1	1.52	0.58
1:D:81:VAL:HG23	1:D:136:ILE:HD13	1.87	0.56
1:D:1:MSE:HG2	1:D:2:LEU:N	2.22	0.55
1:C:112:GLN:NE2	1:C:114:TRP:HE1	2.05	0.53
1:C:140:GLU:HG2	3:C:199:HOH:O	2.08	0.53
1:C:91:ARG:HG2	1:C:126:CYS:HA	1.91	0.52
1:C:60:ALA:HA	1:D:56:GLY:HA3	1.91	0.52
1:A:112:GLN:HE21	1:A:114:TRP:HE1	1.58	0.52
1:D:1:MSE:N	1:D:1:MSE:SE	2.92	0.52
1:A:59:VAL:HG11	1:B:59:VAL:HG11	1.92	0.51
1:C:56:GLY:HA3	1:D:60:ALA:HA	1.93	0.50
1:D:115:GLN:HG3	3:D:162:HOH:O	2.12	0.50
1:D:52:VAL:HG12	1:D:94:ARG:HG2	1.94	0.50
1:A:137:ASN:HB3	1:A:140:GLU:HG3	1.94	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:70:GLY:CA	1:D:112:GLN:HE22	2.25	0.49
1:A:44:HIS:CA	1:A:47:MSE:HE2	2.40	0.49
1:A:70:GLY:CA	1:A:112:GLN:HE22	2.26	0.48
1:C:38:ALA:HB2	1:C:65:ILE:HD13	1.95	0.48
1:B:84:ASP:OD1	1:B:132:THR:CB	2.62	0.47
1:C:59:VAL:HG11	1:D:59:VAL:HG11	1.96	0.47
1:C:142:HIS:O	1:C:142:HIS:HD2	1.97	0.46
1:D:7:PHE:CD2	1:D:72:LEU:HD22	2.51	0.46
1:C:112:GLN:HE21	1:C:114:TRP:HE1	1.62	0.45
1:B:38:ALA:HB2	1:B:65:ILE:HD13	1.99	0.45
2:A:147:PEG:H31	2:A:147:PEG:H12	1.46	0.45
1:C:83:LEU:HD13	1:C:134:SER:HB2	1.99	0.45
1:B:70:GLY:HA2	1:B:112:GLN:HE22	1.82	0.44
1:B:70:GLY:CA	1:B:112:GLN:HE22	2.30	0.44
1:B:83:LEU:HD13	1:B:134:SER:HB2	1.99	0.44
1:D:44:HIS:CA	1:D:47:MSE:HE2	2.43	0.43
1:B:4:LYS:HG3	1:B:73:CYS:O	2.19	0.43
1:C:70:GLY:HA2	1:C:112:GLN:HE22	1.84	0.43
1:A:48:GLN:NE2	3:A:201:HOH:O	2.53	0.42
3:C:214:HOH:O	1:D:47:MSE:HE3	2.19	0.42
1:C:2:LEU:HD22	1:C:73[A]:CYS:SG	2.60	0.41
1:D:79:THR:HG23	1:D:136:ILE:HD11	2.02	0.41
1:D:1:MSE:CG	1:D:2:LEU:H	2.27	0.41
1:D:47:MSE:HB2	1:D:47:MSE:HE3	1.94	0.41
1:A:118:ILE:HB	1:A:127:CYS:HB3	2.03	0.41
1:B:139:LEU:O	1:B:140:GLU:CB	2.60	0.41
1:C:131:LEU:HD21	1:C:133:LEU:HD21	2.02	0.41
1:A:43:ASP:O	1:A:47:MSE:HG3	2.21	0.41
1:B:2:LEU:HG	1:B:73:CYS:SG	2.61	0.40
1:D:41:PRO:HG3	1:D:45:ARG:NH2	2.36	0.40
1:A:70:GLY:HA2	1:A:112:GLN:HE22	1.86	0.40
1:C:70:GLY:CA	1:C:112:GLN:HE22	2.35	0.40
1:A:47:MSE:HE3	3:A:215:HOH:O	2.21	0.40
1:C:23:HIS:CE1	1:D:23:HIS:CE1	3.09	0.40
1:D:139:LEU:HD12	1:D:139:LEU:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/146 (97%)	140 (99%)	2 (1%)	0	100	100
1	B	139/146 (95%)	137 (99%)	2 (1%)	0	100	100
1	C	142/146 (97%)	139 (98%)	3 (2%)	0	100	100
1	D	139/146 (95%)	135 (97%)	4 (3%)	0	100	100
All	All	562/584 (96%)	551 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/123 (101%)	122 (98%)	2 (2%)	62	76
1	B	121/123 (98%)	113 (93%)	8 (7%)	16	19
1	C	124/123 (101%)	120 (97%)	4 (3%)	39	50
1	D	121/123 (98%)	116 (96%)	5 (4%)	30	39
All	All	490/492 (100%)	471 (96%)	19 (4%)	32	41

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	123	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	6	THR
1	B	14	GLN
1	B	33	GLU
1	B	48	GLN
1	B	83	LEU
1	B	109	ARG
1	B	139	LEU
1	B	140	GLU
1	C	2	LEU
1	C	48	GLN
1	C	83	LEU
1	C	94	ARG
1	D	1	MSE
1	D	6	THR
1	D	14	GLN
1	D	83	LEU
1	D	92	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	44	HIS
1	A	48	GLN
1	A	112	GLN
1	A	123	ASN
1	B	11	ASN
1	B	48	GLN
1	B	89	HIS
1	B	112	GLN
1	C	11	ASN
1	C	14	GLN
1	C	48	GLN
1	C	89	HIS
1	C	112	GLN
1	C	142	HIS
1	C	143	HIS
1	D	11	ASN
1	D	44	HIS
1	D	89	HIS
1	D	112	GLN
1	D	115	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	141	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PEG	A	147	-	6,6,6	0.75	0	5,5,5	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	147	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	147	PEG	C1-C2-O2-C3
2	A	147	PEG	O1-C1-C2-O2
2	A	147	PEG	O2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	147	PEG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/146 (95%)	-0.33	0 100 100	13, 22, 45, 57	0
1	B	137/146 (93%)	-0.30	3 (2%) 62 59	13, 23, 49, 86	0
1	C	140/146 (95%)	-0.24	4 (2%) 51 49	11, 24, 54, 92	0
1	D	138/146 (94%)	0.08	10 (7%) 15 14	12, 33, 67, 140	0
All	All	555/584 (95%)	-0.20	17 (3%) 49 47	11, 25, 57, 140	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	143	HIS	5.8
1	D	141	HIS	5.3
1	C	139	LEU	4.5
1	D	139	LEU	3.6
1	D	50	PHE	3.3
1	D	140	GLU	3.0
1	C	141	HIS	2.8
1	D	138	LEU	2.7
1	B	140	GLU	2.4
1	D	7	PHE	2.4
1	C	50	PHE	2.3
1	B	6	THR	2.3
1	D	109	ARG	2.3
1	D	9	LEU	2.2
1	D	76	GLU	2.1
1	D	77	GLY	2.1
1	B	139	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PEG	A	147	7/7	0.95	0.16	20,33,48,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.