



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:18 pm BST

PDB ID : 5M65
Title : Crystal structure of S-adenosyl-L-homocysteine hydrolase from *Bradyrhizobium elkanii* in complex with adenine
Authors : Manszewski, T.; Jaskolski, M.
Deposited on : 2016-10-24
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div></div> </div> </div>
1	B	479	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div></div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EDO	A	506	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylhomocysteinase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	469	Total	C	N	O	S	0	6	0
			3659	2318	627	692	22			
1	B	468	Total	C	N	O	S	0	3	0
			3632	2301	621	686	24			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is ADENINE (three-letter code: ADE) (formula: $C_5H_5N_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			10	5	5		
3	B	1	Total	C	N	0	0
			10	5	5		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Br	0	0
			1	1		
5	A	1	Total	Br	0	0
			1	1		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	282	Total	O	0	0
			282	282		
7	B	288	Total	O	0	0
			288	288		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenosylhomocysteinase



• Molecule 1: Adenosylhomocysteinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	100.67Å 103.05Å 90.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.87 – 1.95 45.87 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.87-1.95) 99.1 (45.87-1.94)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.177 , 0.212 0.177 , 0.210	Depositor DCC
R_{free} test set	1009 reflections (1.46%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7993	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BR, ADE, EDO, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/3745	0.56	0/5055
1	B	0.41	1/3708 (0.0%)	0.59	2/5006 (0.0%)
All	All	0.39	1/7453 (0.0%)	0.57	2/10061 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	181	LYS	CB-CG	6.27	1.69	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	181	LYS	CB-CG-CD	7.55	131.22	111.60
1	B	181	LYS	CD-CE-NZ	5.59	124.55	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3659	0	3678	42	0
1	B	3632	0	3646	45	0
2	A	44	0	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	0	26	0	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	8	0	11	4	0
6	B	12	0	18	1	0
7	A	282	0	0	4	0
7	B	288	0	0	2	0
All	All	7993	0	7413	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181:LYS:HG2	1:B:182:GLY:N	1.96	0.81
1:A:471:TYR:HA	6:A:506:EDO:H12	1.71	0.70
1:B:258[A]:MET:HE2	1:B:281:MET:HB2	1.74	0.67
1:B:164:GLU:HG2	7:B:769:HOH:O	1.95	0.66
1:B:28:ALA:HB1	1:B:32[A]:MET:HE3	1.77	0.65
1:B:162:SER:HB3	1:B:165:GLU:HB3	1.79	0.64
1:B:328:MET:HG2	1:B:331[A]:MET:HE1	1.81	0.63
1:A:303:MET:HE3	1:A:306:ALA:HB3	1.80	0.63
1:A:435:VAL:O	1:A:439:HIS:HD2	1.82	0.61
1:B:258[A]:MET:CE	1:B:281:MET:HB2	2.29	0.61
1:B:115:ASP:OD1	1:B:174:ARG:NH1	2.28	0.60
1:A:222:ASN:HA	1:A:227:LYS:HD2	1.85	0.59
1:A:222:ASN:HD21	1:A:232:ASN:HD21	1.53	0.56
1:A:144:HIS:CE1	1:A:215:LEU:H	2.23	0.56
1:A:258[A]:MET:HG2	1:A:303:MET:HE1	1.88	0.56
1:B:324:THR:HA	1:B:348:GLN:HG3	1.88	0.55
1:B:222:ASN:HD22	1:B:227:LYS:HZ2	1.55	0.55
1:A:303:MET:HA	1:A:303:MET:HE3	1.89	0.54
1:B:258[A]:MET:HE3	1:B:306:ALA:HB2	1.90	0.53
1:B:25:ILE:O	1:B:29:GLU:HG3	2.10	0.52
6:A:506:EDO:H21	1:B:223:ASP:CG	2.30	0.51
1:A:325:ILE:O	1:A:329:ARG:HG3	2.09	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:HIS:HE1	1:A:215:LEU:H	1.59	0.51
1:B:177:LYS:N	1:B:177:LYS:HD3	2.26	0.51
1:A:144:HIS:HE1	1:A:215:LEU:N	2.09	0.50
1:A:237:ARG:HA	1:A:272:SER:HB2	1.92	0.50
1:B:242:ASP:O	1:B:246:ARG:HG3	2.11	0.50
1:A:110:LEU:HD13	1:A:164:GLU:HG2	1.95	0.49
1:B:20:PHE:CD1	6:B:507:EDO:H12	2.48	0.49
1:B:325:ILE:O	1:B:329:ARG:HG3	2.13	0.49
1:A:179:LYS:HD2	1:A:183:TRP:CE2	2.47	0.48
1:B:319:ASN:HB3	1:B:322:ILE:HD11	1.95	0.48
1:A:258[A]:MET:HE3	1:A:306:ALA:HB2	1.94	0.48
1:A:173:LYS:O	7:A:601:HOH:O	2.19	0.47
1:A:337:VAL:HG12	1:A:347:ILE:HD13	1.95	0.47
1:A:303:MET:CE	1:A:306:ALA:HB3	2.44	0.47
1:A:471:TYR:CA	6:A:506:EDO:H12	2.42	0.47
1:B:258[A]:MET:HE1	1:B:301:VAL:HG22	1.97	0.47
1:A:24[B]:GLU:OE1	1:A:90:HIS:HE1	1.97	0.46
1:B:222:ASN:HD21	1:B:232:ASN:HD21	1.64	0.46
1:B:222:ASN:O	1:B:228:SER:HB3	2.16	0.46
1:B:222:ASN:HD22	1:B:227:LYS:NZ	2.12	0.46
1:A:258[A]:MET:HE2	1:A:258[A]:MET:HB2	1.82	0.45
1:A:446:LYS:HD2	7:A:660:HOH:O	2.16	0.45
1:A:207:GLU:HG3	1:A:211:LYS:HE3	1.98	0.45
1:B:226:THR:HA	1:B:230:PHE:CD2	2.52	0.45
1:A:222:ASN:HD22	1:A:227:LYS:HZ2	1.65	0.45
1:A:122:ASP:OD1	1:A:179:LYS:NZ	2.35	0.45
1:A:380:GLU:HG3	7:A:799:HOH:O	2.17	0.45
1:B:61:ILE:H	1:B:88:GLN:NE2	2.15	0.45
1:B:193:GLY:HA3	1:B:411:LEU:HD13	1.99	0.44
1:A:229:LYS:HE2	7:B:696:HOH:O	2.16	0.44
1:B:32[A]:MET:HE1	1:B:62:GLN:HA	1.99	0.44
1:B:32[A]:MET:CE	1:B:62:GLN:HA	2.47	0.44
1:A:352:LEU:HD13	1:A:367:ILE:HD13	2.00	0.44
1:A:144:HIS:CE1	1:A:215:LEU:N	2.85	0.44
1:B:55:GLY:HA3	1:B:79:TRP:CE3	2.53	0.44
1:A:398:SER:OG	1:A:439:HIS:HE1	2.01	0.43
1:A:42:TYR:O	1:A:47:PRO:HD3	2.17	0.43
1:B:28:ALA:O	1:B:32[B]:MET:HG3	2.18	0.43
1:B:403:ASN:HD22	1:B:427:LEU:HD12	1.83	0.43
1:A:258[B]:MET:HG2	1:A:303:MET:HE1	1.99	0.43
1:B:62:GLN:CD	1:B:62:GLN:H	2.21	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ASN:C	1:B:220:ASN:HD22	2.22	0.43
1:A:462:GLN:HG3	7:A:677:HOH:O	2.19	0.43
1:B:372:LYS:HE2	1:B:372:LYS:HA	2.01	0.43
1:B:258[A]:MET:HE3	1:B:306:ALA:CB	2.49	0.43
1:A:155:ALA:HB1	1:A:159:LYS:HZ1	1.84	0.42
1:B:331[A]:MET:HE3	1:B:375:ILE:HG21	2.02	0.42
1:B:88:GLN:HB3	1:B:90:HIS:CE1	2.54	0.42
1:A:221:VAL:HG13	1:A:403:ASN:HB3	2.01	0.42
1:B:328:MET:HA	1:B:331[A]:MET:HE2	2.00	0.42
2:A:501:NAD:H2D	2:A:501:NAD:H6N	1.87	0.42
1:A:207:GLU:OE2	1:A:207:GLU:HA	2.19	0.42
6:A:506:EDO:H21	1:B:223:ASP:OD2	2.19	0.42
1:A:62:GLN:HG3	1:A:394:SER:HA	2.02	0.42
1:B:222:ASN:HA	1:B:227:LYS:HD2	2.02	0.42
1:A:62:GLN:CD	1:A:62:GLN:H	2.23	0.41
1:B:58:HIS:CE1	1:B:82:CYS:SG	3.13	0.41
1:B:339:ASN:ND2	1:B:341:GLY:H	2.18	0.41
1:B:402:THR:O	1:B:406:LEU:HG	2.20	0.41
1:B:122:ASP:OD1	1:B:179:LYS:NZ	2.48	0.41
1:A:155:ALA:HB1	1:A:159:LYS:NZ	2.35	0.41
1:A:339:ASN:ND2	1:A:341:GLY:H	2.19	0.41
1:A:186:GLU:HA	1:A:189:LYS:HG2	2.03	0.41
1:B:114:TRP:HB3	1:B:171:LEU:HD22	2.03	0.40
1:B:225:VAL:HG21	1:B:473:TYR:CE1	2.57	0.40
1:A:398:SER:CB	1:A:439:HIS:HE1	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	473/479 (99%)	458 (97%)	14 (3%)	1 (0%)	47 39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	469/479 (98%)	455 (97%)	13 (3%)	1 (0%)	47	39
All	All	942/958 (98%)	913 (97%)	27 (3%)	2 (0%)	47	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	PRO
1	B	58	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/387 (100%)	382 (99%)	3 (1%)	81	78
1	B	381/387 (98%)	376 (99%)	5 (1%)	69	62
All	All	766/774 (99%)	758 (99%)	8 (1%)	73	71

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	HIS
1	A	216	PHE
1	A	468	SER
1	B	90	HIS
1	B	177	LYS
1	B	211	LYS
1	B	220	ASN
1	B	353	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	144	HIS
1	A	222	ASN
1	A	339	ASN
1	A	392	HIS
1	A	404	GLN
1	A	408	GLN
1	A	439	HIS
1	B	88	GLN
1	B	220	ASN
1	B	222	ASN
1	B	339	ASN
1	B	348	GLN
1	B	359	ASN
1	B	403	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	505	-	3,3,3	0.95	0	2,2,2	1.40	0
6	EDO	B	507	-	3,3,3	0.66	0	2,2,2	0.62	0
6	EDO	B	506	-	3,3,3	0.74	0	2,2,2	0.21	0
2	NAD	B	501	-	42,48,48	1.87	10 (23%)	50,73,73	1.29	6 (12%)
6	EDO	A	506	-	3,3,3	0.42	0	2,2,2	0.34	0
3	ADE	B	502	-	9,11,11	1.01	1 (11%)	7,15,15	1.11	1 (14%)
3	ADE	A	502	-	9,11,11	0.95	1 (11%)	7,15,15	1.13	1 (14%)
6	EDO	B	505	-	3,3,3	0.80	0	2,2,2	0.18	0
2	NAD	A	501	-	42,48,48	1.91	12 (28%)	50,73,73	1.23	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	505	-	-	0/1/1/1	-
6	EDO	B	507	-	-	1/1/1/1	-
6	EDO	B	506	-	-	0/1/1/1	-
2	NAD	B	501	-	-	5/26/62/62	0/5/5/5
6	EDO	A	506	-	-	0/1/1/1	-
3	ADE	B	502	-	-	-	0/2/2/2
3	ADE	A	502	-	-	-	0/2/2/2
6	EDO	B	505	-	-	1/1/1/1	-
2	NAD	A	501	-	-	5/26/62/62	0/5/5/5

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAD	C2B-C3B	-5.01	1.39	1.53
2	A	501	NAD	C2B-C3B	-4.43	1.41	1.53
2	A	501	NAD	C7N-N7N	4.23	1.41	1.33
2	B	501	NAD	C7N-N7N	4.03	1.40	1.33
2	B	501	NAD	C2B-C1B	-3.51	1.48	1.53
2	A	501	NAD	C6A-N6A	3.42	1.46	1.34
2	B	501	NAD	C6A-N6A	3.37	1.46	1.34
2	A	501	NAD	C5N-C4N	3.32	1.45	1.38
2	B	501	NAD	C5N-C4N	3.26	1.45	1.38
2	A	501	NAD	O4B-C4B	-3.17	1.37	1.45
2	B	501	NAD	C2D-C1D	-3.09	1.49	1.53
2	A	501	NAD	C2D-C1D	-2.98	1.49	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NAD	C2B-C1B	-2.94	1.49	1.53
2	B	501	NAD	O4B-C4B	-2.89	1.38	1.45
2	B	501	NAD	C2N-N1N	2.69	1.38	1.35
2	A	501	NAD	C2N-N1N	2.67	1.38	1.35
2	A	501	NAD	O4B-C1B	-2.41	1.37	1.41
2	A	501	NAD	C4N-C3N	2.39	1.43	1.39
2	A	501	NAD	O3D-C3D	-2.39	1.37	1.43
2	A	501	NAD	C3D-C4D	-2.28	1.47	1.53
3	B	502	ADE	C4-N9	2.19	1.38	1.34
3	A	502	ADE	C4-N9	2.17	1.38	1.34
2	B	501	NAD	C3D-C4D	-2.10	1.47	1.53
2	B	501	NAD	O4B-C1B	-2.01	1.38	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAD	N3A-C2A-N1A	-4.43	121.76	128.68
2	B	501	NAD	N3A-C2A-N1A	-4.42	121.77	128.68
2	B	501	NAD	PN-O3-PA	-2.79	123.25	132.83
2	A	501	NAD	PN-O3-PA	-2.75	123.38	132.83
2	B	501	NAD	O7N-C7N-N7N	-2.69	118.76	122.58
2	B	501	NAD	C4A-C5A-N7A	-2.47	106.82	109.40
2	B	501	NAD	C3N-C7N-N7N	2.39	120.62	117.75
2	A	501	NAD	O4B-C1B-C2B	-2.21	103.69	106.93
3	A	502	ADE	C5-C6-N6	2.15	123.61	120.35
2	A	501	NAD	C4A-C5A-N7A	-2.08	107.23	109.40
3	B	502	ADE	C5-C6-N6	2.08	123.51	120.35
2	B	501	NAD	O4B-C1B-C2B	-2.07	103.90	106.93

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	NAD	O4D-C1D-N1N-C2N
2	B	501	NAD	O4D-C1D-N1N-C6N
2	B	501	NAD	C2D-C1D-N1N-C2N
2	B	501	NAD	C2D-C1D-N1N-C6N
2	A	501	NAD	O4D-C1D-N1N-C2N
2	A	501	NAD	O4D-C1D-N1N-C6N
2	A	501	NAD	C2D-C1D-N1N-C2N
2	A	501	NAD	C2D-C1D-N1N-C6N
6	B	507	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	B	505	EDO	O1-C1-C2-O2
2	A	501	NAD	O4B-C4B-C5B-O5B
2	B	501	NAD	O4B-C4B-C5B-O5B


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	507	EDO	1	0
6	A	506	EDO	4	0
2	A	501	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	469/479 (97%)	0.06	23 (4%)	29 37	13, 28, 59, 88	1 (0%)
1	B	468/479 (97%)	-0.07	11 (2%)	59 66	11, 27, 48, 81	0
All	All	937/958 (97%)	-0.01	34 (3%)	42 50	11, 27, 54, 88	1 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	151	GLN	5.2
1	A	156	PHE	4.8
1	A	154	THR	4.6
1	A	160	PRO	4.5
1	A	157	LEU	4.4
1	B	181	LYS	3.9
1	A	176	LEU	3.8
1	B	416	LYS	3.7
1	B	182	GLY	3.4
1	A	147	TYR	3.4
1	A	159	LYS	3.3
1	A	185	ALA	3.0
1	A	177	LYS	3.0
1	B	164	GLU	3.0
1	A	149	ALA	2.8
1	B	369	PHE	2.8
1	A	152	GLY	2.7
1	A	155	ALA	2.7
1	A	209	ALA	2.5
1	B	354	ASN	2.5
1	A	215	LEU	2.5
1	B	161	GLY	2.4
1	A	182	GLY	2.4
1	B	417	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	175	LEU	2.2
1	A	372	LYS	2.2
1	A	153	ASP	2.1
1	A	158	ASP	2.1
1	B	414	ASN	2.1
1	B	163	GLU	2.1
1	A	162	SER	2.1
1	A	211	LYS	2.1
1	B	147	TYR	2.0
1	A	169	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	B	507	4/4	0.77	0.28	46,49,52,52	0
6	EDO	A	505	4/4	0.81	0.25	35,35,37,37	0
6	EDO	A	506	4/4	0.83	0.35	47,51,51,51	0
6	EDO	B	505	4/4	0.85	0.18	43,45,48,49	0
3	ADE	B	502	10/10	0.93	0.16	21,35,42,44	0
4	NA	B	503	1/1	0.94	0.23	39,39,39,39	0
6	EDO	B	506	4/4	0.95	0.12	17,18,19,21	0
3	ADE	A	502	10/10	0.96	0.12	21,27,33,35	0
5	BR	A	504	1/1	0.97	0.05	49,49,49,49	0
2	NAD	B	501	44/44	0.98	0.07	10,17,23,25	0
2	NAD	A	501	44/44	0.98	0.08	13,21,25,28	0
4	NA	A	503	1/1	0.98	0.07	34,34,34,34	0
5	BR	B	504	1/1	0.98	0.09	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.