



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 2, 2021 – 08:10 PM JST

PDB ID : 6M60
Title : Plumbagin in complex with CRM1#-Ran-RanBP1
Authors : Sun, Q.; Lei, Y.
Deposited on : 2020-03-12
Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.19
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.19


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>16%</div> <div>90%</div> <div>6%</div> <div>.</div> </div>
2	B	140	<div> <div>13%</div> <div>81%</div> <div>5%</div> <div>14%</div> </div>
3	C	1003	<div> <div>7%</div> <div>88%</div> <div>11%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	302	-	-	-	X

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 11383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	0	0	0
			1663	1074	284	299	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	LEU	GLN	engineered mutation	UNP P62826
A	182	ALA	LEU	engineered mutation	UNP P62826

- Molecule 2 is a protein called Ran-specific GTPase-activating protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	121	Total	C	N	O	S	0	0	0
			992	628	174	186	4			

- Molecule 3 is a protein called Exportin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	995	Total	C	N	O	S	0	2	0
			8055	5178	1328	1507	42			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P30822
C	-1	GLY	-	expression tag	UNP P30822
C	0	SER	-	expression tag	UNP P30822
C	27	GLU	SER	engineered mutation	UNP P30822
C	49	GLU	GLN	engineered mutation	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	ARG	deletion	UNP P30822

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	LEU	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	THR	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	MET	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	ILE	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	GLN	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	ILE	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	THR	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	GLY	deletion	UNP P30822
C	?	-	ALA	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	ASN	deletion	UNP P30822
C	?	-	PRO	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	TYR	deletion	UNP P30822
C	?	-	MET	deletion	UNP P30822
C	?	-	LYS	deletion	UNP P30822
C	?	-	ARG	deletion	UNP P30822
C	?	-	PHE	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	LEU	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	ASN	deletion	UNP P30822
C	?	-	ASP	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	GLY	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	ILE	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	ARG	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	PHE	deletion	UNP P30822
C	?	-	VAL	deletion	UNP P30822
C	?	-	LYS	deletion	UNP P30822
C	?	-	GLU	deletion	UNP P30822
C	?	-	SER	deletion	UNP P30822
C	?	-	ASP	deletion	UNP P30822
C	?	-	THR	deletion	UNP P30822
C	522	LYS	ASP	engineered mutation	UNP P30822
C	537	GLY	ASP	engineered mutation	UNP P30822
C	539	CYS	THR	engineered mutation	UNP P30822
C	540	GLU	VAL	engineered mutation	UNP P30822
C	541	GLN	LYS	engineered mutation	UNP P30822
C	553	ARG	SER	engineered mutation	UNP P30822
C	561	GLU	GLN	engineered mutation	UNP P30822
C	741	THR	ALA	engineered mutation	UNP P30822
C	1022	CYS	TYR	engineered mutation	UNP P30822

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is NITRATE ION (three-letter code: NO3) (formula: NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	N	O	0	0
			4	1	3		
6	A	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 7 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	3	Total Cl 3 3	0	0

- Molecule 9 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	C	1	Total C O S 4 2 1 1	0	0
9	C	1	Total C O S 4 2 1 1	0	0
9	C	1	Total C O S 4 2 1 1	0	0

- Molecule 10 is (2 {R})-2-methyl-5-oxidanyl-2,3-dihydronaphthalene-1,4-dione (three-letter code: F2X) (formula: $C_{11}H_{10}O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	C	1	Total	C	O	0	0
			14	11	3		
10	C	1	Total	C	O	0	0
			14	11	3		
10	C	1	Total	C	O	0	0
			14	11	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	104	Total	O	0	0
			104	104		
11	B	25	Total	O	0	0
			25	25		
11	C	400	Total	O	0	0
			400	400		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTP-binding nuclear protein Ran




- Molecule 2: Ran-specific GTPase-activating protein 1



- Molecule 3: Exportin-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.59Å 105.59Å 305.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.72 – 2.17 31.72 – 2.17	Depositor EDS
% Data completeness (in resolution range)	99.9 (31.72-2.17) 100.0 (31.72-2.17)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.181 , 0.213 0.188 , 0.217	Depositor DCC
R_{free} test set	4620 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	54.7	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11383	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, CL, MG, GTP, GOL, F2X, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/1704	0.82	0/2310
2	B	0.64	0/1009	0.81	0/1347
3	C	0.66	0/8211	0.75	0/11125
All	All	0.66	0/10924	0.76	0/14782

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1663	0	1666	8	0
2	B	992	0	992	4	0
3	C	8055	0	8146	57	0
4	A	1	0	0	0	0
5	A	6	0	8	0	0
5	C	12	0	16	1	0
6	A	12	0	0	0	0
6	C	24	0	0	0	0
7	A	32	0	12	0	0
8	C	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	C	12	0	18	0	0
10	C	42	0	0	2	0
11	A	104	0	0	0	0
11	B	25	0	0	0	0
11	C	400	0	0	3	0
All	All	11383	0	10858	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:981:VAL:HG22	3:C:982:PRO:HD2	1.58	0.86
3:C:197[A]:CYS:SG	11:C:1224:HOH:O	2.44	0.74
3:C:674:MET:O	3:C:678:THR:HG23	1.89	0.73
3:C:502:TRP:HE1	3:C:547:ASN:HD22	1.38	0.71
3:C:502:TRP:HE1	3:C:547:ASN:ND2	1.92	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/216 (95%)	202 (98%)	4 (2%)	0	100	100
2	B	119/140 (85%)	115 (97%)	4 (3%)	0	100	100
3	C	995/1003 (99%)	976 (98%)	19 (2%)	0	100	100
All	All	1320/1359 (97%)	1293 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/184 (97%)	178 (99%)	1 (1%)	86	92
2	B	103/121 (85%)	101 (98%)	2 (2%)	57	68
3	C	911/914 (100%)	877 (96%)	34 (4%)	34	40
All	All	1193/1219 (98%)	1156 (97%)	37 (3%)	40	48

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	628	ASP
3	C	983	GLN
3	C	688	THR
3	C	876	GLU
3	C	136	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	988	GLN
3	C	1010	GLN
3	C	56	GLN
3	C	541	GLN
3	C	547	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 4 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	GOL	C	1107	-	5,5,5	0.15	0	5,5,5	0.42	0
9	DMS	C	1114	-	3,3,3	0.29	0	3,3,3	0.10	0
6	NO3	C	1101	-	1,3,3	0.56	0	0,3,3	0.00	-
7	GTP	A	305	4	26,34,34	1.05	1 (3%)	33,54,54	2.14	4 (12%)
6	NO3	C	1106	-	1,3,3	0.11	0	0,3,3	0.00	-
9	DMS	C	1112	-	3,3,3	0.27	0	3,3,3	0.19	0
10	F2X	C	1115	3	14,15,15	1.55	2 (14%)	17,22,22	2.05	6 (35%)
9	DMS	C	1105	-	3,3,3	0.24	0	3,3,3	0.12	0
6	NO3	A	304	-	1,3,3	0.50	0	0,3,3	0.00	-
5	GOL	A	302	-	5,5,5	0.16	0	5,5,5	0.36	0
6	NO3	A	303	-	1,3,3	0.76	0	0,3,3	0.00	-
6	NO3	C	1113	-	1,3,3	0.14	0	0,3,3	0.00	-
6	NO3	C	1110	-	1,3,3	0.27	0	0,3,3	0.00	-
5	GOL	C	1108	-	5,5,5	0.13	0	5,5,5	0.40	0
10	F2X	C	1116	3	14,15,15	1.31	2 (14%)	17,22,22	1.67	5 (29%)
6	NO3	C	1111	-	1,3,3	0.20	0	0,3,3	0.00	-
6	NO3	C	1109	-	1,3,3	0.10	0	0,3,3	0.00	-
10	F2X	C	1117	3	14,15,15	1.13	2 (14%)	17,22,22	1.58	5 (29%)
6	NO3	A	306	-	1,3,3	0.23	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	302	-	-	1/4/4/4	-
5	GOL	C	1107	-	-	2/4/4/4	-
10	F2X	C	1117	3	-	-	0/2/2/2
10	F2X	C	1115	3	-	-	0/2/2/2
5	GOL	C	1108	-	-	0/4/4/4	-
10	F2X	C	1116	3	-	-	0/2/2/2
7	GTP	A	305	4	-	2/18/38/38	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	1115	F2X	C1-C2	-3.94	1.35	1.41
7	A	305	GTP	C6-N1	3.74	1.39	1.33
10	C	1115	F2X	C2-C6	-3.56	1.35	1.41
10	C	1117	F2X	C2-C6	-3.24	1.36	1.41
10	C	1116	F2X	C1-C2	-2.97	1.36	1.41

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	305	GTP	C5-C6-N1	-8.86	111.31	123.43
7	A	305	GTP	C6-N1-C2	6.03	125.51	115.93
10	C	1115	F2X	C1-C2-C6	4.12	122.24	118.41
10	C	1115	F2X	C7-C5-C2	3.60	123.42	117.12
10	C	1115	F2X	O3-C5-C2	-3.56	116.74	122.38

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1107	GOL	O1-C1-C2-C3
5	C	1107	GOL	O1-C1-C2-O2
7	A	305	GTP	O4'-C4'-C5'-O5'
5	A	302	GOL	C1-C2-C3-O3
7	A	305	GTP	PA-O3A-PB-O2B

There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1107	GOL	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	C	1115	F2X	1	0
10	C	1117	F2X	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	208/216 (96%)	0.84	34 (16%)	1 1	39, 56, 122, 151	0
2	B	121/140 (86%)	0.62	18 (14%)	2 2	55, 70, 102, 131	0
3	C	995/1003 (99%)	0.42	72 (7%)	15 16	39, 63, 91, 142	0
All	All	1324/1359 (97%)	0.51	124 (9%)	8 8	39, 64, 96, 151	0

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	81	GLU	7.7
3	C	689	LEU	6.9
3	C	616	ARG	5.6
3	C	684	THR	5.5
3	C	543	ARG	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NO3	C	1110	4/4	0.65	0.27	99,100,105,107	0
5	GOL	A	302	6/6	0.67	0.41	73,89,95,98	0
8	CL	C	1103	1/1	0.76	0.15	91,91,91,91	0
9	DMS	C	1114	4/4	0.77	0.32	129,139,141,142	0
8	CL	C	1102	1/1	0.80	0.17	92,92,92,92	0
6	NO3	C	1106	4/4	0.81	0.21	106,106,111,115	0
6	NO3	C	1109	4/4	0.82	0.51	81,85,89,93	0
5	GOL	C	1108	6/6	0.83	0.25	72,80,84,85	0
6	NO3	C	1101	4/4	0.84	0.48	74,76,81,85	0
5	GOL	C	1107	6/6	0.84	0.35	77,86,87,88	0
6	NO3	C	1113	4/4	0.84	0.28	83,89,92,95	0
6	NO3	C	1111	4/4	0.85	0.17	92,93,97,100	0
10	F2X	C	1116	14/14	0.85	0.20	96,108,110,112	0
6	NO3	A	304	4/4	0.87	0.20	82,86,92,93	0
9	DMS	C	1112	4/4	0.87	0.30	139,139,140,140	0
9	DMS	C	1105	4/4	0.88	0.28	110,111,113,114	0
10	F2X	C	1117	14/14	0.88	0.12	86,99,103,103	0
6	NO3	A	306	4/4	0.91	0.17	82,83,85,95	0
6	NO3	A	303	4/4	0.91	0.19	62,64,66,74	0
8	CL	C	1104	1/1	0.93	0.17	97,97,97,97	0
10	F2X	C	1115	14/14	0.94	0.13	69,71,74,76	0
7	GTP	A	305	32/32	0.98	0.10	40,46,49,52	0
4	MG	A	301	1/1	0.99	0.06	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around F2X C 1116:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F2X C 1117:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around F2X C 1115:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.