



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 04:52 AM BST

PDB ID : 6M77  
Title : GH31 alpha-N-acetylgalactosaminidase from *Enterococcus faecalis* in complex with N-acetylgalactosamine  
Authors : Miyazaki, T.  
Deposited on : 2020-03-17  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.4.dev1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	963	<div> <div>7%</div> <div>84%</div> <div>10%</div> <div>6%</div> </div>

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LPXTG-motif cell wall anchor domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	907	Total	C	N	O	S	0	2	0
			7101	4491	1183	1411	16			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-galactopyranose (three-letter code: NGA) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	309	Total O 309 309	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LPXTG-motif cell wall anchor domain protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.60Å 83.23Å 147.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.09 – 1.90 40.06 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.09-1.90) 99.7 (40.06-1.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.37 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.217 , 0.261 0.224 , 0.267	Depositor DCC
$R_{free}$ test set	4011 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.048 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7446	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NGA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	1/7272 (0.0%)	0.90	0/9868

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	751	GLU	CD-OE2	-5.57	1.19	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7101	0	6797	60	0
2	A	15	0	15	0	0
3	A	20	0	30	1	0
4	A	1	0	0	0	0
5	A	309	0	0	3	0
All	All	7446	0	6842	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:768:ASN:HD22	1:A:882:LYS:HA	1.38	0.85
1:A:421[B]:THR:HG22	1:A:423:SER:H	1.46	0.81
1:A:910:THR:HB	1:A:911:PRO:CD	2.15	0.77
1:A:910:THR:HB	1:A:911:PRO:HD2	1.67	0.74
1:A:421[A]:THR:HG21	1:A:452:LEU:HD11	1.68	0.74
1:A:313:ASP:O	1:A:350:GLU:HG2	1.93	0.69
1:A:506:THR:OG1	1:A:520:HIS:HE1	1.77	0.67
1:A:420:TRP:CH2	1:A:422:GLN:HB3	2.34	0.63
1:A:468:ASN:ND2	5:A:1102:HOH:O	2.35	0.59
1:A:132:THR:HA	1:A:144:GLU:O	2.03	0.58
1:A:247:LYS:O	1:A:265:HIS:HD2	1.88	0.56
1:A:421[A]:THR:CG2	1:A:452:LEU:HD11	2.36	0.55
1:A:972:GLU:O	1:A:974:ILE:HD12	2.06	0.55
1:A:421[A]:THR:HG21	1:A:425:LEU:HD21	1.87	0.54
1:A:516:TYR:O	1:A:520:HIS:HD2	1.89	0.54
1:A:251:TYR:OH	1:A:265:HIS:HE1	1.90	0.54
1:A:421[B]:THR:CG2	1:A:423:SER:O	2.56	0.54
1:A:217:ASN:HD21	1:A:493:ASP:CB	2.22	0.53
1:A:545:GLY:HA3	1:A:573:ASN:HD21	1.75	0.52
1:A:93:TYR:CZ	3:A:1004:EDO:H12	2.46	0.51
1:A:600:PRO:HB2	1:A:710:PRO:HB2	1.93	0.51
1:A:133:ASN:O	1:A:143:LEU:HA	2.12	0.50
1:A:311:ASN:HD21	1:A:392:GLN:HG3	1.75	0.50
1:A:461:TYR:HB2	1:A:465:PHE:CD2	2.47	0.49
1:A:308:ASN:HB3	1:A:310:TYR:CE2	2.47	0.49
1:A:756:ALA:HA	1:A:781:LYS:O	2.13	0.49
1:A:119:LYS:NZ	1:A:289:GLU:OE2	2.44	0.48
1:A:924:ALA:HA	1:A:962:VAL:O	2.14	0.48
1:A:421[B]:THR:HG21	1:A:425:LEU:CD2	2.44	0.47
1:A:768:ASN:ND2	1:A:883:GLY:H	2.12	0.47
1:A:900:ALA:HB3	1:A:919:ASP:HB2	1.97	0.47
1:A:817:LEU:HA	1:A:831:ASN:O	2.14	0.47
1:A:860:VAL:HG11	1:A:874:ILE:HD12	1.97	0.47
1:A:913:SER:OG	1:A:913:SER:O	2.33	0.46
1:A:353:ASN:HD21	1:A:406:PHE:CA	2.29	0.46
1:A:421[B]:THR:HG21	1:A:423:SER:O	2.16	0.46
1:A:467:LEU:HA	1:A:470:VAL:HG22	1.98	0.44
1:A:353:ASN:HD22	1:A:364:ARG:HE	1.64	0.44
1:A:61:ILE:CD1	1:A:152:PHE:HB3	2.48	0.44
1:A:208:HIS:HD2	5:A:1225:HOH:O	2.01	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:MET:HB2	1:A:302:PHE:CE1	2.52	0.44
1:A:353:ASN:HD21	1:A:406:PHE:HB2	1.83	0.44
1:A:421[B]:THR:HG22	1:A:423:SER:O	2.18	0.43
1:A:426:HIS:HD2	5:A:1139:HOH:O	2.00	0.43
1:A:429:ASP:C	1:A:429:ASP:OD1	2.57	0.43
1:A:947:ASP:N	1:A:947:ASP:OD1	2.51	0.43
1:A:112:HIS:HB3	1:A:464:SER:HB2	2.00	0.43
1:A:311:ASN:ND2	1:A:392:GLN:HG3	2.34	0.43
1:A:639:MET:HA	1:A:644:LEU:O	2.19	0.42
1:A:306:HIS:C	1:A:307:LEU:HD12	2.40	0.42
1:A:88:ASP:O	1:A:147:GLY:HA3	2.20	0.42
1:A:376:PRO:HA	1:A:716:ASN:HA	2.02	0.42
1:A:307:LEU:HD12	1:A:307:LEU:N	2.36	0.41
1:A:421[B]:THR:OG1	1:A:425:LEU:HD21	2.20	0.41
1:A:654:GLN:HA	1:A:659:ASN:O	2.20	0.41
1:A:198:GLY:O	1:A:208:HIS:HE1	2.03	0.41
1:A:353:ASN:HD21	1:A:406:PHE:HA	1.86	0.41
1:A:421[A]:THR:HG22	1:A:438:ARG:HH12	1.86	0.41
1:A:838:GLU:HA	1:A:854:ASN:HB3	2.02	0.41
1:A:918:TRP:O	1:A:942:ASN:HB2	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	901/963 (94%)	865 (96%)	34 (4%)	2 (0%)	47 38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	912	SER

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	910	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	759/805 (94%)	749 (99%)	10 (1%)	69	68

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	LYS
1	A	275	PHE
1	A	346	ASN
1	A	379	TRP
1	A	492	LEU
1	A	505	TRP
1	A	519	PHE
1	A	603	TYR
1	A	894	ASP
1	A	941	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	GLN
1	A	208	HIS
1	A	214	GLN
1	A	217	ASN
1	A	250	ASN
1	A	265	HIS
1	A	311	ASN
1	A	332	ASN
1	A	353	ASN
1	A	402	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	411	GLN
1	A	426	HIS
1	A	468	ASN
1	A	520	HIS
1	A	573	ASN
1	A	715	ASN
1	A	739	ASN
1	A	768	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	A	1002	-	3,3,3	0.27	0	2,2,2	0.38	0
3	EDO	A	1006	-	3,3,3	0.21	0	2,2,2	0.15	0
3	EDO	A	1005	-	3,3,3	0.18	0	2,2,2	0.30	0
3	EDO	A	1004	-	3,3,3	0.12	0	2,2,2	0.29	0
2	NGA	A	1001	-	15,15,15	1.19	1 (6%)	21,21,21	2.51	6 (28%)
3	EDO	A	1003	-	3,3,3	0.26	0	2,2,2	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1002	-	-	0/1/1/1	-
3	EDO	A	1006	-	-	0/1/1/1	-
3	EDO	A	1005	-	-	0/1/1/1	-
3	EDO	A	1004	-	-	0/1/1/1	-
2	NGA	A	1001	-	-	1/6/26/26	0/1/1/1
3	EDO	A	1003	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	NGA	O5-C5	-2.74	1.37	1.44

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	NGA	O5-C1-C2	6.79	116.34	109.52
2	A	1001	NGA	C4-C3-C2	-5.46	102.35	110.34
2	A	1001	NGA	O1-C1-O5	-4.46	96.98	110.38
2	A	1001	NGA	C1-C2-N2	3.08	114.30	110.73
2	A	1001	NGA	C3-C4-C5	3.06	115.70	110.24
2	A	1001	NGA	C8-C7-N2	2.01	119.49	116.10

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	NGA	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1004	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	907/963 (94%)	0.52	71 (7%) 13 14	15, 30, 64, 111	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	911	PRO	6.8
1	A	899	PRO	6.2
1	A	941	THR	6.2
1	A	974	ILE	5.3
1	A	900	ALA	5.2
1	A	918	TRP	5.2
1	A	915	THR	5.1
1	A	916	LEU	5.0
1	A	917	GLN	5.0
1	A	914	LEU	4.9
1	A	920	GLN	4.7
1	A	957	PHE	4.7
1	A	894	ASP	4.6
1	A	898	ILE	4.5
1	A	895	GLN	4.4
1	A	897	THR	4.4
1	A	964	LYS	4.3
1	A	955	HIS	4.1
1	A	921	VAL	4.0
1	A	943	THR	3.8
1	A	949	PHE	3.8
1	A	956	THR	3.7
1	A	970	TRP	3.6
1	A	971	SER	3.6
1	A	925	THR	3.6
1	A	927	TYR	3.6
1	A	908	LYS	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	944	ALA	3.6
1	A	945	THR	3.5
1	A	946	PHE	3.5
1	A	874	ILE	3.5
1	A	909	THR	3.3
1	A	963	GLY	3.2
1	A	661	ILE	3.2
1	A	972	GLU	3.2
1	A	924	ALA	3.2
1	A	933	GLY	3.2
1	A	256	HIS	3.1
1	A	940	GLN	3.1
1	A	919	ASP	3.0
1	A	959	VAL	3.0
1	A	165	LYS	2.9
1	A	817	LEU	2.9
1	A	942	ASN	2.8
1	A	975	LYS	2.7
1	A	936	PHE	2.7
1	A	912	SER	2.6
1	A	939	ILE	2.6
1	A	431	LYS	2.6
1	A	958	ARG	2.6
1	A	830	THR	2.5
1	A	910	THR	2.5
1	A	57	SER	2.5
1	A	901	ASN	2.5
1	A	76	ALA	2.5
1	A	166	LYS	2.4
1	A	892	VAL	2.4
1	A	947	ASP	2.4
1	A	812	GLY	2.3
1	A	62	VAL	2.3
1	A	525	ILE	2.3
1	A	824	GLU	2.3
1	A	821	ALA	2.3
1	A	950	SER	2.3
1	A	973	PRO	2.2
1	A	430	PRO	2.2
1	A	559	THR	2.1
1	A	535	VAL	2.1
1	A	813	THR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	888	GLY	2.0
1	A	561	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	1006	4/4	0.89	0.16	38,40,41,44	0
3	EDO	A	1004	4/4	0.91	0.12	40,41,41,42	0
3	EDO	A	1005	4/4	0.92	0.16	31,40,41,44	0
3	EDO	A	1002	4/4	0.93	0.12	32,35,35,36	0
3	EDO	A	1003	4/4	0.95	0.14	20,22,23,24	0
2	NGA	A	1001	15/15	0.96	0.09	13,18,20,20	0
4	CA	A	1007	1/1	0.98	0.04	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around NGA A 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.