



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 19, 2021 – 10:08 AM JST

PDB ID : 6M7A
Title : Structure of REV7-R124A complexed with SHLD3(28-73)
Authors : Ma, Y.Z.; Li, Y.; Wu, B.X.; Huang, H.D.
Deposited on : 2020-03-18
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>8%</div> <div>81% 12% • 5%</div> </div>
1	B	211	<div> <div>8%</div> <div>75% 19% 5%</div> </div>
2	C	46	<div> <div>9%</div> <div>67% 11% 22%</div> </div>
2	D	46	<div> <div>7%</div> <div>74% • 22%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitotic spindle assembly checkpoint protein MAD2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1620	1046	272	292	10			
1	B	200	Total	C	N	O	S	0	0	0
			1616	1044	269	293	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	ALA	ARG	engineered mutation	UNP Q9UI95
A	209	LYS	-	expression tag	UNP Q9UI95
A	210	GLY	-	expression tag	UNP Q9UI95
A	211	SER	-	expression tag	UNP Q9UI95
B	124	ALA	ARG	engineered mutation	UNP Q9UI95
B	209	LYS	-	expression tag	UNP Q9UI95
B	210	GLY	-	expression tag	UNP Q9UI95
B	211	SER	-	expression tag	UNP Q9UI95

- Molecule 2 is a protein called Shieldin complex subunit 3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	36	Total	C	N	O	0	0	0
			296	196	47	53			
2	C	36	Total	C	N	O	0	0	0
			296	196	47	53			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	161	Total	O	0	0
			161	161		
3	D	39	Total	O	0	0
			39	39		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	144	Total 144	O 144	0	0
3	C	37	Total 37	O 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2B




- Molecule 1: Mitotic spindle assembly checkpoint protein MAD2B



- Molecule 2: Shieldin complex subunit 3



- Molecule 2: Shieldin complex subunit 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.52Å 62.12Å 130.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 39.81 – 1.90	Depositor EDS
% Data completeness (in resolution range)	91.7 (40.00-1.90) 91.7 (39.81-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.16-3549	Depositor
R, R_{free}	0.180 , 0.232 0.192 , 0.246	Depositor DCC
R_{free} test set	1768 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4209	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1656	0.61	1/2253 (0.0%)
1	B	0.45	1/1651 (0.1%)	0.61	0/2246
2	C	0.32	0/307	0.47	0/419
2	D	0.39	0/307	0.55	0/419
All	All	0.43	1/3921 (0.0%)	0.60	1/5337 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	93	ARG	NE-CZ	5.42	1.40	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	LEU	CA-CB-CG	-6.48	100.40	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1620	0	1648	34	0
1	B	1616	0	1652	45	0
2	C	296	0	294	7	0
2	D	296	0	294	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	161	0	0	3	0
3	B	144	0	0	8	0
3	C	37	0	0	1	0
3	D	39	0	0	1	0
All	All	4209	0	3888	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:11:PHE:HB2	1:B:107:LEU:HD22	1.45	0.99
1:B:42:PHE:O	3:B:301:HOH:O	1.88	0.91
1:B:43:GLN:HE21	1:B:45:ARG:HE	1.16	0.90
2:D:52:ARG:NH2	3:D:101:HOH:O	2.07	0.86
1:A:116:LEU:HG	1:A:117:SER:H	1.46	0.78
1:B:43:GLN:NE2	1:B:45:ARG:HE	1.83	0.77
1:B:79:ASP:OD1	3:B:302:HOH:O	2.05	0.72
1:A:111:SER:HA	2:C:71:TYR:HE2	1.56	0.71
1:B:56:CYS:SG	3:B:320:HOH:O	2.49	0.69
1:A:116:LEU:HG	1:A:117:SER:N	2.08	0.68
1:B:77:LYS:HA	1:B:77:LYS:HE3	1.76	0.68
1:B:107:LEU:HD12	1:B:107:LEU:O	1.94	0.68
1:A:111:SER:HA	2:C:71:TYR:CE2	2.29	0.66
1:B:11:PHE:O	1:B:15:VAL:HG23	1.97	0.64
1:B:15:VAL:HG22	1:B:196:ILE:HG22	1.78	0.64
1:B:78:ASN:ND2	1:B:104:GLN:HB2	2.14	0.63
2:C:38:PHE:N	3:C:102:HOH:O	2.34	0.61
1:A:45:ARG:NH1	3:A:306:HOH:O	2.33	0.60
1:B:114:SER:HB3	1:B:117:SER:HB3	1.82	0.60
1:A:69:HIS:CD2	1:B:69:HIS:CE1	2.90	0.60
1:A:122:LEU:HD13	1:A:192:MET:HG3	1.83	0.60
1:A:69:HIS:CD2	1:B:69:HIS:HE1	2.22	0.58
1:A:161:GLU:N	3:A:308:HOH:O	2.36	0.58
1:A:110:ILE:HD12	1:A:110:ILE:C	2.24	0.57
1:B:110:ILE:H	1:B:110:ILE:HD12	1.74	0.53
1:A:116:LEU:CG	1:A:117:SER:H	2.20	0.53
1:B:15:VAL:HG22	1:B:196:ILE:CG2	2.39	0.53
1:B:15:VAL:CG2	1:B:196:ILE:HG22	2.38	0.53
1:B:78:ASN:HD22	1:B:104:GLN:HB2	1.72	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ALA:HB3	1:A:173:LEU:HD21	1.90	0.52
1:B:10:ASN:OD1	3:B:303:HOH:O	2.19	0.52
1:B:115:LEU:HD11	1:B:196:ILE:HG13	1.91	0.51
1:B:111:SER:HB3	1:B:113:ASP:OD1	2.11	0.51
1:B:149:LEU:HD13	1:B:179:VAL:HG22	1.93	0.51
1:A:33:VAL:HG23	1:A:34:ARG:HG2	1.94	0.50
1:A:58:PRO:HB3	1:B:13:GLN:HG2	1.92	0.50
1:B:113:ASP:OD1	1:B:113:ASP:N	2.41	0.50
1:A:110:ILE:C	1:A:110:ILE:CD1	2.81	0.48
1:B:190:LYS:HD3	1:B:191:THR:H	1.79	0.48
1:A:58:PRO:CB	1:B:13:GLN:HG2	2.44	0.48
1:B:14:VAL:O	1:B:18:VAL:HG23	2.13	0.48
1:A:11:PHE:O	1:A:15:VAL:HG12	2.14	0.47
1:B:73:PRO:HB2	1:B:162:LYS:HD2	1.95	0.47
1:B:162:LYS:HA	1:B:162:LYS:HE2	1.95	0.47
1:B:162:LYS:O	1:B:162:LYS:HD3	2.14	0.47
1:B:45:ARG:NH1	3:B:320:HOH:O	2.47	0.47
1:A:51:PRO:HG3	3:A:402:HOH:O	2.15	0.47
3:B:301:HOH:O	2:C:69:LYS:NZ	2.48	0.47
1:A:167:LYS:HG2	1:B:165:VAL:O	2.16	0.46
1:A:113:ASP:O	1:A:116:LEU:HD21	2.16	0.45
1:A:77:LYS:CE	1:A:162:LYS:HE2	2.47	0.45
1:A:124:ALA:O	1:A:128:LEU:HG	2.17	0.45
1:A:122:LEU:HB3	1:A:192:MET:SD	2.56	0.45
1:A:91:GLU:HB2	1:A:93:ARG:NE	2.33	0.44
1:B:129:LYS:HG3	1:B:190:LYS:HB2	1.99	0.44
1:B:177:GLN:NE2	1:B:178:ASP:OD1	2.38	0.44
1:B:156:ALA:O	2:C:51:LEU:HD21	2.18	0.43
1:B:126:PHE:CE1	1:B:190:LYS:HG3	2.53	0.43
1:A:32:TYR:CE1	1:A:39:VAL:HG23	2.54	0.43
1:B:19:LEU:HA	1:B:19:LEU:HD23	1.81	0.43
1:A:167:LYS:HG2	1:B:165:VAL:HA	2.01	0.43
1:A:116:LEU:HD13	2:C:71:TYR:O	2.19	0.43
1:B:73:PRO:HG3	1:B:165:VAL:HG21	2.01	0.42
2:C:62:GLU:H	2:C:62:GLU:HG3	1.64	0.42
1:A:167:LYS:HG2	1:B:165:VAL:C	2.40	0.42
1:B:116:LEU:HA	1:B:119:VAL:HG12	2.01	0.42
1:B:190:LYS:NZ	3:B:323:HOH:O	2.52	0.42
1:A:32:TYR:HE1	1:A:39:VAL:HG23	1.84	0.41
1:A:11:PHE:HB3	1:A:107:LEU:HD22	2.03	0.41
1:A:156:ALA:HB1	2:D:51:LEU:HG	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:LYS:NZ	3:B:324:HOH:O	2.53	0.40
1:A:115:LEU:O	1:A:116:LEU:HB3	2.21	0.40
1:B:108:LEU:O	1:B:110:ILE:HD12	2.22	0.40
1:A:77:LYS:HE2	1:A:162:LYS:HE2	2.03	0.40
1:A:69:HIS:HD2	1:B:69:HIS:CE1	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/211 (94%)	191 (96%)	7 (4%)	1 (0%)	29	18
1	B	198/211 (94%)	193 (98%)	4 (2%)	1 (0%)	29	18
2	C	34/46 (74%)	34 (100%)	0	0	100	100
2	D	34/46 (74%)	33 (97%)	1 (3%)	0	100	100
All	All	465/514 (90%)	451 (97%)	12 (3%)	2 (0%)	34	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	LEU
1	B	9	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/198 (94%)	182 (97%)	5 (3%)	44	38
1	B	188/198 (95%)	186 (99%)	2 (1%)	73	73
2	C	33/43 (77%)	33 (100%)	0	100	100
2	D	33/43 (77%)	33 (100%)	0	100	100
All	All	441/482 (92%)	434 (98%)	7 (2%)	62	60

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	HIS
1	A	76	GLU
1	A	77	LYS
1	A	93	ARG
1	A	111	SER
1	B	33	VAL
1	B	177	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	HIS
1	B	43	GLN
1	B	53	GLN
1	B	61	ASN
1	B	62	GLN
1	B	69	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/211 (95%)	0.24	16 (7%) 12 13	3, 19, 61, 95	0
1	B	200/211 (94%)	0.24	17 (8%) 10 12	4, 21, 60, 84	0
2	C	36/46 (78%)	0.79	4 (11%) 5 6	16, 32, 56, 62	0
2	D	36/46 (78%)	0.43	3 (8%) 11 13	7, 26, 52, 53	0
All	All	473/514 (92%)	0.30	40 (8%) 10 12	3, 22, 60, 95	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	SER	8.7
1	A	110	ILE	7.0
1	B	112	SER	5.8
2	C	38	PHE	5.7
1	A	114	SER	5.3
2	C	39	ILE	5.2
1	B	109	SER	5.1
1	A	11	PHE	4.8
1	A	115	LEU	4.6
1	A	113	ASP	4.5
1	A	108	LEU	4.3
1	A	111	SER	4.2
1	A	107	LEU	4.2
1	B	8	ASP	4.1
1	B	156	ALA	4.1
1	B	158	ARG	4.0
1	B	207	ALA	4.0
1	A	9	LEU	3.9
1	A	112	SER	3.7
1	B	106	PRO	3.6
2	D	38	PHE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	71	TYR	3.4
1	B	196	ILE	3.4
1	A	116	LEU	3.3
2	D	39	ILE	3.2
1	B	157	THR	3.1
2	C	71	TYR	3.0
1	B	107	LEU	2.9
1	B	111	SER	2.9
2	C	70	GLN	2.9
1	B	182	HIS	2.8
1	B	108	LEU	2.8
1	A	178	ASP	2.8
1	B	110	ILE	2.6
1	A	106	PRO	2.5
1	B	183	ASP	2.2
1	B	185	ARG	2.2
1	B	114	SER	2.1
1	A	8	ASP	2.1
1	A	208	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.