



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:52 am BST

PDB ID : 3M8B
Title : Crystal structure of spin-labeled BtuB V10R1 in the apo state
Authors : Freed, D.M.; Horanyi, P.S.; Wiener, M.C.; Cafiso, D.S.
Deposited on : 2010-03-17
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C8E	A	607	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4605 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vitamin B12 transporter btuB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	549	Total	C	N	O	S	0	0	0
			4329	2727	744	855	3			

- Molecule 2 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: C₁₆H₃₄O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			21	16	5		
2	A	1	Total	C	O	0	0
			21	16	5		

- Molecule 3 is S-[(1-oxyl-2,2,5,5-tetramethyl-2,5-dihydro-1H-pyrrol-3-yl)methyl] methanesulfonothioate (three-letter code: MTN) (formula: $C_{10}H_{18}NO_3S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	9	1	1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Mg	0	0
			4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	113	Total	O	0	0
			113	113		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vitamin B12 transporter *btuB*



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	81.29 Å 81.29 Å 226.56 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.13 – 2.44 44.13 – 2.44	Depositor EDS
% Data completeness (in resolution range)	97.6 (44.13-2.44) 97.6 (44.13-2.44)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.45 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.219 , 0.248 0.218 , 0.245	Depositor DCC
R_{free} test set	1646 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	52.1	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4605	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MTN, C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	3/4433 (0.1%)	0.92	7/6030 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	464	VAL	CB-CG1	5.26	1.63	1.52
1	A	38	GLN	CD-NE2	-5.20	1.19	1.32
1	A	32	GLN	CG-CD	5.06	1.62	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	487	ARG	NE-CZ-NH2	-6.87	116.86	120.30
1	A	31	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	498	ARG	NE-CZ-NH1	6.33	123.46	120.30
1	A	31	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	535	PRO	N-CA-C	-5.77	97.11	112.10
1	A	487	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	A	306	VAL	CB-CA-C	-5.04	101.82	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	534	TYR	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4329	0	4074	40	0
2	A	147	0	238	10	0
3	A	12	0	15	4	0
4	A	4	0	0	0	0
5	A	113	0	0	3	0
All	All	4605	0	4327	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:608:MTN:C5	3:A:608:MTN:C6	1.85	1.55
1:A:38:GLN:OE1	1:A:562:ARG:HB3	1.69	0.92
1:A:160:LEU:HD13	1:A:166:VAL:HG23	1.49	0.92
1:A:38:GLN:NE2	1:A:564:LYS:HB2	1.87	0.90
1:A:267:THR:HG23	1:A:298:VAL:HG22	1.57	0.86
2:A:603:C8E:H52	2:A:603:C8E:H13	1.58	0.84
1:A:506:GLN:OE1	1:A:519:THR:HB	1.85	0.77
1:A:422:PHE:CG	2:A:602:C8E:H11	2.25	0.71
3:A:608:MTN:N1	3:A:608:MTN:C6	2.54	0.70
1:A:48:ARG:HH12	1:A:519:THR:HG21	1.58	0.69
2:A:603:C8E:H171	2:A:603:C8E:H11	1.73	0.69
1:A:86:ASN:HB2	5:A:733:HOH:O	1.92	0.68
1:A:527:TYR:CE1	1:A:540:LYS:HG3	2.29	0.67
1:A:19:ARG:HE	1:A:26:THR:CG2	2.11	0.63
3:A:608:MTN:C3	3:A:608:MTN:C6	2.76	0.61
1:A:198:LEU:C	1:A:198:LEU:HD23	2.21	0.59
1:A:100:PRO:HG2	1:A:103:LEU:HD12	1.83	0.59
1:A:19:ARG:HE	1:A:26:THR:HG23	1.68	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:PRO:O	1:A:21:THR:HG22	2.04	0.58
1:A:267:THR:CG2	1:A:298:VAL:HG22	2.33	0.57
1:A:551:VAL:HG22	1:A:552:ALA:N	2.22	0.54
2:A:603:C8E:C5	2:A:603:C8E:H13	2.35	0.53
2:A:603:C8E:H11	2:A:603:C8E:C17	2.39	0.53
1:A:160:LEU:HD13	1:A:166:VAL:CG2	2.31	0.52
1:A:304:VAL:O	1:A:304:VAL:HG23	2.09	0.52
1:A:38:GLN:HE22	1:A:564:LYS:HB2	1.73	0.51
1:A:19:ARG:NH2	1:A:26:THR:O	2.43	0.51
1:A:304:VAL:HG23	1:A:306:VAL:CG2	2.41	0.51
1:A:227:THR:HB	1:A:243:ARG:HB3	1.94	0.49
1:A:519:THR:HG23	1:A:548:ASP:HB2	1.93	0.49
2:A:605:C8E:H141	5:A:747:HOH:O	2.13	0.49
2:A:603:C8E:C1	2:A:603:C8E:H52	2.36	0.48
1:A:496:LEU:O	1:A:497:ARG:HB2	2.14	0.48
1:A:460:ARG:NH1	5:A:763:HOH:O	2.46	0.47
1:A:160:LEU:CD1	1:A:166:VAL:HG23	2.35	0.46
1:A:321:THR:HA	1:A:333:TYR:O	2.17	0.44
1:A:19:ARG:NE	1:A:26:THR:HG23	2.33	0.43
1:A:422:PHE:CB	2:A:602:C8E:H11	2.47	0.43
1:A:90:VAL:HG13	1:A:326:THR:HG22	2.00	0.43
1:A:323:THR:HG22	1:A:324:PRO:O	2.19	0.42
1:A:53:ASP:HB3	1:A:67:PHE:HB2	2.01	0.42
1:A:213:ALA:HB1	1:A:256:TYR:CE1	2.54	0.42
1:A:587:THR:HG22	1:A:588:LEU:N	2.33	0.42
1:A:422:PHE:CD2	2:A:602:C8E:H21	2.55	0.42
1:A:38:GLN:HE21	1:A:564:LYS:HB2	1.75	0.41
1:A:26:THR:CG2	1:A:26:THR:O	2.68	0.41
3:A:608:MTN:C7	3:A:608:MTN:C6	2.87	0.41
1:A:198:LEU:HD23	1:A:199:SER:N	2.36	0.40
1:A:433:ILE:HG23	1:A:433:ILE:O	2.21	0.40
1:A:304:VAL:O	1:A:304:VAL:CG2	2.70	0.40
2:A:603:C8E:C5	2:A:603:C8E:C1	2.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	541/594 (91%)	523 (97%)	16 (3%)	2 (0%)	34	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	456	GLU
1	A	535	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/495 (92%)	443 (98%)	10 (2%)	52	64

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	THR
1	A	26	THR
1	A	91	SER
1	A	158	GLN
1	A	206	LEU
1	A	242	THR
1	A	267	THR
1	A	270	SER
1	A	519	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	540	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	C8E	A	603	-	20,20,20	0.48	0	19,19,19	0.60	0
2	C8E	A	607	-	20,20,20	0.49	0	19,19,19	0.42	0
3	MTN	A	608	1	9,12,16	5.03	5 (55%)	9,20,27	6.79	4 (44%)
2	C8E	A	604	-	20,20,20	0.45	0	19,19,19	0.48	0
2	C8E	A	605	-	20,20,20	0.43	0	19,19,19	0.37	0
2	C8E	A	602	-	20,20,20	0.42	0	19,19,19	0.73	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	C8E	A	606	-	20,20,20	0.59	0	19,19,19	0.45	0
2	C8E	A	601	-	20,20,20	0.38	0	19,19,19	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C8E	A	603	-	-	7/18/18/18	-
2	C8E	A	607	-	-	9/18/18/18	-
3	MTN	A	608	1	-	0/0/25/29	0/1/1/1
2	C8E	A	604	-	-	13/18/18/18	-
2	C8E	A	605	-	-	11/18/18/18	-
2	C8E	A	602	-	-	12/18/18/18	-
2	C8E	A	606	-	-	10/18/18/18	-
2	C8E	A	601	-	-	12/18/18/18	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	608	MTN	C6-C5	13.46	1.85	1.53
3	A	608	MTN	C9-C1	3.75	1.60	1.53
3	A	608	MTN	C5-C3	-3.23	1.46	1.51
3	A	608	MTN	C1-C2	2.67	1.54	1.50
3	A	608	MTN	C2-C3	2.41	1.35	1.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	608	MTN	C1-C2-C3	-13.30	103.23	113.64
3	A	608	MTN	C8-C1-N1	-9.89	97.36	110.00
3	A	608	MTN	C9-C1-N1	8.93	121.41	110.00
3	A	608	MTN	C2-C1-N1	7.14	104.87	99.37

There are no chirality outliers.

All (74) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	605	C8E	C11-C10-O9-C8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	605	C8E	O15-C16-C17-O18
2	A	604	C8E	O9-C10-C11-O12
2	A	602	C8E	O15-C16-C17-O18
2	A	606	C8E	O15-C16-C17-O18
2	A	602	C8E	O9-C10-C11-O12
2	A	607	C8E	O15-C16-C17-O18
2	A	601	C8E	O15-C16-C17-O18
2	A	602	C8E	C6-C7-C8-O9
2	A	607	C8E	O18-C19-C20-O21
2	A	604	C8E	O18-C19-C20-O21
2	A	605	C8E	O18-C19-C20-O21
2	A	602	C8E	O18-C19-C20-O21
2	A	606	C8E	O18-C19-C20-O21
2	A	601	C8E	C3-C4-C5-C6
2	A	601	C8E	O18-C19-C20-O21
2	A	601	C8E	C2-C3-C4-C5
2	A	602	C8E	C4-C5-C6-C7
2	A	601	C8E	C5-C6-C7-C8
2	A	604	C8E	C4-C5-C6-C7
2	A	607	C8E	C5-C6-C7-C8
2	A	601	C8E	C4-C5-C6-C7
2	A	601	C8E	O9-C10-C11-O12
2	A	605	C8E	O9-C10-C11-O12
2	A	605	C8E	C5-C6-C7-C8
2	A	603	C8E	C2-C3-C4-C5
2	A	607	C8E	C11-C10-O9-C8
2	A	607	C8E	C4-C5-C6-C7
2	A	605	C8E	C1-C2-C3-C4
2	A	605	C8E	C2-C3-C4-C5
2	A	605	C8E	C4-C5-C6-C7
2	A	602	C8E	C11-C10-O9-C8
2	A	606	C8E	O12-C13-C14-O15
2	A	605	C8E	C17-C16-O15-C14
2	A	601	C8E	C6-C7-C8-O9
2	A	607	C8E	C20-C19-O18-C17
2	A	604	C8E	C16-C17-O18-C19
2	A	604	C8E	C7-C8-O9-C10
2	A	606	C8E	C17-C16-O15-C14
2	A	602	C8E	C14-C13-O12-C11
2	A	601	C8E	C14-C13-O12-C11
2	A	604	C8E	C11-C10-O9-C8
2	A	606	C8E	C20-C19-O18-C17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	C8E	C17-C16-O15-C14
2	A	605	C8E	O12-C13-C14-O15
2	A	603	C8E	C7-C8-O9-C10
2	A	607	C8E	O12-C13-C14-O15
2	A	606	C8E	C6-C7-C8-O9
2	A	604	C8E	O15-C16-C17-O18
2	A	605	C8E	C6-C7-C8-O9
2	A	603	C8E	C20-C19-O18-C17
2	A	604	C8E	O12-C13-C14-O15
2	A	604	C8E	C14-C13-O12-C11
2	A	604	C8E	C3-C4-C5-C6
2	A	604	C8E	C5-C6-C7-C8
2	A	602	C8E	C13-C14-O15-C16
2	A	604	C8E	C17-C16-O15-C14
2	A	602	C8E	C1-C2-C3-C4
2	A	601	C8E	C13-C14-O15-C16
2	A	606	C8E	C5-C6-C7-C8
2	A	607	C8E	C2-C3-C4-C5
2	A	606	C8E	C14-C13-O12-C11
2	A	603	C8E	C10-C11-O12-C13
2	A	606	C8E	C10-C11-O12-C13
2	A	601	C8E	C1-C2-C3-C4
2	A	607	C8E	O9-C10-C11-O12
2	A	603	C8E	C6-C7-C8-O9
2	A	603	C8E	O12-C13-C14-O15
2	A	602	C8E	C20-C19-O18-C17
2	A	602	C8E	C10-C11-O12-C13
2	A	606	C8E	O9-C10-C11-O12
2	A	603	C8E	O9-C10-C11-O12
2	A	604	C8E	C13-C14-O15-C16
2	A	602	C8E	O12-C13-C14-O15


There are no ring outliers.





4 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	C8E	6	0
3	A	608	MTN	4	0
2	A	605	C8E	1	0
2	A	602	C8E	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	549/594 (92%)	0.09	24 (4%) 34 32	14, 27, 46, 68	1 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	511	LEU	4.2
1	A	328	TYR	4.2
1	A	325	GLY	4.1
1	A	160	LEU	3.9
1	A	144	TRP	3.9
1	A	512	TYR	3.4
1	A	594	PHE	3.2
1	A	277	TYR	3.1
1	A	324	PRO	3.0
1	A	404	PHE	3.0
1	A	329	VAL	3.0
1	A	162	ASP	2.9
1	A	330	GLU	2.9
1	A	212	ASP	2.9
1	A	288	ALA	2.6
1	A	333	TYR	2.6
1	A	449	HIS	2.6
1	A	331	ASP	2.6
1	A	177	GLY	2.5
1	A	6	ASP	2.4
1	A	161	GLY	2.4
1	A	275	TYR	2.3
1	A	158	GLN	2.1
1	A	289	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	C8E	A	607	21/21	0.48	0.43	54,83,119,119	0
3	MTN	A	608	12/16	0.70	0.40	40,51,53,54	12
2	C8E	A	604	21/21	0.73	0.35	64,76,89,90	0
2	C8E	A	602	21/21	0.73	0.27	69,78,87,88	0
2	C8E	A	606	21/21	0.74	0.30	72,88,93,94	0
4	MG	A	611	1/1	0.86	0.30	65,65,65,65	0
2	C8E	A	605	21/21	0.87	0.16	44,50,59,64	0
4	MG	A	612	1/1	0.91	0.06	76,76,76,76	0
4	MG	A	610	1/1	0.91	0.21	66,66,66,66	0
2	C8E	A	601	21/21	0.91	0.19	40,45,57,61	0
4	MG	A	609	1/1	0.92	0.09	65,65,65,65	0
2	C8E	A	603	21/21	0.93	0.25	36,46,65,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C8E A 607:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MTN A 608:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C8E A 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C8E A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C8E A 606:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C8E A 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C8E A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C8E A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.