



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 03:15 am BST

PDB ID : 4MA9
Title : Wild type Salmonella Alkyl Hydroperoxide Reductase C in its substrate-ready conformation
Authors : Perkins, A.; Karplus, P.A.
Deposited on : 2013-08-15
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	<div> <div>2%</div> <div>85%</div> <div>12%</div> </div>
1	B	186	<div> <div>13%</div> <div>97%</div> </div>
1	C	186	<div> <div>16%</div> <div>98%</div> </div>
1	D	186	<div> <div>12%</div> <div>99%</div> </div>
1	E	186	<div> <div>19%</div> <div>100%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14783 atoms, of which 7029 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkyl hydroperoxide reductase subunit C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	163	Total	C	H	N	O	S	0	4	0
			2550	826	1248	216	257	3			
1	B	186	Total	C	H	N	O	S	0	12	0
			3026	980	1490	254	297	5			
1	C	186	Total	C	H	N	O	S	0	2	0
			2889	933	1424	242	286	4			
1	D	186	Total	C	H	N	O	S	0	2	0
			2886	932	1422	242	286	4			
1	E	186	Total	C	H	N	O	S	0	0	0
			2865	926	1413	241	281	4			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	E	1	Total	Cl	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	K	0	0
			1	1		
3	A	1	Total	K	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total K 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 14 3 8 3	0	0
4	B	1	Total C H O 14 3 8 3	0	0
4	C	1	Total C H O 14 3 8 3	0	0
4	D	1	Total C H O 14 3 8 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	127	Total O 127 127	0	4
5	B	133	Total O 133 133	0	2
5	C	99	Total O 99 99	0	0
5	D	102	Total O 102 102	0	2

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	42	Total	O	0	0
			42	42		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alkyl hydroperoxide reductase subunit C




- Molecule 1: Alkyl hydroperoxide reductase subunit C




- Molecule 1: Alkyl hydroperoxide reductase subunit C



- Molecule 1: Alkyl hydroperoxide reductase subunit C



- Molecule 1: Alkyl hydroperoxide reductase subunit C



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	126.84Å 171.15Å 135.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.76 – 1.82 36.76 – 1.82	Depositor EDS
% Data completeness (in resolution range)	96.0 (36.76-1.82) 96.0 (36.76-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.82Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.204 , 0.240 0.214 , 0.238	Depositor DCC
R_{free} test set	6384 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	29.9	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 67.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14783	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1348	0.69	0/1827
1	B	0.55	0/1586	0.67	0/2151
1	C	0.53	0/1506	0.66	0/2041
1	D	0.50	0/1506	0.65	0/2041
1	E	0.43	0/1486	0.59	0/2014
All	All	0.51	0/7432	0.65	0/10074

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1302	1248	1230	3	0
1	B	1536	1490	1477	4	0
1	C	1465	1424	1415	2	0
1	D	1464	1422	1416	1	0
1	E	1452	1413	1414	0	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	A	6	8	8	0	0
4	B	6	8	8	0	0
4	C	6	8	8	0	0
4	D	6	8	8	0	0
5	A	127	0	0	0	0
5	B	133	0	0	0	0
5	C	99	0	0	0	0
5	D	102	0	0	0	0
5	E	42	0	0	0	0
All	All	7754	7029	6984	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:VAL:HG11	1:B:139:GLY:HA3	1.88	0.55
1:B:31:ARG:HD3	1:B:66:ASP:OD2	2.08	0.53
1:D:88:ILE:HG23	1:D:91:ILE:HD12	2.02	0.42
1:B:43[A]:THR:HG22	1:C:76:PHE:CD2	2.55	0.41
1:B:47[A]:PRO:CG	1:B:88:ILE:HD11	2.51	0.41
1:A:39:PRO:HD2	1:A:46:CYS:SG	2.62	0.40
1:A:81:TRP:CD2	1:A:88:ILE:HG13	2.57	0.40
1:C:65:VAL:HG21	1:C:151:ILE:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/186 (89%)	161 (98%)	4 (2%)	0	100	100
1	B	196/186 (105%)	192 (98%)	4 (2%)	0	100	100
1	C	186/186 (100%)	181 (97%)	5 (3%)	0	100	100
1	D	186/186 (100%)	182 (98%)	4 (2%)	0	100	100
1	E	184/186 (99%)	178 (97%)	6 (3%)	0	100	100
All	All	917/930 (99%)	894 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/154 (91%)	140 (100%)	0	100	100
1	B	164/154 (106%)	164 (100%)	0	100	100
1	C	156/154 (101%)	156 (100%)	0	100	100
1	D	156/154 (101%)	156 (100%)	0	100	100
1	E	153/154 (99%)	153 (100%)	0	100	100
All	All	769/770 (100%)	769 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	D	203	-	5,5,5	0.21	0	5,5,5	0.66	0
4	GOL	A	203	-	5,5,5	0.16	0	5,5,5	0.16	0
4	GOL	B	203	-	5,5,5	0.22	0	5,5,5	0.22	0
4	GOL	C	202	-	5,5,5	0.13	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	203	-	-	4/4/4/4	-
4	GOL	A	203	-	-	0/4/4/4	-
4	GOL	B	203	-	-	2/4/4/4	-
4	GOL	C	202	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	203	GOL	O1-C1-C2-C3
4	D	203	GOL	C1-C2-C3-O3
4	B	203	GOL	O1-C1-C2-C3
4	D	203	GOL	O2-C2-C3-O3
4	D	203	GOL	O1-C1-C2-O2
4	B	203	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	163/186 (87%)	0.05	4 (2%) 57 52	24, 38, 61, 71	0
1	B	186/186 (100%)	0.49	24 (12%) 3 2	24, 39, 81, 101	0
1	C	186/186 (100%)	0.71	30 (16%) 1 1	28, 45, 101, 123	0
1	D	186/186 (100%)	0.59	22 (11%) 4 3	33, 46, 80, 103	0
1	E	186/186 (100%)	1.06	35 (18%) 1 0	42, 67, 103, 128	0
All	All	907/930 (97%)	0.59	115 (12%) 3 2	24, 46, 93, 128	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	172	GLY	9.9
1	E	174	ALA	9.2
1	C	176	LEU	8.7
1	B	182	LEU	8.4
1	B	174	ALA	8.3
1	C	172	GLY	7.7
1	E	182	LEU	7.4
1	D	176	LEU	6.7
1	D	181	ASP	6.6
1	B	180	LEU	6.4
1	B	172	GLY	6.3
1	B	181	ASP	6.2
1	E	176	LEU	6.1
1	C	170	LYS	6.1
1	C	185	LYS	6.1
1	C	171	GLU	6.0
1	D	171	GLU	5.8
1	D	172	GLY	5.8
1	E	159	ALA	5.7
1	E	171	GLU	5.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	174	ALA	5.4
1	E	173	GLU	5.4
1	B	177	ALA	5.4
1	B	179	SER	5.3
1	E	175	THR	5.3
1	B	185	LYS	5.2
1	C	174	ALA	5.2
1	D	179	SER	5.1
1	B	170	LYS	4.9
1	E	185	LYS	4.8
1	C	169	TRP	4.7
1	D	180	LEU	4.7
1	D	182	LEU	4.7
1	E	181	ASP	4.7
1	D	175	THR	4.6
1	B	46[A]	CYS	4.5
1	E	179	SER	4.4
1	B	175	THR	4.3
1	E	169	TRP	4.1
1	C	182	LEU	4.1
1	D	178	PRO	4.1
1	E	135	VAL	3.9
1	B	44[A]	PHE	3.9
1	D	185	LYS	3.8
1	D	177	ALA	3.8
1	E	148	LEU	3.8
1	E	177	ALA	3.7
1	E	180	LEU	3.7
1	E	183	VAL	3.7
1	B	176	LEU	3.7
1	E	161	PRO	3.7
1	E	62	LYS	3.7
1	C	164	VAL	3.6
1	B	169	TRP	3.5
1	E	170	LYS	3.5
1	C	184	GLY	3.4
1	D	133	ILE	3.4
1	E	133	ILE	3.4
1	E	156	TYR	3.4
1	B	171	GLU	3.4
1	E	178	PRO	3.4
1	C	161	PRO	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	180	LEU	3.3
1	B	173	GLU	3.3
1	E	186	ILE	3.2
1	C	62	LYS	3.2
1	D	161	PRO	3.2
1	E	184	GLY	3.2
1	B	164	VAL	3.2
1	C	135	VAL	3.2
1	B	186	ILE	3.1
1	B	162	GLY	3.1
1	C	37	PHE	3.0
1	C	186	ILE	3.0
1	C	168	LYS	3.0
1	B	178	PRO	2.9
1	C	166	PRO	2.9
1	C	120	ALA	2.9
1	C	181	ASP	2.8
1	E	134	GLU	2.8
1	D	169	TRP	2.8
1	C	38	TYR	2.8
1	D	121	THR	2.7
1	B	161	PRO	2.7
1	C	63	LEU	2.7
1	E	58	GLU	2.7
1	C	165	CYS	2.7
1	E	121	THR	2.7
1	B	183	VAL	2.6
1	B	45[A]	VAL	2.6
1	C	121	THR	2.6
1	D	183	VAL	2.5
1	C	183	VAL	2.5
1	C	173	GLU	2.4
1	A	135	VAL	2.4
1	C	36	PHE	2.4
1	D	186	ILE	2.3
1	E	136	THR	2.3
1	E	26	LYS	2.3
1	D	162	GLY	2.3
1	E	164	VAL	2.2
1	A	120	ALA	2.2
1	C	134	GLU	2.2
1	D	135	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	162	GLY	2.1
1	E	92	LYS	2.1
1	D	132	ALA	2.1
1	E	120	ALA	2.1
1	E	44	PHE	2.1
1	C	159	ALA	2.1
1	C	136	THR	2.0
1	A	18	GLY	2.0
1	B	163	GLU	2.0
1	D	36	PHE	2.0
1	A	146[A]	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	203	6/6	0.69	0.22	69,76,76,77	0
4	GOL	C	202	6/6	0.74	0.23	69,77,82,82	0
4	GOL	D	203	6/6	0.88	0.14	60,66,71,72	0
4	GOL	B	203	6/6	0.91	0.29	63,73,80,81	0
3	K	D	202	1/1	0.93	0.09	71,71,71,71	0
3	K	B	202	1/1	0.94	0.14	57,57,57,57	0
3	K	A	202	1/1	0.94	0.22	59,59,59,59	1
2	CL	D	201	1/1	0.99	0.04	46,46,46,46	0
2	CL	E	201	1/1	0.99	0.07	60,60,60,60	0
2	CL	C	201	1/1	0.99	0.04	43,43,43,43	0
2	CL	A	201	1/1	0.99	0.05	37,37,37,37	0
2	CL	B	201	1/1	1.00	0.05	41,41,41,41	0

6.5 Other polymers

There are no such residues in this entry.