



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:25 pm BST

PDB ID : 3MC9
Title : POTRA1-2 of the periplasmic domain of Omp85 from Anabaena
Authors : Koenig, P.; Schleiff, E.; Sinning, I.; Tews, I.
Deposited on : 2010-03-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div></div> <div>42% 8% • 49%</div> </div>
1	B	316	<div> <div>%</div> <div>42% 8% • 48%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alr2269 protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	0	0	0
			1241	781	215	245			
1	B	165	Total	C	N	O	0	0	0
			1270	797	221	252			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	MET	-	EXPRESSION TAG	UNP Q8YUR6
A	468	LEU	-	EXPRESSION TAG	UNP Q8YUR6
A	469	GLU	-	EXPRESSION TAG	UNP Q8YUR6
A	470	HIS	-	EXPRESSION TAG	UNP Q8YUR6
A	471	HIS	-	EXPRESSION TAG	UNP Q8YUR6
A	472	HIS	-	EXPRESSION TAG	UNP Q8YUR6
A	473	HIS	-	EXPRESSION TAG	UNP Q8YUR6
A	474	HIS	-	EXPRESSION TAG	UNP Q8YUR6
A	475	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	160	MET	-	EXPRESSION TAG	UNP Q8YUR6
B	468	LEU	-	EXPRESSION TAG	UNP Q8YUR6
B	469	GLU	-	EXPRESSION TAG	UNP Q8YUR6
B	470	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	471	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	472	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	473	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	474	HIS	-	EXPRESSION TAG	UNP Q8YUR6
B	475	HIS	-	EXPRESSION TAG	UNP Q8YUR6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	85	Total	O	0	0
			85	85		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	68	Total	O	0	0
			68	68		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	108.37Å 108.37Å 60.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.20 46.92 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.20) 100.0 (46.92-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.195 , 0.245 0.201 , 0.258	Depositor DCC
R_{free} test set	1056 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.067 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2664	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/1260	0.67	0/1718
1	B	0.53	0/1290	0.76	3/1762 (0.2%)
All	All	0.55	0/2550	0.72	3/3480 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	248	ARG	NE-CZ-NH2	-7.09	116.75	120.30
1	B	227	LEU	CA-CB-CG	6.19	129.53	115.30
1	B	248	ARG	NE-CZ-NH1	5.78	123.19	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1241	0	1250	28	0
1	B	1270	0	1279	24	0
2	A	85	0	0	12	0
2	B	68	0	0	0	0
All	All	2664	0	2529	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:343:LYS:HE3	1:B:343:LYS:HA	1.47	0.95
1:B:322:ASP:O	1:B:326:ARG:HB2	1.77	0.83
1:B:217:THR:HG23	1:B:218:GLU:H	1.44	0.82
1:B:322:ASP:HB3	1:B:326:ARG:NH1	2.03	0.73
1:A:336:ARG:HG2	2:A:145:HOH:O	1.87	0.73
1:A:316:LEU:HB3	1:A:320:THR:HG21	1.71	0.72
1:A:348:ARG:HH12	1:A:352:GLN:HE21	1.35	0.72
1:A:322:ASP:O	1:A:326:ARG:HB2	1.93	0.69
1:A:236:THR:HG23	1:A:239:LEU:HB2	1.76	0.68
1:B:217:THR:HG23	1:B:218:GLU:N	2.08	0.67
1:B:312:VAL:HG11	1:B:379:VAL:HG22	1.77	0.65
1:B:315:VAL:HG21	1:B:354:TYR:CZ	2.32	0.65
1:A:305:GLN:OE1	1:A:374:GLN:NE2	2.24	0.64
1:A:236:THR:HG21	2:A:78:HOH:O	1.98	0.64
1:A:319:ALA:O	1:A:323:GLU:HG3	1.98	0.63
1:A:236:THR:HG22	2:A:53:HOH:O	1.98	0.63
1:A:368:ASN:HB2	2:A:137:HOH:O	1.99	0.62
1:A:317:PRO:O	1:A:320:THR:CG2	2.48	0.62
1:B:217:THR:CG2	1:B:218:GLU:H	2.12	0.61
2:A:106:HOH:O	1:B:257:ARG:HD2	2.00	0.61
1:B:315:VAL:HG21	1:B:354:TYR:OH	2.01	0.60
1:B:343:LYS:CE	1:B:343:LYS:HA	2.24	0.59
1:A:236:THR:CG2	1:A:239:LEU:HD12	2.34	0.56
1:B:364:GLN:HA	1:B:364:GLN:HE21	1.71	0.56
1:B:322:ASP:HB3	1:B:326:ARG:CZ	2.36	0.55
1:A:317:PRO:O	1:A:320:THR:HG22	2.06	0.55
1:A:308:PRO:HD2	2:A:138:HOH:O	2.09	0.53
1:B:217:THR:CG2	1:B:218:GLU:N	2.74	0.50
1:A:317:PRO:O	1:A:320:THR:HG23	2.12	0.49
1:A:231:GLN:HB3	1:A:294:GLN:HE21	1.77	0.49
1:A:348:ARG:HH12	1:A:352:GLN:NE2	2.04	0.49
1:A:366:SER:CB	2:A:137:HOH:O	2.61	0.49
1:B:364:GLN:CA	1:B:364:GLN:HE21	2.25	0.48
1:A:218:GLU:OE2	1:A:258:SER:OG	2.28	0.48
1:B:311:ASN:N	1:B:311:ASN:OD1	2.47	0.47
1:A:366:SER:HB2	2:A:137:HOH:O	2.13	0.47
1:A:307:ASN:CB	2:A:82:HOH:O	2.63	0.46
1:A:368:ASN:CB	2:A:137:HOH:O	2.61	0.46
1:A:236:THR:HG23	1:A:239:LEU:CB	2.45	0.46
1:A:348:ARG:NH1	1:A:352:GLN:HE21	2.12	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:345:LEU:HD22	1:B:349:TYR:CE2	2.53	0.43
1:B:364:GLN:NE2	1:B:364:GLN:HA	2.32	0.43
1:B:222:LEU:HG	1:B:223:VAL:N	2.34	0.42
1:B:316:LEU:HA	1:B:316:LEU:HD23	1.87	0.42
1:B:301:LYS:HA	1:B:329:TYR:CE1	2.55	0.42
1:A:296:ASN:HB3	1:A:333:LEU:HB3	2.02	0.41
1:A:368:ASN:C	2:A:137:HOH:O	2.59	0.41
1:A:368:ASN:OD1	1:A:368:ASN:N	2.53	0.41
1:B:234:GLN:HE21	1:B:234:GLN:N	2.20	0.40
1:B:231:GLN:HB3	1:B:294:GLN:NE2	2.36	0.40
1:B:312:VAL:HG11	1:B:379:VAL:CG2	2.47	0.40
1:A:308:PRO:C	2:A:148:HOH:O	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/316 (50%)	156 (99%)	1 (1%)	0	100	100
1	B	163/316 (52%)	158 (97%)	4 (2%)	1 (1%)	25	26
All	All	320/632 (51%)	314 (98%)	5 (2%)	1 (0%)	41	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	326	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/277 (50%)	132 (94%)	8 (6%)	20	24
1	B	144/277 (52%)	130 (90%)	14 (10%)	8	7
All	All	284/554 (51%)	262 (92%)	22 (8%)	13	13

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	221	VAL
1	A	234	GLN
1	A	236	THR
1	A	301	LYS
1	A	320	THR
1	A	345	LEU
1	A	364	GLN
1	A	368	ASN
1	B	227	LEU
1	B	234	GLN
1	B	285	LEU
1	B	315	VAL
1	B	336	ARG
1	B	338	LEU
1	B	339	GLN
1	B	345	LEU
1	B	348	ARG
1	B	352	GLN
1	B	358	ASN
1	B	364	GLN
1	B	368	ASN
1	B	379	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	GLN
1	A	275	ASN
1	A	294	GLN
1	A	352	GLN
1	B	231	GLN
1	B	234	GLN
1	B	250	GLN
1	B	275	ASN
1	B	294	GLN
1	B	364	GLN
1	B	374	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/316 (50%)	-0.28	1 (0%) 89 88	27, 41, 57, 65	0
1	B	165/316 (52%)	-0.10	2 (1%) 79 77	25, 43, 61, 74	0
All	All	326/632 (51%)	-0.19	3 (0%) 84 83	25, 42, 59, 74	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	311	ASN	4.0
1	A	308	PRO	3.1
1	B	312	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.