



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 08:02 am BST

PDB ID : 6MDB  
Title : Non-receptor Protein Tyrosine Phosphatase SHP2 in Complex with Allosteric Inhibitor Pyrazolo-pyrimidinone 5  
Authors : Fodor, M.; Stams, T.  
Deposited on : 2018-09-04  
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	<div> <div>8%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>7%</div> </div> </div>
1	B	526	<div> <div>8%</div> <div> <div></div> <div>80%</div> <div>10%</div> <div>9%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	0	2	0
			3943	2486	699	739	19			
1	B	481	Total	C	N	O	S	0	3	0
			3912	2469	695	728	20			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 6-(4-amino-4-methylpiperidin-1-yl)-3-(2,3-dichlorophenyl)-5-methyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one (three-letter code: JE4) (formula: C<sub>18</sub>H<sub>20</sub>Cl<sub>2</sub>N<sub>6</sub>O) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	
			27	18	2	6	1	
2	B	1	Total	C	Cl	N	O	
			27	18	2	6	1	

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O P	0	0
			5	4 1		
3	A	1	Total	O P	0	0
			5	4 1		
3	B	1	Total	O P	0	0
			5	4 1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	267	Total	O	0	0
			267	267		
4	B	257	Total	O	0	0
			257	257		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.27Å 213.52Å 55.76Å 90.00° 96.43° 90.00°	Depositor
Resolution (Å)	23.59 – 2.34 23.59 – 2.34	Depositor EDS
% Data completeness (in resolution range)	98.7 (23.59-2.34) 98.7 (23.59-2.34)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.60 (at 2.33Å)	Xtriage
Refinement program	BUSTER	Depositor
R, $R_{free}$	0.175 , 0.236 0.181 , 0.245	Depositor DCC
$R_{free}$ test set	2160 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 59.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8448	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, JE4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/4029	0.68	1/5430 (0.0%)
1	B	0.51	0/4000	0.68	0/5389
All	All	0.51	0/8029	0.68	1/10819 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	SER	C-N-CA	5.42	135.24	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3943	0	3888	27	0
1	B	3912	0	3870	25	0
2	A	27	0	0	3	0
2	B	27	0	0	3	0
3	A	10	0	0	0	0
3	B	5	0	0	0	0
4	A	267	0	0	0	0
4	B	257	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8448	0	7758	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:GLU:HB3	1:A:301:VAL:HG11	1.47	0.96
1:B:330:THR:HG23	1:B:458:HIS:HB3	1.60	0.82
1:A:294:ASP:HB2	1:A:343:ARG:HE	1.54	0.73
1:B:47:ARG:HG3	1:B:96:ILE:HD11	1.73	0.70
1:A:272:GLU:CB	1:A:301:VAL:HG11	2.25	0.64
1:A:390:GLU:HG2	1:A:399:ARG:HG2	1.81	0.62
1:A:125:LEU:HB3	1:A:216:LEU:HD21	1.83	0.60
1:A:140:SER:HA	1:A:141:GLN:HB3	1.83	0.60
1:A:65:LEU:HD23	1:A:68:GLY:HA3	1.82	0.59
2:A:601:JE4:C5	2:A:601:JE4:C10	2.81	0.58
1:B:125:LEU:HB3	1:B:216:LEU:HD21	1.86	0.58
1:B:114:HIS:HE1	1:B:218:THR:HG21	1.69	0.58
1:A:294:ASP:HB2	1:A:343:ARG:NE	2.19	0.58
1:A:309:ILE:HD13	1:A:328:ILE:HG12	1.89	0.55
1:A:495:GLN:HG2	2:A:601:JE4:CL1	2.46	0.53
1:A:330:THR:HG23	1:A:458:HIS:HB3	1.91	0.52
1:B:133:GLY:HA3	1:B:213:LYS:HB2	1.93	0.51
1:B:117:LEU:O	1:B:138:ARG:HD2	2.12	0.50
2:B:601:JE4:C5	2:B:601:JE4:C10	2.91	0.49
1:B:268:GLY:HA2	1:B:304:TYR:CE2	2.48	0.49
1:B:222:ASN:O	1:B:225:GLU:HG2	2.13	0.49
1:A:257:GLN:HG3	2:A:601:JE4:CL2	2.50	0.49
1:A:326:SER:HB3	1:A:454:PRO:HB3	1.94	0.48
1:A:41:PHE:HB2	1:A:56:ILE:HB	1.94	0.48
1:A:360:VAL:HG21	1:B:394:HIS:CD2	2.49	0.47
1:A:491:PRO:HG3	1:A:511:TYR:OH	2.13	0.47
1:A:399:ARG:HD2	1:A:417:GLN:OE1	2.15	0.47
1:B:134:SER:HA	1:B:214:GLN:O	2.15	0.46
1:A:377:LEU:HD11	1:A:384:ARG:HG2	1.96	0.46
1:A:149:LEU:HB2	1:A:172:ILE:HD11	1.98	0.46
1:A:135:PHE:HB3	1:A:151:VAL:HG13	1.97	0.46
1:B:126:LEU:HD23	1:B:216:LEU:HD13	1.98	0.46
1:A:11:ILE:HD12	1:A:16:ALA:HB2	1.98	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:HIS:CE1	1:B:218:THR:HG21	2.51	0.45
1:B:264:SER:HB2	1:B:266:LYS:HD3	1.98	0.45
1:A:65:LEU:HD21	1:A:88:LEU:HD13	1.98	0.45
1:B:124:LYS:O	1:B:128:GLU:HG2	2.18	0.45
1:B:8:HIS:CD2	1:B:104:CYS:HB2	2.52	0.45
1:B:432:PRO:HG3	1:B:516[A]:MET:HG2	1.99	0.44
1:B:254:LEU:HA	2:B:601:JE4:CL2	2.53	0.44
1:A:56:ILE:HG12	1:A:65:LEU:HD12	1.99	0.44
1:B:495:GLN:HG2	2:B:601:JE4:CL1	2.55	0.43
1:A:285:PHE:H	1:A:308:ASN:HD21	1.66	0.43
1:B:65:LEU:HD21	1:B:88:LEU:HD13	2.02	0.42
1:B:217:ASN:OD1	1:B:220:ARG:HB3	2.19	0.42
1:B:290:VAL:HG11	1:B:344:MET:HG3	2.02	0.42
1:B:273:ASN:HA	1:B:276:LYS:HD2	2.02	0.41
1:A:366:LYS:HD3	1:A:366:LYS:HA	1.86	0.41
1:B:65:LEU:HD23	1:B:68:GLY:HA3	2.02	0.41
1:B:17:GLU:HG3	1:B:51:VAL:HB	2.03	0.41
1:A:9:PRO:HB2	1:A:248:TRP:HH2	1.86	0.40
1:B:4:ARG:HD3	1:B:256:GLN:HA	2.02	0.40
1:A:294:ASP:CB	1:A:343:ARG:HE	2.30	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/526 (90%)	455 (96%)	18 (4%)	2 (0%)	34	38
1	B	470/526 (89%)	455 (97%)	14 (3%)	1 (0%)	47	55
All	All	945/1052 (90%)	910 (96%)	32 (3%)	3 (0%)	41	47

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	295	GLY
1	A	505	VAL
1	B	505	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/468 (93%)	412 (95%)	22 (5%)	24	29
1	B	431/468 (92%)	414 (96%)	17 (4%)	32	41
All	All	865/936 (92%)	826 (96%)	39 (4%)	28	34

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	VAL
1	A	17	GLU
1	A	48	ASN
1	A	55	LYS
1	A	87	GLN
1	A	128	GLU
1	A	140	SER
1	A	152	ARG
1	A	165	SER
1	A	221	ILE
1	A	249	GLU
1	A	256	GLN
1	A	280	LYS
1	A	326	SER
1	A	410	ASN
1	A	413	ARG
1	A	439	LEU
1	A	447	GLU
1	A	477	ASP
1	A	484	VAL
1	A	516[A]	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	516[B]	MET
1	B	47	ARG
1	B	87	GLN
1	B	94	ASP
1	B	96	ILE
1	B	166	LYS
1	B	171	MET
1	B	177	LEU
1	B	199	LYS
1	B	264	SER
1	B	280	LYS
1	B	325	LYS
1	B	413	ARG
1	B	451	ASP
1	B	477	ASP
1	B	481	GLU
1	B	502	SER
1	B	508	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	87	GLN
1	A	211	GLN
1	A	281	ASN
1	A	410	ASN
1	B	87	GLN
1	B	114	HIS
1	B	275	ASN
1	B	394	HIS
1	B	520	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	PO4	A	602	-	4,4,4	2.46	1 (25%)	6,6,6	0.48	0
2	JE4	A	601	-	25,30,30	1.47	2 (8%)	23,46,46	1.64	5 (21%)
3	PO4	A	603	-	4,4,4	2.49	1 (25%)	6,6,6	0.51	0
2	JE4	B	601	-	25,30,30	1.51	3 (12%)	23,46,46	1.85	6 (26%)
3	PO4	B	602	-	4,4,4	2.47	1 (25%)	6,6,6	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JE4	A	601	-	-	2/6/20/20	0/4/4/4
2	JE4	B	601	-	-	3/6/20/20	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	JE4	C4-C1	4.51	1.49	1.41
3	A	603	PO4	P-O1	4.14	1.60	1.50
2	A	601	JE4	C4-C1	4.10	1.49	1.41
3	B	602	PO4	P-O1	4.10	1.60	1.50
3	A	602	PO4	P-O1	4.06	1.60	1.50
2	B	601	JE4	C13-C12	-2.59	1.46	1.49
2	A	601	JE4	C2-N5	-2.53	1.30	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	JE4	C2-N5	-2.11	1.31	1.34

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	JE4	C14-C15-CL1	-4.75	115.89	120.52
2	A	601	JE4	C9-C10-N3	4.47	118.25	110.34
2	B	601	JE4	C9-C10-N3	4.01	117.44	110.34
2	B	601	JE4	C6-N3-C10	3.20	118.57	111.52
2	A	601	JE4	C13-C12-N6	2.77	125.52	120.78
2	A	601	JE4	C10-C9-C8	2.56	117.69	113.22
2	A	601	JE4	C6-N3-C10	2.45	116.92	111.52
2	B	601	JE4	C7-C6-N3	2.34	114.48	110.34
2	B	601	JE4	C13-C12-N6	2.28	124.68	120.78
2	B	601	JE4	C16-C15-CL1	2.21	122.84	118.41
2	A	601	JE4	C14-C15-CL1	-2.19	118.39	120.52

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	JE4	N2-C3-N3-C6
2	B	601	JE4	N1-C3-N3-C10
2	B	601	JE4	N1-C3-N3-C6
2	A	601	JE4	N1-C3-N3-C10
2	A	601	JE4	N1-C3-N3-C6


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	JE4	3	0
2	B	601	JE4	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	487/526 (92%)	0.44	44 (9%) <b>9</b> <b>15</b>	25, 47, 77, 108	0
1	B	481/526 (91%)	0.44	41 (8%) <b>10</b> <b>16</b>	24, 48, 75, 114	0
All	All	968/1052 (92%)	0.44	85 (8%) <b>10</b> <b>15</b>	24, 47, 77, 114	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	THR	6.1
1	A	177	LEU	5.4
1	A	86	GLY	5.0
1	B	86	GLY	4.7
1	B	177	LEU	4.7
1	B	93	GLY	4.6
1	B	426	HIS	4.1
1	B	205	THR	4.1
1	A	95	VAL	4.0
1	A	120	LYS	4.0
1	A	206	LEU	4.0
1	B	12	THR	3.7
1	B	92	ASN	3.7
1	B	207	GLY	3.7
1	A	297	PRO	3.7
1	A	94	ASP	3.4
1	A	128	GLU	3.3
1	B	123	GLU	3.3
1	B	85	HIS	3.2
1	A	199	LYS	3.2
1	A	296	ASP	3.1
1	A	83	GLU	3.1
1	B	362	ARG	3.1
1	A	85	HIS	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	232	GLU	3.0
1	A	116	HIS	3.0
1	B	116	HIS	2.9
1	B	121	GLU	2.9
1	A	174	CYS	2.9
1	A	525	LEU	2.9
1	B	311	MET	2.9
1	A	457	VAL	2.9
1	B	485	ASP	2.8
1	B	206	LEU	2.8
1	B	523	GLU	2.8
1	A	524	THR	2.8
1	B	199	LYS	2.7
1	B	425	ASP	2.7
1	B	260	LYS	2.7
1	B	137	VAL	2.7
1	A	450	MET	2.7
1	A	523	GLU	2.7
1	B	363	GLY	2.7
1	A	375	TYR	2.6
1	B	38	PRO	2.6
1	A	165	SER	2.6
1	A	87	GLN	2.6
1	A	66	TYR	2.6
1	B	323	PRO	2.5
1	A	137	VAL	2.5
1	B	120	LYS	2.5
1	A	38	PRO	2.5
1	A	353	ILE	2.5
1	B	301	VAL	2.5
1	A	246	GLY	2.5
1	A	235	LYS	2.4
1	A	354	VAL	2.4
1	A	93	GLY	2.3
1	B	375	TYR	2.3
1	A	151	VAL	2.3
1	A	481	GLU	2.3
1	B	149	LEU	2.3
1	A	149	LEU	2.3
1	B	128	GLU	2.3
1	B	450	MET	2.2
1	B	345	VAL	2.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	456	VAL	2.2
1	A	231	ARG	2.2
1	B	14	VAL	2.2
1	A	409	GLY	2.2
1	B	127	THR	2.2
1	B	353	ILE	2.1
1	A	464	GLY	2.1
1	B	524	THR	2.1
1	B	364	LYS	2.1
1	A	136	LEU	2.1
1	B	212	LEU	2.1
1	B	144	PRO	2.1
1	A	121	GLU	2.1
1	A	43	LEU	2.1
1	A	394	HIS	2.1
1	A	171	MET	2.1
1	A	295	GLY	2.1
1	B	451	ASP	2.0
1	A	35	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	JE4	A	601	27/27	0.84	0.19	43,51,58,67	0
3	PO4	A	603	5/5	0.85	0.37	113,113,114,114	0
2	JE4	B	601	27/27	0.86	0.20	42,51,59,69	0
3	PO4	A	602	5/5	0.92	0.16	103,103,104,105	0

*Continued on next page...*




*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	602	5/5	0.97	0.09	63,63,64,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around JE4 A 601:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.