



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 08:18 pm BST

PDB ID : 1ME0
Title : Chimeric hairpin with 2',5'-linked RNA loop and DNA stem
Authors : Denisov, A.Y.; Hannoush, R.N.; Gehring, K.; Damha, M.J.
Deposited on : 2002-08-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	12	 25% 42% 33%

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 377 atoms, of which 133 are hydrogens and 0 are deuteriums.

- Molecule 1 is DNA/RNA hybrid called DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3').


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	12	377	114	133	43	76	11	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')



4.2.2 Score per residue for model 2

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')



4.2.3 Score per residue for model 3


- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*G P*TP*CP*C)-3')

Chain A: 



4.2.4 Score per residue for model 4

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*G P*TP*CP*C)-3')

Chain A: 



4.2.5 Score per residue for model 5

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*G P*TP*CP*C)-3')


Chain A: 



4.2.6 Score per residue for model 6


- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*G P*TP*CP*C)-3')

Chain A: 



4.2.7 Score per residue for model 7

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*G P*TP*CP*C)-3')




4.2.8 Score per residue for model 8

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')




4.2.9 Score per residue for model 9

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')



4.2.10 Score per residue for model 10

- Molecule 1: DNA/RNA (5'-D(*GP*GP*AP*C)-R(P*(U25)P*(U25)P*(C25)P*(5GP))-D(P*GP*TP*CP*C)-3')



5 Refinement protocol and experimental data overview [i](#)

The models were refined using the following method: *simulated annealing*.

Of the 20 calculated structures, 10 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	244	133	133	6±2
All	All	2440	1330	1330	59

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:5GP:H8	1:A:8:5GP:O5'	0.66	1.91	2	5
1:A:8:5GP:O5'	1:A:8:5GP:H8	0.60	1.96	8	3
1:A:3:DA:C5	1:A:4:DC:C4	0.53	2.97	7	4
1:A:8:5GP:C6	1:A:9:DG:C5	0.52	2.92	2	8
1:A:8:5GP:C2	1:A:9:DG:C4	0.51	2.94	9	5
1:A:2:DG:H2''	1:A:3:DA:O5'	0.49	2.08	9	5
1:A:3:DA:H2''	1:A:4:DC:O5'	0.46	2.10	3	3
1:A:6:U5P:O4'	1:A:8:5GP:C2	0.46	2.63	4	1
1:A:5:U5P:C4	1:A:6:U5P:C4	0.45	2.95	2	3
1:A:1:DG:C2	1:A:2:DG:C5	0.45	3.04	5	2
1:A:8:5GP:N3	1:A:9:DG:C8	0.44	2.86	9	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:5GP:C2	1:A:9:DG:N9	0.44	2.80	4	1
1:A:8:5GP:N1	1:A:9:DG:C4	0.43	2.86	3	4
1:A:4:DC:H2''	1:A:5:U5P:O5'	0.43	2.14	6	3
1:A:11:DC:H2''	1:A:12:DC:O5'	0.43	2.12	3	5
1:A:8:5GP:N2	1:A:9:DG:C1'	0.42	2.82	4	1
1:A:9:DG:H2''	1:A:10:DT:O5'	0.42	2.14	7	2
1:A:3:DA:C6	1:A:4:DC:C4	0.42	3.08	7	1
1:A:8:5GP:N1	1:A:9:DG:C5	0.41	2.88	9	1
1:A:6:U5P:O4'	1:A:8:5GP:N1	0.40	2.54	2	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.2.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	3-A	4
1	4-A	4
1	5-A	4
1	8-A	4
1	1-A	4
1	6-A	4
1	2-A	4
1	10-A	4
1	9-A	4
1	7-A	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	6:U5P	O3'	7:C5P	P	4.18
2	A	6:U5P	O3'	7:C5P	P	4.18
4	A	6:U5P	O3'	7:C5P	P	4.18
5	A	6:U5P	O3'	7:C5P	P	4.18
6	A	6:U5P	O3'	7:C5P	P	4.18
3	A	6:U5P	O3'	7:C5P	P	4.17
8	A	6:U5P	O3'	7:C5P	P	4.17
9	A	6:U5P	O3'	7:C5P	P	4.17
10	A	6:U5P	O3'	7:C5P	P	4.17
7	A	6:U5P	O3'	7:C5P	P	4.16
9	A	8:5GP	O3'	9:DG	P	3.82
5	A	8:5GP	O3'	9:DG	P	3.77
4	A	8:5GP	O3'	9:DG	P	3.76
8	A	8:5GP	O3'	9:DG	P	3.67
3	A	8:5GP	O3'	9:DG	P	3.56
10	A	8:5GP	O3'	9:DG	P	3.55
6	A	8:5GP	O3'	9:DG	P	3.44
7	A	8:5GP	O3'	9:DG	P	3.41
2	A	8:5GP	O3'	9:DG	P	3.40
8	A	5:U5P	O3'	6:U5P	P	3.40
2	A	5:U5P	O3'	6:U5P	P	3.39

Continued on next page...

Continued from previous page...

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
9	A	5:U5P	O3'	6:U5P	P	3.37
1	A	5:U5P	O3'	6:U5P	P	3.35
3	A	5:U5P	O3'	6:U5P	P	3.34
1	A	8:5GP	O3'	9:DG	P	3.33
10	A	5:U5P	O3'	6:U5P	P	3.29
4	A	5:U5P	O3'	6:U5P	P	3.28
7	A	5:U5P	O3'	6:U5P	P	3.28
10	A	7:C5P	O3'	8:5GP	P	3.26
3	A	7:C5P	O3'	8:5GP	P	3.25
5	A	5:U5P	O3'	6:U5P	P	3.17
6	A	5:U5P	O3'	6:U5P	P	3.17
8	A	7:C5P	O3'	8:5GP	P	3.17
5	A	7:C5P	O3'	8:5GP	P	3.14
1	A	7:C5P	O3'	8:5GP	P	3.12
2	A	7:C5P	O3'	8:5GP	P	3.12
4	A	7:C5P	O3'	8:5GP	P	3.12
6	A	7:C5P	O3'	8:5GP	P	3.12
7	A	7:C5P	O3'	8:5GP	P	3.11
9	A	7:C5P	O3'	8:5GP	P	3.11

6 Chemical shift validation

No chemical shift data were provided