



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:11 am BST

PDB ID : 3ME2
Title : Crystal structure of mouse RANKL-RANK complex
Authors : Walter, S.W.; Liu, C.Z.; Zhu, X.K.; Wu, Y.; Owens, R.J.; Stuart, D.I.; Gao, B.; Ren, J.
Deposited on : 2010-03-31
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	
2	R	216	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	0	0
			1232	787	210	231	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	GLY	-	EXPRESSION TAG	UNP O35235
A	147	PRO	-	EXPRESSION TAG	UNP O35235
A	148	LEU	-	EXPRESSION TAG	UNP O35235
A	149	GLY	-	EXPRESSION TAG	UNP O35235
A	150	SER	-	EXPRESSION TAG	UNP O35235
A	151	PRO	-	EXPRESSION TAG	UNP O35235
A	152	GLU	-	EXPRESSION TAG	UNP O35235
A	153	PHE	-	EXPRESSION TAG	UNP O35235
A	154	PRO	-	EXPRESSION TAG	UNP O35235
A	155	GLY	-	EXPRESSION TAG	UNP O35235
A	156	ARG	-	EXPRESSION TAG	UNP O35235
A	157	ARG	-	EXPRESSION TAG	UNP O35235

- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member 11A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	R	166	Total	C	N	O	S	0	0	0
			1262	769	223	249	21			

There are 31 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	3	MET	-	EXPRESSION TAG	UNP O35305
R	4	GLY	-	EXPRESSION TAG	UNP O35305
R	5	SER	-	EXPRESSION TAG	UNP O35305
R	6	SER	-	EXPRESSION TAG	UNP O35305

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	7	HIS	-	EXPRESSION TAG	UNP O35305
R	8	HIS	-	EXPRESSION TAG	UNP O35305
R	9	HIS	-	EXPRESSION TAG	UNP O35305
R	10	HIS	-	EXPRESSION TAG	UNP O35305
R	11	HIS	-	EXPRESSION TAG	UNP O35305
R	12	HIS	-	EXPRESSION TAG	UNP O35305
R	13	SER	-	EXPRESSION TAG	UNP O35305
R	14	SER	-	EXPRESSION TAG	UNP O35305
R	15	GLY	-	EXPRESSION TAG	UNP O35305
R	16	LEU	-	EXPRESSION TAG	UNP O35305
R	17	VAL	-	EXPRESSION TAG	UNP O35305
R	18	PRO	-	EXPRESSION TAG	UNP O35305
R	19	ARG	-	EXPRESSION TAG	UNP O35305
R	20	GLY	-	EXPRESSION TAG	UNP O35305
R	21	SER	-	EXPRESSION TAG	UNP O35305
R	22	HIS	-	EXPRESSION TAG	UNP O35305
R	23	MET	-	EXPRESSION TAG	UNP O35305
R	24	HIS	-	EXPRESSION TAG	UNP O35305
R	25	MET	-	EXPRESSION TAG	UNP O35305
R	211	LEU	-	EXPRESSION TAG	UNP O35305
R	212	GLU	-	EXPRESSION TAG	UNP O35305
R	213	HIS	-	EXPRESSION TAG	UNP O35305
R	214	HIS	-	EXPRESSION TAG	UNP O35305
R	215	HIS	-	EXPRESSION TAG	UNP O35305
R	216	HIS	-	EXPRESSION TAG	UNP O35305
R	217	HIS	-	EXPRESSION TAG	UNP O35305
R	218	HIS	-	EXPRESSION TAG	UNP O35305

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	R	1	Total Cl 1 1	0	0
3	A	3	Total Cl 3 3	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	R	1	Total Na 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	23	Total 23	O 23	0	0
5	R	22	Total 22	O 22	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	121.22Å 121.22Å 95.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.80 29.12 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.80) 98.8 (29.12-2.78)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.174 , 0.205 0.178 , 0.169	Depositor DCC
R_{free} test set	1005 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	48.7	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.054 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2544	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1267	0.52	0/1718
2	R	0.34	0/1293	0.50	0/1756
All	All	0.36	0/2560	0.51	0/3474

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	233	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1232	0	1182	7	0
2	R	1262	0	1163	5	0
3	A	3	0	0	1	0
3	R	1	0	0	0	0
4	R	1	0	0	0	0
5	A	23	0	0	0	0
5	R	22	0	0	0	0
All	All	2544	0	2345	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:62:THR:HB	2:R:63:PRO:HD2	1.88	0.55
1:A:211:GLY:HA3	1:A:311:LYS:HE3	1.89	0.55
2:R:103:ASP:HB3	2:R:112:ARG:HB2	1.95	0.49
1:A:181:VAL:HG22	1:A:295:PRO:HG3	1.96	0.48
1:A:244:THR:HG22	1:A:245:SER:H	1.81	0.45
1:A:200:LEU:HD11	3:A:5:CL:CL	2.55	0.44
2:R:185:HIS:HB3	2:R:194:VAL:CG2	2.48	0.43
1:A:281:LYS:HE3	1:A:315:ILE:HD12	2.01	0.42
1:A:162:GLN:HA	1:A:163:PRO:HD3	1.77	0.41
2:R:160:PHE:CE2	2:R:171:LYS:HG3	2.56	0.41
2:R:185:HIS:HB3	2:R:194:VAL:HG22	2.02	0.41
1:A:301:ASP:HB3	1:A:303:ASP:OD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/171 (90%)	147 (96%)	6 (4%)	1 (1%)	25	56
2	R	164/216 (76%)	159 (97%)	5 (3%)	0	100	100
All	All	318/387 (82%)	306 (96%)	11 (4%)	1 (0%)	41	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	231	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/147 (92%)	131 (97%)	4 (3%)	41	75
2	R	146/191 (76%)	142 (97%)	4 (3%)	44	78
All	All	281/338 (83%)	273 (97%)	8 (3%)	43	77

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	LEU
1	A	222	ARG
1	A	244	THR
1	A	316	ASP
2	R	108	THR
2	R	157	LEU
2	R	169	LYS
2	R	171	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	262	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/171 (91%)	-0.16	4 (2%) 56 46	28, 43, 89, 106	0
2	R	166/216 (76%)	0.08	9 (5%) 25 17	42, 60, 121, 177	0
All	All	322/387 (83%)	-0.04	13 (4%) 38 28	28, 54, 102, 177	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	R	36	THR	5.0
2	R	35	CYS	4.9
1	A	316	ASP	4.8
2	R	37	GLN	4.6
2	R	47	CYS	4.3
1	A	161	ALA	3.6
2	R	38	GLU	3.5
2	R	39	ARG	2.9
2	R	157	LEU	2.6
2	R	34	PRO	2.6
1	A	232	THR	2.5
2	R	44	LEU	2.5
1	A	231	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	A	6	1/1	0.95	0.10	52,52,52,52	0
3	CL	R	219	1/1	0.97	0.08	52,52,52,52	0
4	NA	R	1	1/1	0.98	0.04	36,36,36,36	0
3	CL	A	5	1/1	0.98	0.22	69,69,69,69	0
3	CL	A	7	1/1	1.00	0.16	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.