



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:33 am BST

PDB ID : 4MET  
Title : Assigning the EPR Fine Structure Parameters of the Mn(II) Centers in *Bacillus subtilis* Oxalate Decarboxylase by Site-Directed Mutagenesis and DFT/MM Calculations  
Authors : Campomanes, P.; Kellett, W.F.; Easthon, L.M.; Ozarowski, A.; Allen, K.N.; Angerhofer, A.; Rothlisberger, U.; Richards, N.G.J.  
Deposited on : 2013-08-27  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div>4%</div> <div>93%</div> <div>5%</div> <div>• • •</div> </div>
1	B	388	<div> <div>2%</div> <div>92%</div> <div>5%</div> <div>•</div> </div>
1	C	388	<div> <div>4%</div> <div>91%</div> <div>5%</div> <div>•</div> </div>
1	D	388	<div> <div>2%</div> <div>91%</div> <div>6%</div> <div>•</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxalate decarboxylase OxdC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			2997	1914	503	572	8			
1	B	377	Total	C	N	O	S	0	0	0
			2997	1914	503	572	8			
1	C	377	Total	C	N	O	S	0	0	0
			2997	1914	503	572	8			
1	D	377	Total	C	N	O	S	0	0	0
			2997	1914	503	572	8			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	PHE	TRP	engineered mutation	UNP O34714
A	383	HIS	-	expression tag	UNP O34714
A	384	HIS	-	expression tag	UNP O34714
A	385	HIS	-	expression tag	UNP O34714
A	386	HIS	-	expression tag	UNP O34714
A	387	HIS	-	expression tag	UNP O34714
A	388	HIS	-	expression tag	UNP O34714
B	132	PHE	TRP	engineered mutation	UNP O34714
B	383	HIS	-	expression tag	UNP O34714
B	384	HIS	-	expression tag	UNP O34714
B	385	HIS	-	expression tag	UNP O34714
B	386	HIS	-	expression tag	UNP O34714
B	387	HIS	-	expression tag	UNP O34714
B	388	HIS	-	expression tag	UNP O34714
C	132	PHE	TRP	engineered mutation	UNP O34714
C	383	HIS	-	expression tag	UNP O34714
C	384	HIS	-	expression tag	UNP O34714
C	385	HIS	-	expression tag	UNP O34714
C	386	HIS	-	expression tag	UNP O34714
C	387	HIS	-	expression tag	UNP O34714
C	388	HIS	-	expression tag	UNP O34714

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	132	PHE	TRP	engineered mutation	UNP O34714
D	383	HIS	-	expression tag	UNP O34714
D	384	HIS	-	expression tag	UNP O34714
D	385	HIS	-	expression tag	UNP O34714
D	386	HIS	-	expression tag	UNP O34714
D	387	HIS	-	expression tag	UNP O34714
D	388	HIS	-	expression tag	UNP O34714

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Co 2 2	0	0
2	A	2	Total Co 2 2	0	0
2	D	2	Total Co 2 2	0	0
2	C	2	Total Co 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	97	Total O 97 97	0	0
3	B	116	Total O 116 116	0	0
3	C	79	Total O 79 79	0	0
3	D	97	Total O 97 97	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Oxalate decarboxylase OxdC




- Molecule 1: Oxalate decarboxylase OxdC



- Molecule 1: Oxalate decarboxylase OxdC



- Molecule 1: Oxalate decarboxylase OxdC



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.97Å 156.97Å 330.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.39 – 2.10 49.06 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.39-2.10) 94.3 (49.06-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.196 , 0.227 0.198 , 0.228	Depositor DCC
$R_{free}$ test set	2000 reflections (2.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 31.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12385	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 65.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.0239e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/3079	0.56	0/4182
1	B	0.43	0/3079	0.58	0/4182
1	C	0.40	0/3079	0.52	0/4182
1	D	0.41	0/3079	0.55	0/4182
All	All	0.42	0/12316	0.55	0/16728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2997	0	2873	13	0
1	B	2997	0	2873	8	0
1	C	2997	0	2873	12	0
1	D	2997	0	2873	14	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	97	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	116	0	0	0	0
3	C	79	0	0	0	0
3	D	97	0	0	2	0
All	All	12385	0	11492	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:SER:HB2	1:C:164:SER:HB3	1.72	0.70
1:B:8:PRO:HD2	1:B:25:ILE:HD11	1.73	0.68
1:B:162:GLU:HG2	1:B:200:TYR:OH	1.94	0.67
1:A:167:GLN:NE2	1:A:198:GLU:OE1	2.29	0.66
1:B:161:SER:OG	1:B:164:SER:HB3	2.00	0.62
1:A:160:PHE:HZ	1:A:162:GLU:HG3	1.68	0.57
1:D:162:GLU:HG2	1:D:200:TYR:OH	2.05	0.56
1:A:161:SER:HB3	3:A:687:HOH:O	2.05	0.55
1:A:161:SER:OG	1:A:164:SER:HB3	2.08	0.54
1:D:165:THR:HA	3:D:696:HOH:O	2.07	0.53
1:D:67:GLU:HB3	1:D:82:VAL:HG22	1.91	0.53
1:C:164:SER:OG	1:C:164:SER:O	2.28	0.51
1:A:197:LYS:NZ	1:A:198:GLU:O	2.37	0.51
1:D:197:LYS:HG3	3:D:692:HOH:O	2.11	0.50
1:D:12:ARG:NH1	1:D:37:PRO:O	2.45	0.50
1:D:238:SER:HB3	1:D:378:VAL:O	2.12	0.50
1:C:238:SER:HB3	1:C:378:VAL:O	2.11	0.49
1:C:160:PHE:CE2	1:C:162:GLU:HB2	2.47	0.49
1:D:82:VAL:HG21	1:D:162:GLU:OE1	2.13	0.49
1:A:8:PRO:HD2	1:A:25:ILE:HD11	1.96	0.48
1:B:146:GLU:CD	1:B:146:GLU:H	2.18	0.47
1:A:160:PHE:CZ	1:A:162:GLU:HG3	2.49	0.47
1:A:6:ASP:O	1:A:29:ARG:NH1	2.48	0.46
1:B:231:LEU:HD21	1:B:261:LEU:HD23	1.97	0.46
1:C:111:ARG:NH1	1:C:213:ASP:OD2	2.45	0.46
1:D:231:LEU:HA	1:D:231:LEU:HD23	1.77	0.45
1:C:67:GLU:HB3	1:C:82:VAL:HG22	1.99	0.45
1:D:160:PHE:HE2	1:D:162:GLU:HB2	1.82	0.45
1:A:6:ASP:O	1:A:29:ARG:NH2	2.48	0.45
1:A:129:GLY:HA3	1:A:231:LEU:HD12	1.98	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:160:PHE:HE2	1:C:162:GLU:HB2	1.82	0.44
1:D:146:GLU:HG3	1:D:206:LEU:HD13	1.98	0.44
1:B:49:LYS:NZ	1:B:304:ASN:O	2.48	0.44
1:A:162:GLU:O	3:A:628:HOH:O	2.20	0.44
1:C:244:TYR:O	1:C:260:ALA:HA	2.17	0.44
1:D:231:LEU:HD21	1:D:261:LEU:HD22	2.00	0.43
1:A:197:LYS:HB2	1:A:197:LYS:HE2	1.71	0.43
1:A:14:ASP:N	1:A:14:ASP:OD1	2.51	0.43
1:B:367:PHE:O	1:B:370:VAL:HG22	2.17	0.43
1:B:161:SER:HG	1:B:164:SER:HB3	1.82	0.43
1:C:154:VAL:HG11	1:C:334:ILE:HG21	2.00	0.43
1:D:160:PHE:CE2	1:D:162:GLU:HB2	2.55	0.42
1:C:116:ASP:HA	1:C:138:LEU:HD13	2.02	0.41
1:C:292:THR:HB	1:C:320:TYR:HE2	1.86	0.41
1:D:367:PHE:O	1:D:370:VAL:HG22	2.21	0.40
1:C:111:ARG:NH1	1:C:207:PRO:HB2	2.35	0.40
1:D:10:PRO:HA	1:D:36:VAL:HG22	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/388 (97%)	366 (98%)	9 (2%)	0	100	100
1	B	375/388 (97%)	369 (98%)	6 (2%)	0	100	100
1	C	375/388 (97%)	363 (97%)	12 (3%)	0	100	100
1	D	375/388 (97%)	365 (97%)	9 (2%)	1 (0%)	41	41
All	All	1500/1552 (97%)	1463 (98%)	36 (2%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	164	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/336 (96%)	317 (99%)	4 (1%)	71	77
1	B	321/336 (96%)	314 (98%)	7 (2%)	52	57
1	C	321/336 (96%)	317 (99%)	4 (1%)	71	77
1	D	321/336 (96%)	318 (99%)	3 (1%)	78	84
All	All	1284/1344 (96%)	1266 (99%)	18 (1%)	67	73

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	SER
1	A	97	HIS
1	A	161	SER
1	A	167	GLN
1	B	26	GLU
1	B	27	ARG
1	B	83	ASN
1	B	97	HIS
1	B	167	GLN
1	B	203	GLU
1	B	342	ASP
1	C	27	ARG
1	C	83	ASN
1	C	161	SER
1	C	167	GLN
1	D	83	ASN
1	D	161	SER
1	D	167	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	377/388 (97%)	-0.20	15 (3%) 38 44	9, 17, 35, 54	0
1	B	377/388 (97%)	-0.36	7 (1%) 66 71	9, 15, 31, 47	0
1	C	377/388 (97%)	0.12	14 (3%) 41 48	16, 28, 50, 75	0
1	D	377/388 (97%)	-0.06	8 (2%) 63 68	14, 24, 42, 58	0
All	All	1508/1552 (97%)	-0.13	44 (2%) 51 57	9, 21, 44, 75	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	163	ASN	5.6
1	A	216	GLU	4.2
1	C	6	ASP	4.2
1	C	215	VAL	4.1
1	A	164	SER	4.1
1	A	163	ASN	4.0
1	A	215	VAL	4.0
1	C	214	ILE	3.5
1	D	163	ASN	3.2
1	A	214	ILE	3.2
1	B	164	SER	3.1
1	D	162	GLU	3.0
1	B	163	ASN	3.0
1	A	239	GLU	2.9
1	A	381	LYS	2.9
1	C	239	GLU	2.8
1	C	216	GLU	2.8
1	B	382	LYS	2.8
1	B	161	SER	2.8
1	C	164	SER	2.8
1	D	6	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	216	GLU	2.7
1	B	162	GLU	2.5
1	C	366	ASP	2.5
1	C	162	GLU	2.5
1	D	189	GLU	2.5
1	A	382	LYS	2.5
1	D	164	SER	2.5
1	A	162	GLU	2.4
1	B	197	LYS	2.4
1	D	374	GLU	2.4
1	A	161	SER	2.3
1	A	366	ASP	2.3
1	A	212	ASP	2.3
1	A	211	LYS	2.2
1	A	326	ASP	2.2
1	C	374	GLU	2.2
1	A	210	LEU	2.1
1	C	367	PHE	2.1
1	D	214	ILE	2.1
1	C	218	PRO	2.1
1	D	146	GLU	2.0
1	C	212	ASP	2.0
1	C	382	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CO	C	500	1/1	0.99	0.14	24,24,24,24	0
2	CO	D	500	1/1	0.99	0.13	20,20,20,20	0
2	CO	D	501	1/1	1.00	0.13	18,18,18,18	0
2	CO	A	500	1/1	1.00	0.14	20,20,20,20	0
2	CO	B	500	1/1	1.00	0.15	17,17,17,17	0
2	CO	B	501	1/1	1.00	0.10	15,15,15,15	0
2	CO	C	501	1/1	1.00	0.10	22,22,22,22	0
2	CO	A	501	1/1	1.00	0.12	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.