



Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 09:52 pm BST

PDB ID : 5MEF
Title : Cyanothece lipxygenase 2 (CspLOX2) variant - L304F
Authors : Newie, J.; Neumann, P.; Werner, M.; Mata, R.A.; Ficner, R.; Feussner, I.
Deposited on : 2016-11-14
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	569	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div>.</div> </div> </div>
1	B	569	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IMD	B	1004	-	-	-	X
4	IMD	B	1005	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9503 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arachidonate 15-lipoxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	569	Total	C	N	O	S	0	0	0
			4554	2920	772	846	16			
1	B	566	Total	C	N	O	S	0	0	0
			4531	2905	769	841	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	304	PHE	LEU	engineered mutation	UNP B7JX99
B	304	PHE	LEU	engineered mutation	UNP B7JX99

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N 5 3 2	0	0
4	A	1	Total C N 5 3 2	0	0
4	B	1	Total C N 5 3 2	0	0
4	B	1	Total C N 5 3 2	0	0
4	B	1	Total C N 5 3 2	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0
5	A	2	Total Cl 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	192	Total O 192 192	0	0
6	B	172	Total O 172 172	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Arachidonate 15-lipoxygenase



• Molecule 1: Arachidonate 15-lipoxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.40 Å 166.26 Å 166.74 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.20 – 2.36 46.20 – 2.36	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.20-2.36) 98.4 (46.20-2.36)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.37 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.191 , 0.216 0.192 , 0.217	Depositor DCC
R_{free} test set	3149 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	52.3	Xtriage
Anisotropy	0.322	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 23.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.449 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9503	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, FE, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/4662	0.47	0/6346
1	B	0.27	0/4637	0.44	0/6311
All	All	0.27	0/9299	0.46	0/12657

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4554	0	4548	43	1
1	B	4531	0	4531	40	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	16	1	0
3	B	12	0	16	0	0
4	A	10	0	10	1	0
4	B	15	0	15	1	0
5	A	2	0	0	0	0
5	B	1	0	0	0	0
6	A	192	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	172	0	0	3	0
All	All	9503	0	9136	83	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:PRO:HA	1:A:418:ARG:HG2	1.52	0.90
1:B:414:PRO:HA	1:B:418:ARG:HG3	1.55	0.87
1:B:109:ALA:HB1	1:B:210:LEU:HD23	1.59	0.82
1:B:395:PRO:HB2	1:B:399:ARG:HH21	1.48	0.77
1:B:3:GLN:NE2	1:B:523:ASP:OD2	2.22	0.72
1:A:165:GLY:O	6:A:1101:HOH:O	2.10	0.69
1:A:521:PHE:O	6:A:1102:HOH:O	2.11	0.67
1:A:410:ARG:O	1:A:413:SER:OG	2.13	0.64
1:A:72:ASN:ND2	6:A:1104:HOH:O	2.19	0.64
1:B:264:ILE:HD11	1:B:444:VAL:HG11	1.81	0.62
1:A:88:GLU:OE1	6:A:1103:HOH:O	2.16	0.62
1:A:264:ILE:HD11	1:A:444:VAL:HG11	1.83	0.61
1:A:147:LYS:NZ	1:A:155:GLN:OE1	2.33	0.60
1:B:285:LEU:HB2	1:B:387:TYR:CE2	2.37	0.58
1:B:257:HIS:O	1:B:261:THR:OG1	2.22	0.57
1:B:451:ALA:HB2	1:B:558:LEU:HD21	1.87	0.56
1:A:508:VAL:HG13	1:A:509:PHE:HD1	1.70	0.56
1:A:117:ASN:HB2	1:A:252:HIS:HB2	1.89	0.55
1:B:3:GLN:HE22	1:B:523:ASP:CG	2.10	0.54
1:A:201:GLU:O	1:A:205:ASP:O	2.26	0.54
1:A:346:LEU:HD13	1:A:370:ARG:HG3	1.88	0.54
1:B:410:ARG:O	1:B:416:GLY:HA3	2.08	0.53
1:A:284:VAL:HG12	1:A:521:PHE:CE1	2.43	0.53
1:A:561:SER:O	4:A:1005:IMD:H2	2.10	0.52
1:B:346:LEU:HD13	1:B:370:ARG:HG3	1.91	0.52
1:A:110:ARG:NH1	1:A:366:ASP:OD2	2.43	0.52
1:A:541:GLU:O	1:A:545:VAL:HG23	2.10	0.52
1:A:284:VAL:HG12	1:A:521:PHE:HE1	1.76	0.51
1:A:318:THR:HG22	3:A:1003:GOL:H32	1.92	0.51
1:A:285:LEU:HB2	1:A:387:TYR:CE2	2.46	0.51
1:A:153:ILE:HG13	1:A:158:LEU:HD12	1.93	0.50
1:A:520:GLU:HG2	1:A:527:GLN:NE2	2.26	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:529:VAL:HA	1:A:532:GLN:HG3	1.94	0.49
1:B:71:SER:HB3	1:B:73:ILE:HG22	1.95	0.49
1:A:253:GLU:OE2	1:A:456:GLN:NE2	2.45	0.49
1:A:444:VAL:HA	1:A:448:GLN:HB3	1.94	0.48
1:B:206:TYR:N	1:B:206:TYR:CD1	2.81	0.48
1:A:272:THR:HG23	1:A:276:LEU:HD12	1.94	0.48
1:B:223:PRO:O	1:B:477:LYS:NZ	2.33	0.47
1:A:3:GLN:NE2	1:A:523:ASP:OD2	2.49	0.46
1:B:257:HIS:CE1	6:B:1174:HOH:O	2.67	0.46
1:B:33:GLU:OE2	1:B:36:GLY:N	2.31	0.46
1:B:117:ASN:HB2	1:B:252:HIS:HB2	1.99	0.45
1:B:444:VAL:HA	1:B:448:GLN:HB3	1.97	0.45
1:B:265:LEU:HD11	1:B:445:SER:HA	1.98	0.45
1:B:380:ILE:O	1:B:384:VAL:HG13	2.17	0.45
1:A:455:PRO:HG3	1:A:557:TYR:CD2	2.52	0.44
1:B:463:ILE:HG13	1:B:496:ALA:HB1	1.99	0.44
1:A:73:ILE:HD13	1:A:497:ALA:HB2	1.98	0.44
1:A:528:GLN:O	1:A:532:GLN:HG3	2.18	0.44
1:A:22:ILE:O	1:A:26:ARG:HG3	2.18	0.44
1:B:410:ARG:O	1:B:413:SER:OG	2.32	0.43
1:B:127:LYS:NZ	6:B:1121:HOH:O	2.51	0.43
1:B:269:VAL:HG22	1:B:290:PHE:CG	2.54	0.43
1:B:262:HIS:HB3	1:B:297:ASN:OD1	2.19	0.43
1:B:1:MET:HG3	1:B:1:MET:O	2.18	0.43
1:A:262:HIS:HB3	1:A:297:ASN:OD1	2.18	0.43
1:B:409:ARG:HA	1:B:412:VAL:HG22	2.01	0.43
1:B:272:THR:HG23	1:B:276:LEU:HD12	2.01	0.43
1:A:260:ARG:HA	1:A:264:ILE:HG12	2.00	0.42
1:B:562:ASN:OD1	4:B:1004:IMD:H2	2.19	0.42
1:B:504:ARG:O	1:B:508:VAL:HG22	2.19	0.42
1:B:353:ILE:HG23	6:B:1198:HOH:O	2.19	0.42
1:A:4:PRO:HG2	6:A:1125:HOH:O	2.19	0.42
1:B:134:VAL:HG12	1:B:149:LEU:HD11	2.01	0.42
1:A:171:PRO:HB3	1:A:189:ILE:HG12	2.02	0.42
1:B:73:ILE:HD13	1:B:497:ALA:HB2	2.01	0.42
1:B:394:SER:OG	1:B:396:VAL:HG12	2.20	0.41
1:A:451:ALA:HB2	1:A:558:LEU:HD21	2.02	0.41
1:A:16:GLU:OE2	1:A:20:ARG:NH2	2.54	0.41
1:A:523:ASP:OD1	1:A:525:ARG:N	2.53	0.41
1:B:142:ILE:HD11	1:B:230:THR:HB	2.01	0.41
1:A:106:ARG:HG3	1:A:203:PHE:CE2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:521:PHE:N	1:A:527:GLN:OE1	2.49	0.41
1:A:112:ARG:O	1:A:118:PRO:HA	2.20	0.41
1:A:269:VAL:HG22	1:A:290:PHE:CG	2.56	0.41
1:A:138:MET:O	1:A:142:ILE:HD13	2.21	0.41
1:B:73:ILE:HD11	1:B:493:GLN:O	2.21	0.41
1:B:206:TYR:HB2	1:B:207:ARG:H	1.76	0.40
1:A:101:THR:HB	1:A:107:VAL:HG21	2.03	0.40
1:B:333:TYR:HA	1:B:340:PHE:CE2	2.56	0.40
1:B:455:PRO:HG3	1:B:557:TYR:CD2	2.56	0.40
1:B:395:PRO:HB2	1:B:399:ARG:NH2	2.25	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:GLU:OE2	1:B:399:ARG:NH1[3_655]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	567/569 (100%)	555 (98%)	12 (2%)	0	100	100
1	B	562/569 (99%)	549 (98%)	12 (2%)	1 (0%)	47	56
All	All	1129/1138 (99%)	1104 (98%)	24 (2%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/493 (100%)	478 (97%)	15 (3%)	41	50
1	B	491/493 (100%)	475 (97%)	16 (3%)	38	46
All	All	984/986 (100%)	953 (97%)	31 (3%)	39	47

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	THR
1	A	84	GLU
1	A	106	ARG
1	A	110	ARG
1	A	144	PHE
1	A	146	LYS
1	A	147	LYS
1	A	201	GLU
1	A	205	ASP
1	A	301	ARG
1	A	304	PHE
1	A	367	TYR
1	A	393	GLN
1	A	513	ARG
1	A	524	VAL
1	B	84	GLU
1	B	178	LYS
1	B	179	ASN
1	B	181	THR
1	B	201	GLU
1	B	206	TYR
1	B	210	LEU
1	B	304	PHE
1	B	367	TYR
1	B	384	VAL
1	B	415	GLU
1	B	423	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	476	ASN
1	B	478	GLU
1	B	508	VAL
1	B	538	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	542	GLN
1	B	3	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 5 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1002	-	5,5,5	0.36	0	5,5,5	0.95	0
4	IMD	A	1005	-	3,5,5	0.39	0	4,5,5	0.54	0
3	GOL	B	1003	-	5,5,5	0.64	0	5,5,5	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IMD	B	1004	-	3,5,5	0.28	0	4,5,5	0.53	0
4	IMD	A	1004	-	3,5,5	0.49	0	4,5,5	0.62	0
4	IMD	B	1005	-	3,5,5	0.43	0	4,5,5	0.62	0
4	IMD	B	1006	-	3,5,5	0.43	0	4,5,5	0.61	0
3	GOL	A	1003	-	5,5,5	0.30	0	5,5,5	0.31	0
3	GOL	B	1002	-	5,5,5	0.29	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1002	-	-	0/4/4/4	-
4	IMD	A	1005	-	-	-	0/1/1/1
3	GOL	B	1003	-	-	4/4/4/4	-
4	IMD	B	1004	-	-	-	0/1/1/1
4	IMD	A	1004	-	-	-	0/1/1/1
4	IMD	B	1005	-	-	-	0/1/1/1
4	IMD	B	1006	-	-	-	0/1/1/1
3	GOL	A	1003	-	-	0/4/4/4	-
3	GOL	B	1002	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1003	GOL	O1-C1-C2-C3
3	B	1003	GOL	C1-C2-C3-O3
3	B	1003	GOL	O2-C2-C3-O3
3	B	1003	GOL	O1-C1-C2-O2
3	B	1002	GOL	C1-C2-C3-O3
3	B	1002	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1005	IMD	1	0
4	B	1004	IMD	1	0
3	A	1003	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	569/569 (100%)	0.30	14 (2%) 57 67	37, 55, 97, 179	0
1	B	566/569 (99%)	0.27	16 (2%) 53 64	36, 54, 94, 134	0
All	All	1135/1138 (99%)	0.29	30 (2%) 56 65	36, 54, 97, 179	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	6.5
1	A	203	PHE	5.6
1	A	205	ASP	5.1
1	A	206	TYR	5.1
1	A	202	GLY	4.6
1	B	358	GLN	4.0
1	B	550	LYS	3.9
1	A	204	GLY	3.7
1	B	224	ILE	3.4
1	B	1	MET	3.2
1	B	552	LEU	3.1
1	B	20	ARG	3.1
1	A	177	GLN	3.0
1	B	206	TYR	2.8
1	B	391	TYR	2.7
1	B	359	ARG	2.6
1	A	552	LEU	2.6
1	A	394	SER	2.5
1	A	72	ASN	2.5
1	A	431	ILE	2.5
1	A	531	HIS	2.4
1	A	24	GLN	2.4
1	B	98	VAL	2.4
1	B	395	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	113	ILE	2.3
1	B	6	LEU	2.2
1	B	135	THR	2.1
1	B	169	LEU	2.0
1	B	207	ARG	2.0
1	B	296	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	1003	6/6	0.59	0.17	87,90,90,91	0
4	IMD	B	1004	5/5	0.70	0.48	81,81,81,81	0
4	IMD	B	1005	5/5	0.72	0.46	81,81,81,82	0
3	GOL	A	1002	6/6	0.79	0.26	75,77,78,80	0
4	IMD	A	1005	5/5	0.85	0.15	78,78,79,79	0
4	IMD	A	1004	5/5	0.87	0.47	81,81,81,81	0
5	CL	B	1007	1/1	0.87	0.22	61,61,61,61	1
4	IMD	B	1006	5/5	0.88	0.23	83,83,83,83	0
5	CL	A	1006	1/1	0.88	0.18	71,71,71,71	1
5	CL	A	1007	1/1	0.89	0.32	65,65,65,65	1
3	GOL	B	1002	6/6	0.91	0.10	50,54,57,58	0
3	GOL	A	1003	6/6	0.92	0.16	56,57,60,61	0
2	FE	B	1001	1/1	0.94	0.05	65,65,65,65	0
2	FE	A	1001	1/1	0.96	0.11	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.