



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 08:00 am BST

PDB ID : 5MFD  
Title : Designed armadillo repeat protein YIIIM”6AII in complex with pD\_(KR)5  
Authors : Hansen, S.; Kiefer, J.; Madhurantakam, C.; Mittl, P.; Plueckthun, A.  
Deposited on : 2016-11-18  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	<div> <div>91%</div> <div>7% .</div> </div>
1	C	328	<div> <div>%</div> <div>95%</div> <div>. .</div> </div>
1	E	328	<div> <div>95%</div> <div>. . .</div> </div>
1	G	328	<div> <div>%</div> <div>93%</div> <div>5% .</div> </div>
1	I	328	<div> <div>%</div> <div>97%</div> <div>. .</div> </div>
1	J	328	<div> <div>97%</div> <div>. .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	328	
1	L	328	
2	B	109	
2	D	109	
2	F	109	
2	H	109	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YIIIM"6AII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2386	1489	406	490	1			
1	C	323	Total	C	N	O	S	0	0	0
			2386	1489	406	490	1			
1	E	323	Total	C	N	O	S	0	2	0
			2414	1511	410	492	1			
1	G	322	Total	C	N	O	S	0	1	0
			2394	1497	407	489	1			
1	I	323	Total	C	N	O	S	0	0	0
			2386	1489	406	490	1			
1	J	324	Total	C	N	O	S	0	1	0
			2410	1506	411	492	1			
1	K	323	Total	C	N	O	S	0	1	0
			2400	1500	408	491	1			
1	L	323	Total	C	N	O	S	0	2	0
			2409	1506	410	492	1			

- Molecule 2 is a protein called Capsid decoration protein,pD\_(KR)5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	105	Total	C	N	O	S	0	0	0
			777	487	139	149	2			
2	D	107	Total	C	N	O	S	0	0	0
			792	495	144	151	2			
2	F	105	Total	C	N	O	S	0	0	0
			777	487	139	149	2			
2	H	104	Total	C	N	O	S	0	0	0
			768	481	137	148	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	18	GLY	-	expression tag	UNP P03712

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	19	PRO	-	expression tag	UNP P03712
D	18	GLY	-	expression tag	UNP P03712
D	19	PRO	-	expression tag	UNP P03712
F	18	GLY	-	expression tag	UNP P03712
F	19	PRO	-	expression tag	UNP P03712
H	18	GLY	-	expression tag	UNP P03712
H	19	PRO	-	expression tag	UNP P03712

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	9	Total Ca 9 9	0	0
3	J	4	Total Ca 4 4	0	0
3	K	10	Total Ca 10 10	0	0
3	E	9	Total Ca 9 9	0	0
3	H	1	Total Ca 1 1	0	0
3	I	4	Total Ca 4 4	0	0
3	C	1	Total Ca 1 1	0	0
3	A	8	Total Ca 8 8	0	0
3	L	2	Total Ca 2 2	0	0
3	F	1	Total Ca 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	29	Total O 29 29	0	0
4	B	19	Total O 19 19	0	0
4	C	58	Total O 58 58	0	0
4	D	16	Total O 16 16	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	60	Total 60	O 60	0	0
4	F	7	Total 7	O 7	0	0
4	G	27	Total 27	O 27	0	0
4	H	2	Total 2	O 2	0	0
4	I	65	Total 65	O 65	0	0
4	J	61	Total 61	O 61	0	0
4	K	98	Total 98	O 98	0	0
4	L	43	Total 43	O 43	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: YIIIM"6AII

Chain A: 




- Molecule 1: YIIIM"6AII

Chain C: 



- Molecule 1: YIIIM"6AII

Chain E: 



- Molecule 1: YIIIM"6AII

Chain G: 



- Molecule 1: YIIIM"6AII

Chain I: 



- Molecule 1: YIIIM"6AII

Chain J:  97% ..




• Molecule 1: YIIIM"6AII

Chain K:  95% ..





• Molecule 1: YIIIM"6AII

Chain L:  91% 6% ..




• Molecule 2: Capsid decoration protein,pD\_(KR)5

Chain B:  3% 86% 8% ..




• Molecule 2: Capsid decoration protein,pD\_(KR)5

Chain D:  10% 92% 6% ..





• Molecule 2: Capsid decoration protein,pD\_(KR)5

Chain F:  2% 91% 6% ..



• Molecule 2: Capsid decoration protein,pD\_(KR)5

Chain H:  28% 89% 5% ..





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	194.66Å 194.66Å 241.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.12 – 2.30 49.12 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.12-2.30) 86.5 (49.12-2.10)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 2.10Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.190 , 0.214 0.206 , 0.233	Depositor DCC
$R_{free}$ test set	13207 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.7	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	22833	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/2412	0.59	0/3289
1	C	0.43	0/2412	0.60	0/3289
1	E	0.43	0/2444	0.60	0/3335
1	G	0.42	0/2422	0.59	0/3304
1	I	0.43	0/2412	0.60	0/3289
1	J	0.43	0/2439	0.59	0/3327
1	K	0.44	0/2428	0.61	0/3312
1	L	0.43	0/2437	0.59	0/3323
2	B	0.47	0/791	0.66	0/1069
2	D	0.40	0/806	0.60	0/1089
2	F	0.44	0/791	0.62	0/1069
2	H	0.43	0/782	0.59	0/1058
All	All	0.43	0/22576	0.60	0/30753

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2386	0	2399	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2386	0	2399	3	0
1	E	2414	0	2416	5	0
1	G	2394	0	2403	10	0
1	I	2386	0	2399	2	0
1	J	2410	0	2415	1	0
1	K	2400	0	2407	4	0
1	L	2409	0	2420	12	0
2	B	777	0	801	8	0
2	D	792	0	816	6	0
2	F	777	0	801	2	0
2	H	768	0	788	10	0
3	A	8	0	0	0	0
3	C	1	0	0	0	0
3	E	9	0	0	0	0
3	F	1	0	0	0	0
3	G	9	0	0	0	0
3	H	1	0	0	0	0
3	I	4	0	0	0	0
3	J	4	0	0	0	0
3	K	10	0	0	0	0
3	L	2	0	0	0	0
4	A	29	0	0	0	0
4	B	19	0	0	0	0
4	C	58	0	0	0	0
4	D	16	0	0	0	0
4	E	60	0	0	0	0
4	F	7	0	0	0	0
4	G	27	0	0	0	0
4	H	2	0	0	0	0
4	I	65	0	0	0	0
4	J	61	0	0	0	0
4	K	98	0	0	0	0
4	L	43	0	0	0	0
All	All	22833	0	22464	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 60 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:113:SER:HB2	2:B:114:PRO:HD3	1.65	0.78
2:H:117:LYS:HA	2:H:117:LYS:HE3	1.69	0.74
1:G:163:ASN:HA	2:H:117:LYS:NZ	2.03	0.74
2:D:81:ARG:HG2	2:D:83:GLU:HG2	1.72	0.69
1:G:198:GLU:O	1:G:202:THR:HG23	1.94	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/328 (98%)	320 (100%)	1 (0%)	0	100	100
1	C	321/328 (98%)	321 (100%)	0	0	100	100
1	E	323/328 (98%)	321 (99%)	2 (1%)	0	100	100
1	G	321/328 (98%)	320 (100%)	1 (0%)	0	100	100
1	I	321/328 (98%)	319 (99%)	2 (1%)	0	100	100
1	J	323/328 (98%)	321 (99%)	2 (1%)	0	100	100
1	K	322/328 (98%)	321 (100%)	1 (0%)	0	100	100
1	L	323/328 (98%)	322 (100%)	0	1 (0%)	41	50
2	B	103/109 (94%)	95 (92%)	5 (5%)	3 (3%)	4	3
2	D	105/109 (96%)	101 (96%)	4 (4%)	0	100	100
2	F	103/109 (94%)	99 (96%)	4 (4%)	0	100	100
2	H	102/109 (94%)	97 (95%)	5 (5%)	0	100	100
All	All	2988/3060 (98%)	2957 (99%)	27 (1%)	4 (0%)	51	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	113	SER
2	B	109	VAL
2	B	114	PRO
1	L	42	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/256 (99%)	248 (98%)	5 (2%)	55	72
1	C	253/256 (99%)	247 (98%)	6 (2%)	49	66
1	E	255/256 (100%)	248 (97%)	7 (3%)	44	61
1	G	253/256 (99%)	245 (97%)	8 (3%)	39	54
1	I	253/256 (99%)	251 (99%)	2 (1%)	81	91
1	J	255/256 (100%)	250 (98%)	5 (2%)	55	72
1	K	254/256 (99%)	251 (99%)	3 (1%)	71	84
1	L	255/256 (100%)	246 (96%)	9 (4%)	36	50
2	B	80/82 (98%)	79 (99%)	1 (1%)	69	82
2	D	81/82 (99%)	80 (99%)	1 (1%)	71	84
2	F	80/82 (98%)	78 (98%)	2 (2%)	47	65
2	H	79/82 (96%)	75 (95%)	4 (5%)	24	33
All	All	2351/2376 (99%)	2298 (98%)	53 (2%)	52	67

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	34	LYS
1	G	307	LEU
1	L	154	LEU
1	G	113	GLN
1	G	212	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such

sidechains are listed below:

Mol	Chain	Res	Type
1	E	236	GLN
1	J	29	GLN
1	L	25	GLN
1	E	29	GLN
1	K	129	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 49 ligands modelled in this entry, 49 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	323/328 (98%)	-0.09	1 (0%) 94 96	61, 83, 112, 137	0
1	C	323/328 (98%)	-0.24	2 (0%) 89 92	50, 65, 112, 134	0
1	E	323/328 (98%)	-0.23	1 (0%) 94 96	49, 67, 93, 121	0
1	G	322/328 (98%)	-0.08	4 (1%) 79 83	54, 77, 112, 138	0
1	I	323/328 (98%)	-0.24	3 (0%) 84 88	52, 64, 103, 128	0
1	J	324/328 (98%)	-0.24	1 (0%) 94 96	50, 64, 97, 125	0
1	K	323/328 (98%)	-0.32	0 100 100	45, 59, 90, 107	0
1	L	323/328 (98%)	-0.16	1 (0%) 94 96	52, 73, 102, 129	0
2	B	105/109 (96%)	0.04	3 (2%) 51 58	57, 81, 119, 129	0
2	D	107/109 (98%)	0.32	11 (10%) 6 9	60, 88, 116, 140	0
2	F	105/109 (96%)	0.17	2 (1%) 66 73	60, 88, 123, 136	0
2	H	104/109 (95%)	1.22	30 (28%) 0 0	75, 112, 146, 156	0
All	All	3005/3060 (98%)	-0.11	59 (1%) 65 71	45, 71, 113, 156	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	122	ARG	10.8
2	H	118	ARG	6.3
2	H	110	GLY	5.8
2	D	18	GLY	5.3
2	H	68	THR	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	A	405	1/1	0.35	0.12	119,119,119,119	0
3	CA	A	406	1/1	0.49	0.30	135,135,135,135	0
3	CA	I	401	1/1	0.79	0.27	112,112,112,112	0
3	CA	K	410	1/1	0.83	0.17	102,102,102,102	0
3	CA	J	404	1/1	0.85	0.09	127,127,127,127	0
3	CA	C	401	1/1	0.85	0.21	118,118,118,118	0
3	CA	J	403	1/1	0.87	0.19	81,81,81,81	0
3	CA	K	406	1/1	0.87	0.22	105,105,105,105	0
3	CA	L	402	1/1	0.87	0.25	115,115,115,115	0
3	CA	E	405	1/1	0.91	0.16	91,91,91,91	0
3	CA	G	406	1/1	0.92	0.11	156,156,156,156	0
3	CA	J	401	1/1	0.92	0.03	113,113,113,113	0
3	CA	A	403	1/1	0.93	0.17	72,72,72,72	0
3	CA	K	409	1/1	0.93	0.12	66,66,66,66	0
3	CA	G	409	1/1	0.93	0.18	81,81,81,81	0
3	CA	G	404	1/1	0.93	0.18	76,76,76,76	0
3	CA	G	403	1/1	0.94	0.11	71,71,71,71	0
3	CA	F	201	1/1	0.94	0.12	108,108,108,108	0
3	CA	E	406	1/1	0.94	0.20	101,101,101,101	0
3	CA	A	404	1/1	0.94	0.14	65,65,65,65	0
3	CA	A	408	1/1	0.94	0.12	77,77,77,77	0
3	CA	A	407	1/1	0.94	0.17	72,72,72,72	0
3	CA	A	402	1/1	0.95	0.13	77,77,77,77	0
3	CA	K	401	1/1	0.95	0.11	67,67,67,67	0
3	CA	I	402	1/1	0.95	0.21	86,86,86,86	0
3	CA	G	405	1/1	0.95	0.22	140,140,140,140	0
3	CA	K	402	1/1	0.96	0.19	66,66,66,66	0
3	CA	G	408	1/1	0.96	0.11	67,67,67,67	0
3	CA	E	403	1/1	0.96	0.19	74,74,74,74	0
3	CA	K	407	1/1	0.96	0.10	100,100,100,100	0
3	CA	K	403	1/1	0.96	0.19	74,74,74,74	0
3	CA	E	407	1/1	0.96	0.15	73,73,73,73	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	K	404	1/1	0.97	0.11	77,77,77,77	0
3	CA	A	401	1/1	0.97	0.14	71,71,71,71	0
3	CA	E	402	1/1	0.97	0.11	62,62,62,62	0
3	CA	H	201	1/1	0.97	0.14	112,112,112,112	0
3	CA	E	409	1/1	0.97	0.16	102,102,102,102	0
3	CA	G	407	1/1	0.98	0.14	72,72,72,72	0
3	CA	G	401	1/1	0.98	0.10	72,72,72,72	0
3	CA	E	401	1/1	0.98	0.14	66,66,66,66	0
3	CA	G	402	1/1	0.98	0.13	61,61,61,61	0
3	CA	J	402	1/1	0.98	0.23	111,111,111,111	0
3	CA	K	408	1/1	0.98	0.16	63,63,63,63	0
3	CA	L	401	1/1	0.98	0.17	107,107,107,107	0
3	CA	I	403	1/1	0.98	0.19	92,92,92,92	0
3	CA	K	405	1/1	0.99	0.15	77,77,77,77	0
3	CA	E	404	1/1	0.99	0.13	67,67,67,67	0
3	CA	I	404	1/1	0.99	0.14	60,60,60,60	0
3	CA	E	408	1/1	0.99	0.11	74,74,74,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.