



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 03:24 am BST

PDB ID : 3MGQ
Title : Binding of Nickel ions to the Nucleosome Core Particle
Authors : Mohideen, K.; Muhammad, R.; Davey, C.A.
Deposited on : 2010-04-07
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	<div> <div>3%</div> <div> <div></div> <div>62%</div> <div>11%</div> <div>27%</div> </div> </div>
1	E	135	<div> <div>4%</div> <div> <div></div> <div>60%</div> <div>12%</div> <div>27%</div> </div> </div>
2	B	102	<div> <div>3%</div> <div> <div></div> <div>64%</div> <div>11%</div> <div>23%</div> </div> </div>
2	F	102	<div> <div>0%</div> <div> <div></div> <div>73%</div> <div>11%</div> <div>15%</div> </div> </div>
3	C	119	<div> <div>3%</div> <div> <div></div> <div>69%</div> <div>17%</div> <div>12%</div> </div> </div>
3	G	119	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>17%</div> <div>11%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	125	
4	H	125	
5	I	147	
6	J	147	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	NI	C	120	-	-	-	X
7	NI	D	123	-	-	-	X
7	NI	F	103	-	-	-	X
7	NI	G	120	-	-	-	X
7	NI	H	125	-	-	-	X
7	NI	I	77	-	-	-	X
7	NI	I	79	-	-	-	X
7	NI	I	81	-	-	-	X
7	NI	I	82	-	-	-	X
7	NI	I	84	-	-	-	X
7	NI	I	85	-	-	-	X
7	NI	I	87	-	-	-	X
7	NI	I	88	-	-	-	X
7	NI	I	90	-	-	-	X
7	NI	I	91	-	-	-	X
7	NI	J	105	-	-	-	X
7	NI	J	80	-	-	-	X
7	NI	J	83	-	-	-	X
7	NI	J	85	-	-	-	X
7	NI	J	86	-	-	-	X
7	NI	J	88	-	-	-	X
7	NI	J	90	-	-	-	X
7	NI	J	91	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	0	0
			817	515	158	141	3			
1	E	99	Total	C	N	O	S	0	0	0
			817	515	158	141	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	SEE REMARK 999	UNP P84233
E	102	ALA	GLY	SEE REMARK 999	UNP P84233

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	79	Total	C	N	O	S	0	0	0
			627	395	121	110	1			
2	F	87	Total	C	N	O	S	0	0	0
			703	442	142	118	1			

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	105	Total	C	N	O		0	0	0
			813	513	159	141				
3	G	106	Total	C	N	O		0	0	0
			818	516	160	142				

- Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	100	Total	C	N	O	S	0	0	0
			789	495	147	145	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	99	Total	C	N	O	S	0	0	0
			785	493	146	144	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	SEE REMARK 999	UNP P02281
H	29	THR	SER	SEE REMARK 999	UNP P02281

- Molecule 5 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	147	Total	C	N	O	P	0	0	0
			3011	1440	546	879	146			

- Molecule 6 is a DNA chain called DNA (147-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	147	Total	C	N	O	P	0	0	0
			3010	1440	543	881	146			





- Molecule 7 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	G	1	Total	Ni	0	0
			1	1		
7	J	19	Total	Ni	0	0
			19	19		
7	D	2	Total	Ni	0	0
			2	2		
7	E	1	Total	Ni	0	0
			1	1		
7	H	3	Total	Ni	0	0
			3	3		
7	I	18	Total	Ni	0	0
			18	18		
7	C	1	Total	Ni	0	0
			1	1		
7	A	1	Total	Ni	0	0
			1	1		
7	F	1	Total	Ni	0	0
			1	1		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	1	Total 1	Cl 1	0	0
8	A	1	Total 1	Cl 1	0	0
8	C	1	Total 1	Cl 1	0	0
8	E	1	Total 1	Cl 1	0	0

- Molecule 1: Histone H3.2







- Molecule 3: Histone H2A




- Molecule 4: Histone H2B 1.1



- Molecule 4: Histone H2B 1.1



- Molecule 5: DNA (147-MER)



- Molecule 6: DNA (147-MER)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.13Å 109.64Å 183.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.00 – 2.65 50.97 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.0 (51.00-2.65) 98.0 (50.97-2.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.65Å)	Xtriage
Refinement program	REFMAC Rigid Body	Depositor
R, R_{free}	0.232 , 0.276 0.236 , 0.274	Depositor DCC
R_{free} test set	1248 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å ²)	68.9	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.020 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12241	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/829	0.65	0/1111
1	E	0.70	0/829	0.72	0/1111
2	B	0.58	0/634	0.68	0/848
2	F	0.71	0/711	0.82	0/948
3	C	0.68	0/823	0.68	0/1110
3	G	0.54	0/828	0.61	0/1117
4	D	0.62	0/800	0.67	0/1070
4	H	0.59	0/796	0.65	0/1065
5	I	0.76	0/3378	1.49	46/5212 (0.9%)
6	J	0.79	0/3376	1.51	43/5209 (0.8%)
All	All	0.71	0/13004	1.20	89/18801 (0.5%)

There are no bond length outliers.

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	-13	DA	O4'-C1'-N9	-9.36	101.45	108.00
5	I	-62	DC	O4'-C1'-N1	9.26	114.48	108.00
6	J	67	DT	O4'-C1'-N1	8.58	114.01	108.00
6	J	54	DA	O4'-C1'-N9	8.40	113.88	108.00
6	J	-14	DC	O4'-C1'-N1	8.12	113.68	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	817	0	858	9	0
1	E	817	0	858	12	0
2	B	627	0	663	8	0
2	F	703	0	755	10	0
3	C	813	0	872	13	0
3	G	818	0	877	20	0
4	D	789	0	828	6	0
4	H	785	0	825	15	0
5	I	3011	0	1662	40	0
6	J	3010	0	1663	53	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
7	D	2	0	0	0	0
7	E	1	0	0	0	0
7	F	1	0	0	0	0
7	G	1	0	0	0	0
7	H	3	0	0	0	0
7	I	18	0	0	0	0
7	J	19	0	0	0	0
8	A	1	0	0	1	0
8	C	1	0	0	0	0
8	E	1	0	0	1	0
8	G	1	0	0	0	0
All	All	12241	0	9861	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:50:TYR:OH	4:H:92:GLN:HG3	1.69	0.91
6:J:-16:DG:OP2	6:J:-15:DG:H5''	1.69	0.91
3:G:13:LYS:HB3	6:J:-43:DG:H4'	1.57	0.86
3:G:13:LYS:HA	6:J:-42:DT:H5''	1.61	0.80
4:D:121:ALA:O	4:D:122:LYS:HB2	1.86	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/135 (72%)	94 (97%)	3 (3%)	0	100	100
1	E	97/135 (72%)	95 (98%)	2 (2%)	0	100	100
2	B	77/102 (76%)	75 (97%)	2 (3%)	0	100	100
2	F	85/102 (83%)	80 (94%)	3 (4%)	2 (2%)	6	7
3	C	103/119 (87%)	100 (97%)	3 (3%)	0	100	100
3	G	104/119 (87%)	99 (95%)	4 (4%)	1 (1%)	15	23
4	D	98/125 (78%)	91 (93%)	4 (4%)	3 (3%)	4	5
4	H	97/125 (78%)	90 (93%)	5 (5%)	2 (2%)	7	10
All	All	758/962 (79%)	724 (96%)	26 (3%)	8 (1%)	14	21

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	26	ARG
4	H	101	GLY
4	D	101	GLY
2	F	18	HIS
3	G	14	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/110 (78%)	84 (98%)	2 (2%)	50	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	86/110 (78%)	82 (95%)	4 (5%)	26	40
2	B	64/78 (82%)	59 (92%)	5 (8%)	12	20
2	F	72/78 (92%)	69 (96%)	3 (4%)	30	45
3	C	84/92 (91%)	75 (89%)	9 (11%)	6	9
3	G	84/92 (91%)	82 (98%)	2 (2%)	49	67
4	D	85/105 (81%)	82 (96%)	3 (4%)	36	52
4	H	85/105 (81%)	78 (92%)	7 (8%)	11	17
All	All	646/770 (84%)	611 (95%)	35 (5%)	22	34

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	119	LYS
1	E	60	LEU
4	H	85	THR
4	D	25	LYS
4	D	30	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	125	GLN
4	H	92	GLN
3	G	110	ASN
1	E	68	GLN
3	G	31	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 51 ligands modelled in this entry, 51 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	99/135 (73%)	0.53	4 (4%)	38 34	41, 60, 93, 110	0
1	E	99/135 (73%)	0.45	6 (6%)	21 18	35, 47, 78, 101	0
2	B	79/102 (77%)	0.51	3 (3%)	40 36	45, 54, 72, 80	0
2	F	87/102 (85%)	0.42	1 (1%)	80 79	36, 47, 66, 102	0
3	C	105/119 (88%)	0.42	3 (2%)	51 48	37, 54, 73, 93	0
3	G	106/119 (89%)	0.39	2 (1%)	66 63	41, 62, 84, 101	0
4	D	100/125 (80%)	0.56	4 (4%)	38 34	41, 56, 108, 120	0
4	H	99/125 (79%)	0.49	4 (4%)	38 34	47, 63, 107, 124	0
5	I	147/147 (100%)	0.26	10 (6%)	17 14	62, 112, 143, 162	0
6	J	147/147 (100%)	0.21	6 (4%)	37 33	67, 113, 147, 154	0
All	All	1068/1256 (85%)	0.41	43 (4%)	38 34	35, 62, 135, 162	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	13	LYS	8.0
3	C	119	LYS	7.1
1	A	135	ALA	6.7
2	B	102	GLY	5.9
4	H	24	LYS	5.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	NI	J	91	1/1	-0.54	1.50	88,88,88,88	1
7	NI	J	86	1/1	-0.49	1.31	158,158,158,158	1
7	NI	I	84	1/1	-0.21	0.60	110,110,110,110	1
7	NI	F	103	1/1	-0.05	0.97	104,104,104,104	1
7	NI	I	87	1/1	0.03	0.41	165,165,165,165	1
7	NI	C	120	1/1	0.11	0.72	111,111,111,111	1
7	NI	J	88	1/1	0.14	0.73	97,97,97,97	1
7	NI	J	82	1/1	0.15	0.31	105,105,105,105	1
7	NI	I	78	1/1	0.19	0.18	142,142,142,142	1
7	NI	J	90	1/1	0.22	0.62	186,186,186,186	1
7	NI	J	80	1/1	0.23	0.52	116,116,116,116	1
7	NI	I	89	1/1	0.29	0.27	163,163,163,163	1
7	NI	J	85	1/1	0.32	0.46	232,232,232,232	1
7	NI	D	123	1/1	0.33	0.42	135,135,135,135	1
7	NI	H	125	1/1	0.41	0.51	100,100,100,100	1
7	NI	I	82	1/1	0.42	0.45	67,67,67,67	1
7	NI	I	85	1/1	0.46	1.28	142,142,142,142	1
7	NI	I	91	1/1	0.52	0.83	318,318,318,318	1
7	NI	J	105	1/1	0.57	0.43	345,345,345,345	1
7	NI	I	90	1/1	0.58	0.63	110,110,110,110	1
7	NI	I	77	1/1	0.60	0.41	111,111,111,111	1
7	NI	A	136	1/1	0.67	0.36	200,200,200,200	1
7	NI	J	74	1/1	0.70	0.35	97,97,97,97	1
7	NI	I	81	1/1	0.73	0.46	84,84,84,84	1
7	NI	J	81	1/1	0.74	0.35	137,137,137,137	1
7	NI	I	79	1/1	0.74	0.68	106,106,106,106	1
7	NI	H	124	1/1	0.74	0.36	106,106,106,106	1
7	NI	I	88	1/1	0.75	0.85	81,81,81,81	1
7	NI	J	89	1/1	0.76	0.28	140,140,140,140	1
7	NI	G	120	1/1	0.76	0.84	82,82,82,82	1
7	NI	J	83	1/1	0.76	0.54	84,84,84,84	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	NI	I	76	1/1	0.80	0.32	103,103,103,103	1
7	NI	J	84	1/1	0.81	0.41	132,132,132,132	1
7	NI	J	76	1/1	0.84	0.39	83,83,83,83	1
7	NI	I	83	1/1	0.84	0.24	77,77,77,77	1
7	NI	J	87	1/1	0.88	0.22	87,87,87,87	1
7	NI	I	74	1/1	0.88	0.48	106,106,106,106	1
7	NI	H	123	1/1	0.88	1.04	72,72,72,72	1
7	NI	J	75	1/1	0.89	0.28	100,100,100,100	1
7	NI	J	79	1/1	0.89	0.35	89,89,89,89	1
7	NI	J	78	1/1	0.89	0.40	66,66,66,66	1
7	NI	I	80	1/1	0.91	0.23	98,98,98,98	1
8	CL	C	3146	1/1	0.92	0.19	72,72,72,72	0
7	NI	E	136	1/1	0.92	0.47	88,88,88,88	0
7	NI	J	77	1/1	0.92	0.40	88,88,88,88	1
8	CL	A	3147	1/1	0.93	0.11	67,67,67,67	1
8	CL	E	3148	1/1	0.94	0.20	65,65,65,65	1
7	NI	I	75	1/1	0.94	0.47	75,75,75,75	1
8	CL	G	3145	1/1	0.95	0.17	78,78,78,78	0
7	NI	D	124	1/1	0.96	0.83	83,83,83,83	1
7	NI	I	86	1/1	0.97	0.27	71,71,71,71	1

6.5 Other polymers

There are no such residues in this entry.