



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 09:04 AM BST

PDB ID : 5MGT
Title : Complex of human NKR-P1 and LLT1 in deglycosylated forms
Authors : Blaha, J.; Skalova, T.; Stransky, J.; Koval, T.; Hasek, J.; Yuguang, Z.; Harlos, K.; Vanek, O.; Dohnalek, J.
Deposited on : 2016-11-22
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	<div> <div>81%</div> <div>14%</div> </div>
1	B	135	<div> <div>80%</div> <div>14%</div> </div>
2	C	146	<div> <div>79%</div> <div>8%</div> <div>13%</div> </div>
2	D	146	<div> <div>81%</div> <div>5%</div> <div>14%</div> </div>
2	E	146	<div> <div>80%</div> <div>6%</div> <div>14%</div> </div>
2	F	146	<div> <div>78%</div> <div>6%</div> <div>14%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7030 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-type lectin domain family 2 member D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	1	0
			956	604	166	180	6			
1	B	116	Total	C	N	O	S	0	2	0
			962	609	165	182	6			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	ILE	-	expression tag	UNP Q9UHP7
A	70	THR	-	expression tag	UNP Q9UHP7
A	71	GLY	-	expression tag	UNP Q9UHP7
A	176	CYS	HIS	engineered mutation	UNP Q9UHP7
A	192	GLY	-	expression tag	UNP Q9UHP7
A	193	THR	-	expression tag	UNP Q9UHP7
A	194	LYS	-	expression tag	UNP Q9UHP7
A	195	HIS	-	expression tag	UNP Q9UHP7
A	196	HIS	-	expression tag	UNP Q9UHP7
A	197	HIS	-	expression tag	UNP Q9UHP7
A	198	HIS	-	expression tag	UNP Q9UHP7
A	199	HIS	-	expression tag	UNP Q9UHP7
A	200	HIS	-	expression tag	UNP Q9UHP7
A	201	HIS	-	expression tag	UNP Q9UHP7
A	202	HIS	-	expression tag	UNP Q9UHP7
A	203	GLY	-	expression tag	UNP Q9UHP7
B	69	ILE	-	expression tag	UNP Q9UHP7
B	70	THR	-	expression tag	UNP Q9UHP7
B	71	GLY	-	expression tag	UNP Q9UHP7
B	176	CYS	HIS	engineered mutation	UNP Q9UHP7
B	192	GLY	-	expression tag	UNP Q9UHP7
B	193	THR	-	expression tag	UNP Q9UHP7
B	194	LYS	-	expression tag	UNP Q9UHP7
B	195	HIS	-	expression tag	UNP Q9UHP7
B	196	HIS	-	expression tag	UNP Q9UHP7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	197	HIS	-	expression tag	UNP Q9UHP7
B	198	HIS	-	expression tag	UNP Q9UHP7
B	199	HIS	-	expression tag	UNP Q9UHP7
B	200	HIS	-	expression tag	UNP Q9UHP7
B	201	HIS	-	expression tag	UNP Q9UHP7
B	202	HIS	-	expression tag	UNP Q9UHP7
B	203	GLY	-	expression tag	UNP Q9UHP7

- Molecule 2 is a protein called Killer cell lectin-like receptor subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	127	Total	C	N	O	S	0	6	0
			1072	678	182	206	6			
2	D	125	Total	C	N	O	S	0	3	0
			1044	663	176	199	6			
2	E	126	Total	C	N	O	S	0	4	0
			1052	667	178	201	6			
2	F	125	Total	C	N	O	S	0	4	0
			1049	665	178	200	6			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	87	GLU	-	expression tag	UNP Q12918
C	88	THR	-	expression tag	UNP Q12918
C	89	GLY	-	expression tag	UNP Q12918
C	226	LYS	-	expression tag	UNP Q12918
C	227	HIS	-	expression tag	UNP Q12918
C	228	HIS	-	expression tag	UNP Q12918
C	229	HIS	-	expression tag	UNP Q12918
C	230	HIS	-	expression tag	UNP Q12918
C	231	HIS	-	expression tag	UNP Q12918
C	232	HIS	-	expression tag	UNP Q12918
D	87	GLU	-	expression tag	UNP Q12918
D	88	THR	-	expression tag	UNP Q12918
D	89	GLY	-	expression tag	UNP Q12918
D	226	LYS	-	expression tag	UNP Q12918
D	227	HIS	-	expression tag	UNP Q12918
D	228	HIS	-	expression tag	UNP Q12918
D	229	HIS	-	expression tag	UNP Q12918
D	230	HIS	-	expression tag	UNP Q12918
D	231	HIS	-	expression tag	UNP Q12918

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	232	HIS	-	expression tag	UNP Q12918
E	87	GLU	-	expression tag	UNP Q12918
E	88	THR	-	expression tag	UNP Q12918
E	89	GLY	-	expression tag	UNP Q12918
E	226	LYS	-	expression tag	UNP Q12918
E	227	HIS	-	expression tag	UNP Q12918
E	228	HIS	-	expression tag	UNP Q12918
E	229	HIS	-	expression tag	UNP Q12918
E	230	HIS	-	expression tag	UNP Q12918
E	231	HIS	-	expression tag	UNP Q12918
E	232	HIS	-	expression tag	UNP Q12918
F	87	GLU	-	expression tag	UNP Q12918
F	88	THR	-	expression tag	UNP Q12918
F	89	GLY	-	expression tag	UNP Q12918
F	226	LYS	-	expression tag	UNP Q12918
F	227	HIS	-	expression tag	UNP Q12918
F	228	HIS	-	expression tag	UNP Q12918
F	229	HIS	-	expression tag	UNP Q12918
F	230	HIS	-	expression tag	UNP Q12918
F	231	HIS	-	expression tag	UNP Q12918
F	232	HIS	-	expression tag	UNP Q12918

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		
3	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	F	1	Total O S 5 4 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Cl 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	87	Total O 87 87	0	0
6	B	124	Total O 124 124	0	0
6	C	137	Total O 137 137	0	0
6	D	118	Total O 118 118	0	0
6	E	107	Total O 107 107	0	0
6	F	118	Total O 118 118	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C-type lectin domain family 2 member D





- Molecule 1: C-type lectin domain family 2 member D




- Molecule 2: Killer cell lectin-like receptor subfamily B member 1




- Molecule 2: Killer cell lectin-like receptor subfamily B member 1



- Molecule 2: Killer cell lectin-like receptor subfamily B member 1



- Molecule 2: Killer cell lectin-like receptor subfamily B member 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.58 Å 80.15 Å 272.95 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.91 – 1.90 76.90 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (76.91-1.90) 100.0 (76.90-1.90)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 1.90 Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.162 , 0.201 0.173 , 0.185	Depositor DCC
R_{free} test set	3915 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7030	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/988	0.92	1/1336 (0.1%)
1	B	1.03	1/997 (0.1%)	0.93	2/1348 (0.1%)
2	C	0.96	0/1106	0.90	2/1498 (0.1%)
2	D	0.97	1/1075 (0.1%)	0.87	2/1457 (0.1%)
2	E	0.90	0/1086	0.88	4/1472 (0.3%)
2	F	0.96	0/1080	0.94	4/1464 (0.3%)
All	All	0.96	2/6332 (0.0%)	0.91	15/8575 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	200	GLU	CD-OE2	5.34	1.31	1.25
1	B	125	TYR	CE1-CZ	-5.03	1.32	1.38

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	134	ASP	CB-CG-OD1	6.76	124.38	118.30
2	F	177	ASP	CB-CG-OD2	-6.64	112.32	118.30
2	E	134	ASP	CB-CG-OD2	-6.18	112.74	118.30
2	D	181	ARG	NE-CZ-NH2	6.17	123.39	120.30
2	C	207	ARG	NE-CZ-NH1	-6.06	117.27	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	956	0	877	3	0
1	B	962	0	886	4	0
2	C	1072	0	1044	6	1
2	D	1044	0	1021	8	1
2	E	1052	0	1029	8	0
2	F	1049	0	1024	7	0
3	A	28	0	26	0	0
3	B	28	0	26	0	0
3	C	28	0	26	6	0
3	D	28	0	26	2	0
3	E	14	0	13	0	0
3	F	42	0	39	5	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
4	F	5	0	0	0	0
5	D	1	0	0	0	0
6	A	87	0	0	1	0
6	B	124	0	0	0	0
6	C	137	0	0	1	0
6	D	118	0	0	1	0
6	E	107	0	0	1	0
6	F	118	0	0	4	0
All	All	7030	0	6037	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:177[A]:ASP:OD2	6:E:601:HOH:O	1.78	1.02
2:E:116[B]:ASN:ND2	3:F:502:NAG:O3	1.99	0.95
2:F:143[B]:ASN:ND2	6:F:602:HOH:O	2.00	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:116[A]:ASN:ND2	3:D:502:NAG:O4	2.16	0.78
2:F:143[B]:ASN:OD1	6:F:601:HOH:O	1.99	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:103:GLU:OE2	2:D:162[B]:GLU:OE2[1_455]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/135 (85%)	110 (96%)	5 (4%)	0	100	100
1	B	116/135 (86%)	112 (97%)	4 (3%)	0	100	100
2	C	131/146 (90%)	128 (98%)	3 (2%)	0	100	100
2	D	126/146 (86%)	124 (98%)	2 (2%)	0	100	100
2	E	128/146 (88%)	126 (98%)	2 (2%)	0	100	100
2	F	127/146 (87%)	124 (98%)	3 (2%)	0	100	100
All	All	743/854 (87%)	724 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/116 (87%)	99 (98%)	2 (2%)	55	51
1	B	103/116 (89%)	101 (98%)	2 (2%)	57	53
2	C	125/138 (91%)	124 (99%)	1 (1%)	81	82
2	D	122/138 (88%)	121 (99%)	1 (1%)	81	82
2	E	123/138 (89%)	121 (98%)	2 (2%)	62	60
2	F	123/138 (89%)	121 (98%)	2 (2%)	62	60
All	All	697/784 (89%)	687 (99%)	10 (1%)	65	65

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	160	LEU
2	D	113	ASN
2	E	174	ASN
1	B	169	LYS
2	E	160	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	167	ASN
2	D	113	ASN
2	E	174	ASN
2	E	176	ASN
2	F	99	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 1 is monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	C	503	-	4,4,4	0.38	0	6,6,6	0.70	0
3	NAG	C	502	2	14,14,15	0.55	0	17,19,21	1.51	3 (17%)
4	SO4	D	503	-	4,4,4	0.33	0	6,6,6	1.28	0
3	NAG	A	502	1	14,14,15	0.66	0	17,19,21	1.50	3 (17%)
3	NAG	F	503	2	14,14,15	0.67	0	17,19,21	1.97	5 (29%)
3	NAG	B	501	1	14,14,15	0.55	0	17,19,21	1.78	5 (29%)
4	SO4	A	503	-	4,4,4	0.36	0	6,6,6	0.27	0
4	SO4	F	504	-	4,4,4	0.47	0	6,6,6	0.44	0
3	NAG	F	502	2	14,14,15	0.92	0	17,19,21	1.56	3 (17%)
3	NAG	B	502	1	14,14,15	0.88	0	17,19,21	1.05	1 (5%)
3	NAG	D	501	2	14,14,15	0.97	1 (7%)	17,19,21	1.20	2 (11%)
3	NAG	C	501	2	14,14,15	0.93	1 (7%)	17,19,21	1.22	1 (5%)
3	NAG	F	501	2	14,14,15	0.89	1 (7%)	17,19,21	1.23	2 (11%)
3	NAG	E	501	2	14,14,15	1.10	1 (7%)	17,19,21	0.94	1 (5%)
4	SO4	B	503	-	4,4,4	0.27	0	6,6,6	0.33	0
3	NAG	D	502	2	14,14,15	0.70	0	17,19,21	1.28	2 (11%)
4	SO4	B	504	-	4,4,4	0.45	0	6,6,6	0.20	0
3	NAG	A	501	1	14,14,15	0.38	0	17,19,21	1.36	3 (17%)
4	SO4	A	504	-	4,4,4	0.33	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	502	2	-	0/6/23/26	0/1/1/1
3	NAG	A	502	1	-	1/6/23/26	0/1/1/1
3	NAG	F	503	2	-	2/6/23/26	0/1/1/1
3	NAG	F	502	2	-	2/6/23/26	0/1/1/1
3	NAG	B	502	1	-	0/6/23/26	0/1/1/1
3	NAG	D	501	2	-	0/6/23/26	0/1/1/1
3	NAG	C	501	2	-	0/6/23/26	0/1/1/1
3	NAG	F	501	2	-	0/6/23/26	0/1/1/1
3	NAG	E	501	2	-	0/6/23/26	0/1/1/1
3	NAG	D	502	2	-	0/6/23/26	0/1/1/1
3	NAG	B	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	501	1	-	2/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	501	NAG	C1-C2	3.00	1.56	1.52
3	C	501	NAG	C1-C2	2.03	1.55	1.52
3	F	501	NAG	C4-C5	2.02	1.57	1.53
3	D	501	NAG	O5-C1	-2.01	1.40	1.43

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	503	NAG	C2-N2-C7	4.00	128.60	122.90
3	B	501	NAG	O5-C1-C2	-3.69	105.46	111.29
3	F	502	NAG	C4-C3-C2	-3.44	105.97	111.02
3	C	502	NAG	C1-C2-N2	3.36	116.22	110.49
3	F	503	NAG	O3-C3-C2	3.33	116.36	109.47

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	502	NAG	C8-C7-N2-C2
3	F	502	NAG	O7-C7-N2-C2
3	A	501	NAG	O5-C5-C6-O6
3	F	503	NAG	C4-C5-C6-O6
3	A	501	NAG	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	502	NAG	6	0
3	F	502	NAG	5	0
3	D	502	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/135 (85%)	-0.18	2 (1%) 70 72	15, 24, 49, 80	1 (0%)
1	B	116/135 (85%)	-0.28	1 (0%) 84 85	13, 20, 44, 67	0
2	C	127/146 (86%)	-0.24	1 (0%) 86 87	14, 20, 38, 66	0
2	D	125/146 (85%)	-0.17	0 100 100	13, 19, 39, 54	1 (0%)
2	E	126/146 (86%)	-0.19	3 (2%) 59 62	14, 22, 50, 65	1 (0%)
2	F	125/146 (85%)	-0.01	6 (4%) 30 33	15, 23, 55, 77	0
All	All	735/854 (86%)	-0.18	13 (1%) 68 71	13, 21, 47, 80	3 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	92	LEU	6.4
1	A	139	GLN	5.3
1	B	189	ILE	3.4
2	F	160	LEU	3.3
1	A	140	GLY	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	502	14/15	0.61	0.37	19,29,33,34	14
3	NAG	D	502	14/15	0.72	0.31	21,27,29,29	14
3	NAG	F	503	14/15	0.79	0.22	23,37,46,47	14
3	NAG	F	502	14/15	0.80	0.20	17,25,29,34	14
3	NAG	A	502	14/15	0.88	0.20	56,62,72,74	0
4	SO4	D	503	5/5	0.88	0.28	33,34,40,41	5
3	NAG	A	501	14/15	0.89	0.18	39,46,51,55	0
3	NAG	B	501	14/15	0.91	0.10	30,41,49,54	0
4	SO4	B	504	5/5	0.92	0.17	62,66,77,81	0
3	NAG	E	501	14/15	0.93	0.09	22,26,32,34	0
4	SO4	F	504	5/5	0.93	0.28	60,63,69,71	0
4	SO4	A	503	5/5	0.94	0.23	64,67,69,69	0
4	SO4	A	504	5/5	0.94	0.24	67,72,75,81	0
3	NAG	C	501	14/15	0.95	0.10	21,24,32,33	0
4	SO4	C	503	5/5	0.95	0.18	62,68,72,73	0
3	NAG	D	501	14/15	0.96	0.08	21,28,42,42	0
4	SO4	B	503	5/5	0.96	0.16	56,58,63,68	0
3	NAG	F	501	14/15	0.96	0.10	23,29,41,45	0
3	NAG	B	502	14/15	0.97	0.07	19,21,24,25	0
5	CL	D	504	1/1	0.99	0.15	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.