



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 1, 2020 – 06:03 PM EST

PDB ID : 6MI1
Title : CRYSTAL STRUCTURE ANALYSIS OF THE VARIANT PLANT EXOHYDROLASE ARG158ALA-GLU161ALA IN COMPLEX WITH METHYL 6-T HIO-BETA-GENTIOBIOSIDE
Authors : Streltsov, V.A.; Luang, S.; Hrmova, M.
Deposited on : 2018-09-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>9%</div> </div> </div>
2	B	2	<div> <div>50%</div> <div>50%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	708	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-D-glucan exohydrolase isoenzyme ExoI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	606	Total	C	N	O	S	0	3	0
			4598	2908	799	865	26			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	HIS	-	expression tag	UNP Q9XEI3
A	-2	HIS	-	expression tag	UNP Q9XEI3
A	-1	ALA	-	expression tag	UNP Q9XEI3
A	158	ALA	ARG	engineered mutation	UNP Q9XEI3
A	161	ALA	GLU	engineered mutation	UNP Q9XEI3
A	320	LYS	ASN	conflict	UNP Q9XEI3

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-6)-methyl 6-thio-beta-D-glucopyranoside.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	S	0	1	0
			25	14	10	1			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	330	Total	O	0	0
			330	330		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-D-glucan exohydrolase isoenzyme ExoI



- Molecule 2: beta-D-glucopyranose-(1-6)-methyl 6-thio-beta-D-glucopyranoside



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.72Å 101.72Å 183.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.13 – 2.30 39.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.13-2.30) 99.4 (39.10-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.36 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0232	Depositor
R, R_{free}	0.168 , 0.214 0.175 , 0.218	Depositor DCC
R_{free} test set	2175 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5070	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U1Y, GOL, BGC, NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	3/4715 (0.1%)	1.02	9/6406 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	220	GLU	CD-OE1	6.52	1.32	1.25
1	A	470	GLU	CD-OE1	5.33	1.31	1.25
1	A	549	GLU	CD-OE2	5.03	1.31	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	ARG	NE-CZ-NH2	-8.56	116.02	120.30
1	A	592	ARG	NE-CZ-NH2	6.92	123.76	120.30
1	A	62	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	A	126	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	A	566	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	A	62	ARG	CG-CD-NE	-6.16	98.88	111.80
1	A	399	THR	N-CA-CB	-5.40	100.04	110.30
1	A	440[A]	ARG	NE-CZ-NH2	5.21	122.90	120.30
1	A	440[B]	ARG	NE-CZ-NH2	5.21	122.90	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-3	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4598	0	4569	47	0
2	B	25	0	10	2	0
3	A	42	0	39	1	0
4	A	60	0	80	7	0
5	A	15	0	0	0	0
6	A	330	0	0	7	0
All	All	5070	0	4698	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440[B]:ARG:NH2	6:A:802:HOH:O	1.71	1.22
1:A:394:ASN:HD21	1:A:404:LEU:H	1.16	0.91
1:A:227[A]:ARG:NH2	6:A:801:HOH:O	1.62	0.84
1:A:100:GLN:HA	1:A:100:GLN:HE21	1.42	0.83
1:A:466[A]:ASN:ND2	1:A:508:THR:OG1	2.16	0.79
1:A:156:TRP:HE1	1:A:219:ASN:HD22	1.28	0.78
1:A:377:HIS:HE1	4:A:708:GOL:H12	1.48	0.78
1:A:97:VAL:H	1:A:101:ASN:HD21	1.40	0.69
1:A:377:HIS:HE1	4:A:708:GOL:C1	2.04	0.69
1:A:227[A]:ARG:NH1	6:A:801:HOH:O	2.08	0.65
1:A:399:THR:HG22	6:A:826:HOH:O	1.96	0.65
1:A:57:GLY:H	2:B:1[B]:U1Y:C7	2.09	0.64
1:A:181:GLN:HE21	1:A:203:ALA:H	1.42	0.64
1:A:377:HIS:CE1	4:A:708:GOL:H12	2.32	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:ASN:HB2	3:A:702:NAG:H82	1.81	0.61
1:A:134:GLU:OE2	1:A:377:HIS:HD2	1.84	0.60
1:A:301:SER:O	1:A:331:HIS:HE1	1.85	0.60
1:A:97:VAL:H	1:A:101:ASN:ND2	2.00	0.59
1:A:167:ARG:HH21	1:A:171:GLN:HE22	1.50	0.58
1:A:100:GLN:HE21	1:A:100:GLN:CA	2.14	0.58
1:A:262:HIS:HE1	1:A:285:ASP:H	1.50	0.57
1:A:124:VAL:HG21	1:A:173:MET:SD	2.46	0.56
1:A:399:THR:CG2	6:A:826:HOH:O	2.52	0.54
1:A:100:GLN:HA	1:A:100:GLN:NE2	2.19	0.53
1:A:377:HIS:CE1	4:A:708:GOL:C1	2.88	0.51
1:A:167:ARG:NH2	1:A:171:GLN:HE22	2.08	0.51
1:A:524:SER:O	1:A:544:TRP:HA	2.11	0.50
1:A:493:LYS:HE3	1:A:493:LYS:HA	1.93	0.49
1:A:262:HIS:CE1	1:A:285:ASP:H	2.31	0.48
1:A:156:TRP:HE1	1:A:219:ASN:ND2	2.03	0.48
1:A:181:GLN:NE2	1:A:203:ALA:H	2.10	0.47
1:A:393:LYS:HE2	1:A:396:LYS:HA	1.97	0.46
1:A:456:ASP:CG	1:A:457:PRO:HD2	2.37	0.45
1:A:134:GLU:OE2	1:A:377:HIS:CD2	2.67	0.45
1:A:167:ARG:HH21	1:A:171:GLN:NE2	2.14	0.44
1:A:234:HIS:HE1	6:A:1067:HOH:O	2.01	0.44
1:A:301:SER:O	1:A:331:HIS:CE1	2.69	0.44
1:A:416:ALA:O	1:A:482:VAL:HA	2.19	0.43
1:A:156:TRP:CE2	1:A:158:ALA:HB3	2.54	0.42
1:A:602:THR:C	6:A:965:HOH:O	2.57	0.42
1:A:-2:HIS:O	1:A:-1:ALA:O	2.37	0.42
1:A:527:PRO:HB3	1:A:571:TRP:CD2	2.55	0.42
1:A:208:PHE:HA	1:A:209:VAL:HA	1.92	0.41
1:A:566:ARG:HH21	4:A:705:GOL:H2	1.86	0.40
1:A:465:GLU:HG2	1:A:466[B]:ASN:ND2	2.37	0.40
1:A:566:ARG:NH2	4:A:705:GOL:H2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	607/609 (100%)	581 (96%)	23 (4%)	3 (0%)	29	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	-2	HIS
1	A	-1	ALA
1	A	505	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/488 (100%)	471 (96%)	17 (4%)	36	50

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLU
1	A	54	LEU
1	A	63	LYS
1	A	90	MET
1	A	98	HIS
1	A	100	GLN
1	A	186	LYS
1	A	187	ASP
1	A	290	ASP
1	A	399	THR
1	A	440[A]	ARG
1	A	440[B]	ARG
1	A	493	LYS
1	A	517	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	586	HIS
1	A	592	ARG
1	A	602	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	100	GLN
1	A	101	ASN
1	A	102	ASN
1	A	112	ASN
1	A	171	GLN
1	A	181	GLN
1	A	199	ASN
1	A	219	ASN
1	A	234	HIS
1	A	262	HIS
1	A	331	HIS
1	A	377	HIS
1	A	394	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	U1Y	B	1[A]	-	13,13,13	1.37	2 (15%)	18,18,18	4.34	14 (77%)
2	U1Y	B	1[B]	-	13,13,13	1.42	2 (15%)	18,18,18	4.28	14 (77%)
2	BGC	B	2	2	11,11,12	0.95	0	15,15,17	1.23	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U1Y	B	1[A]	-	-	4/4/24/24	0/1/1/1
2	U1Y	B	1[B]	-	-	4/4/24/24	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1[B]	U1Y	C6-S6	-3.03	1.75	1.81
2	B	1[A]	U1Y	C6-S6	-3.03	1.75	1.81
2	B	1[B]	U1Y	O5-C5	-2.30	1.38	1.44
2	B	1[A]	U1Y	O5-C5	-2.30	1.38	1.44

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1[B]	U1Y	O1-C1-C2	8.61	118.24	108.15
2	B	1[A]	U1Y	O1-C1-C2	8.61	118.24	108.15
2	B	1[B]	U1Y	O5-C1-O1	-8.40	91.50	110.97
2	B	1[A]	U1Y	O5-C1-O1	-8.40	91.50	110.97
2	B	1[B]	U1Y	O5-C5-C6	-6.47	95.58	106.48
2	B	1[A]	U1Y	O5-C5-C6	-6.47	95.58	106.48
2	B	1[B]	U1Y	C5-C6-S6	-6.00	97.94	114.37
2	B	1[A]	U1Y	C5-C6-S6	-6.00	97.94	114.37
2	B	1[B]	U1Y	O4-C4-C3	-5.62	97.36	110.35
2	B	1[A]	U1Y	O4-C4-C3	-5.62	97.36	110.35
2	B	1[A]	U1Y	C7-O1-C1	-5.59	104.64	113.27
2	B	1[B]	U1Y	C7-O1-C1	-4.57	106.22	113.27
2	B	2	BGC	O5-C1-C2	2.64	114.85	110.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1[B]	U1Y	O3-C3-C4	-2.63	104.28	110.35
2	B	1[A]	U1Y	O3-C3-C4	-2.63	104.28	110.35
2	B	1[B]	U1Y	C3-C4-C5	2.48	114.66	110.24
2	B	1[A]	U1Y	C3-C4-C5	2.48	114.66	110.24
2	B	1[B]	U1Y	O3-C3-C2	2.45	116.01	110.35
2	B	1[A]	U1Y	O3-C3-C2	2.45	116.01	110.35
2	B	1[B]	U1Y	O4-C4-C5	2.38	115.21	109.30
2	B	1[A]	U1Y	O4-C4-C5	2.38	115.21	109.30
2	B	1[B]	U1Y	C1-C2-C3	-2.35	105.10	110.00
2	B	1[A]	U1Y	C1-C2-C3	-2.35	105.10	110.00
2	B	1[B]	U1Y	O2-C2-C3	-2.23	105.19	110.35
2	B	1[A]	U1Y	O2-C2-C3	-2.23	105.19	110.35
2	B	1[B]	U1Y	C1-O5-C5	2.20	118.02	113.69
2	B	1[A]	U1Y	C1-O5-C5	2.20	118.02	113.69
2	B	1[B]	U1Y	C6-C5-C4	2.19	118.32	112.84
2	B	1[A]	U1Y	C6-C5-C4	2.19	118.32	112.84
2	B	2	BGC	O2-C2-C3	2.04	114.23	110.14

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1[B]	U1Y	O5-C1-O1-C7
2	B	1[A]	U1Y	O5-C1-O1-C7
2	B	1[A]	U1Y	C2-C1-O1-C7
2	B	1[B]	U1Y	C2-C1-O1-C7
2	B	1[B]	U1Y	O5-C5-C6-S6
2	B	1[A]	U1Y	O5-C5-C6-S6
2	B	1[B]	U1Y	C4-C5-C6-S6
2	B	1[A]	U1Y	C4-C5-C6-S6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1[B]	U1Y	1	0
2	B	1[A]	U1Y	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	711	-	5,5,5	0.17	0	5,5,5	0.35	0
4	GOL	A	709	-	5,5,5	0.11	0	5,5,5	0.44	0
4	GOL	A	706	-	5,5,5	0.20	0	5,5,5	0.47	0
3	NAG	A	703	1	14,14,15	0.67	0	17,19,21	1.32	3 (17%)
4	GOL	A	705	-	5,5,5	0.17	0	5,5,5	0.42	0
5	SO4	A	716	-	4,4,4	0.31	0	6,6,6	0.14	0
4	GOL	A	710	-	5,5,5	0.29	0	5,5,5	0.54	0
4	GOL	A	713	-	5,5,5	0.13	0	5,5,5	0.31	0
4	GOL	A	707	-	5,5,5	0.18	0	5,5,5	0.41	0
5	SO4	A	715	-	4,4,4	0.25	0	6,6,6	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	708	-	5,5,5	0.10	0	5,5,5	0.34	0
4	GOL	A	712	-	5,5,5	0.14	0	5,5,5	0.37	0
3	NAG	A	702	1	14,14,15	0.75	1 (7%)	17,19,21	1.40	2 (11%)
4	GOL	A	714	-	5,5,5	0.17	0	5,5,5	0.49	0
3	NAG	A	701	1	14,14,15	0.67	0	17,19,21	1.63	3 (17%)
5	SO4	A	717	-	4,4,4	0.35	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	711	-	-	2/4/4/4	-
4	GOL	A	709	-	-	2/4/4/4	-
4	GOL	A	706	-	-	1/4/4/4	-
3	NAG	A	703	1	-	0/6/23/26	0/1/1/1
4	GOL	A	710	-	-	3/4/4/4	-
4	GOL	A	713	-	-	4/4/4/4	-
4	GOL	A	707	-	-	0/4/4/4	-
4	GOL	A	708	-	-	2/4/4/4	-
4	GOL	A	712	-	-	4/4/4/4	-
3	NAG	A	702	1	-	2/6/23/26	0/1/1/1
4	GOL	A	714	-	-	2/4/4/4	-
4	GOL	A	705	-	-	2/4/4/4	-
3	NAG	A	701	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	NAG	C1-C2	2.14	1.55	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAG	C1-O5-C5	3.76	117.28	112.19
3	A	701	NAG	O5-C1-C2	3.66	117.07	111.29
3	A	702	NAG	O5-C5-C6	2.89	111.73	107.20
3	A	701	NAG	O5-C5-C6	2.74	111.50	107.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	703	NAG	C2-N2-C7	2.46	126.41	122.90
3	A	703	NAG	O5-C5-C4	-2.33	105.15	110.83
3	A	703	NAG	O7-C7-N2	2.20	125.99	121.95
3	A	702	NAG	C1-O5-C5	2.07	114.99	112.19

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	709	GOL	O1-C1-C2-C3
4	A	710	GOL	O1-C1-C2-C3
4	A	713	GOL	C1-C2-C3-O3
4	A	714	GOL	O1-C1-C2-O2
4	A	714	GOL	O1-C1-C2-C3
4	A	705	GOL	O1-C1-C2-C3
4	A	712	GOL	O2-C2-C3-O3
3	A	702	NAG	O5-C5-C6-O6
4	A	711	GOL	O1-C1-C2-C3
4	A	708	GOL	O1-C1-C2-C3
4	A	712	GOL	O1-C1-C2-C3
4	A	712	GOL	C1-C2-C3-O3
4	A	709	GOL	O1-C1-C2-O2
4	A	710	GOL	O1-C1-C2-O2
4	A	713	GOL	O2-C2-C3-O3
4	A	712	GOL	O1-C1-C2-O2
4	A	705	GOL	O1-C1-C2-O2
3	A	702	NAG	C4-C5-C6-O6
3	A	701	NAG	C4-C5-C6-O6
3	A	701	NAG	O5-C5-C6-O6
4	A	711	GOL	O1-C1-C2-O2
4	A	713	GOL	O1-C1-C2-C3
4	A	706	GOL	C1-C2-C3-O3
4	A	713	GOL	O1-C1-C2-O2
4	A	710	GOL	C1-C2-C3-O3
4	A	708	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	709	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	705	GOL	2	0
4	A	708	GOL	4	0
3	A	702	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	606/609 (99%)	-0.12	37 (6%) 21 27	27, 37, 63, 130	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-2	HIS	7.8
1	A	586	HIS	5.0
1	A	-3	HIS	4.6
1	A	-1	ALA	3.8
1	A	224	ILE	3.7
1	A	187	ASP	3.6
1	A	585	ALA	3.5
1	A	249	VAL	3.4
1	A	97	VAL	3.2
1	A	294	THR	3.0
1	A	295	PRO	3.0
1	A	186	LYS	3.0
1	A	257	ASN	2.9
1	A	228	GLU	2.9
1	A	226	ASN	2.8
1	A	204	CYS	2.8
1	A	143	ALA	2.8
1	A	206	LYS	2.8
1	A	147	CYS	2.7
1	A	148	ILE	2.7
1	A	258	GLY	2.7
1	A	283	ILE	2.6
1	A	582	VAL	2.5
1	A	250	MET	2.5
1	A	256	TRP	2.4
1	A	161	ALA	2.4
1	A	144	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	ALA	2.3
1	A	203	ALA	2.3
1	A	205	ALA	2.3
1	A	146	PRO	2.3
1	A	2	TYR	2.3
1	A	96	ALA	2.2
1	A	215	VAL	2.2
1	A	296	ALA	2.1
1	A	225	ILE	2.0
1	A	298	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	U1Y	B	1[B]	13/13	0.89	0.14	34,55,61,74	1
2	U1Y	B	1[A]	13/13	0.89	0.14	46,55,61,74	1
2	BGC	B	2	11/12	0.96	0.15	34,44,49,53	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	701	14/15	0.74	0.37	67,90,103,104	0
4	GOL	A	705	6/6	0.76	0.20	68,77,81,84	0
4	GOL	A	714	6/6	0.77	0.29	75,94,97,102	0
4	GOL	A	712	6/6	0.79	0.24	52,69,88,91	0
4	GOL	A	708	6/6	0.80	0.22	61,74,76,81	0
4	GOL	A	707	6/6	0.82	0.16	69,74,83,84	0
3	NAG	A	702	14/15	0.83	0.43	80,95,104,110	0
4	GOL	A	711	6/6	0.83	0.23	57,69,78,79	0
3	NAG	A	703	14/15	0.86	0.32	75,85,93,100	0
4	GOL	A	710	6/6	0.86	0.17	62,64,66,76	0
4	GOL	A	709	6/6	0.86	0.29	76,77,81,82	0
4	GOL	A	706	6/6	0.88	0.31	47,63,65,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	713	6/6	0.88	0.14	59,69,79,82	0
5	SO4	A	716	5/5	0.88	0.37	87,88,101,103	0
5	SO4	A	717	5/5	0.92	0.41	84,102,112,116	0
5	SO4	A	715	5/5	0.96	0.18	49,55,62,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.