



wwPDB X-ray Structure Validation Summary Report ⓘ

May 30, 2020 – 03:08 am BST

PDB ID : 6MJN
Title : Crystal structure of an organic hydroperoxide resistance protein OsmC, predicted redox protein, regulator of sulfide bond formation from Legionella pneumophila
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2018-09-21
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

PERCENTILES INFOmissingINFO

1 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Organic hydroperoxide resistance protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	138	1027	645	180	198	4	0	6	0
1	B	140	1026	644	180	197	5	0	4	0
1	C	137	970	605	165	196	4	0	1	0
1	D	139	952	594	164	189	5	0	1	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A0A2S6F4M7
A	-6	ALA	-	expression tag	UNP A0A2S6F4M7
A	-5	HIS	-	expression tag	UNP A0A2S6F4M7
A	-4	HIS	-	expression tag	UNP A0A2S6F4M7
A	-3	HIS	-	expression tag	UNP A0A2S6F4M7
A	-2	HIS	-	expression tag	UNP A0A2S6F4M7
A	-1	HIS	-	expression tag	UNP A0A2S6F4M7
A	0	HIS	-	expression tag	UNP A0A2S6F4M7
B	-7	MET	-	expression tag	UNP A0A2S6F4M7
B	-6	ALA	-	expression tag	UNP A0A2S6F4M7
B	-5	HIS	-	expression tag	UNP A0A2S6F4M7
B	-4	HIS	-	expression tag	UNP A0A2S6F4M7
B	-3	HIS	-	expression tag	UNP A0A2S6F4M7
B	-2	HIS	-	expression tag	UNP A0A2S6F4M7
B	-1	HIS	-	expression tag	UNP A0A2S6F4M7
B	0	HIS	-	expression tag	UNP A0A2S6F4M7
C	-7	MET	-	expression tag	UNP A0A2S6F4M7
C	-6	ALA	-	expression tag	UNP A0A2S6F4M7
C	-5	HIS	-	expression tag	UNP A0A2S6F4M7
C	-4	HIS	-	expression tag	UNP A0A2S6F4M7
C	-3	HIS	-	expression tag	UNP A0A2S6F4M7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP A0A2S6F4M7
C	-1	HIS	-	expression tag	UNP A0A2S6F4M7
C	0	HIS	-	expression tag	UNP A0A2S6F4M7
D	-7	MET	-	expression tag	UNP A0A2S6F4M7
D	-6	ALA	-	expression tag	UNP A0A2S6F4M7
D	-5	HIS	-	expression tag	UNP A0A2S6F4M7
D	-4	HIS	-	expression tag	UNP A0A2S6F4M7
D	-3	HIS	-	expression tag	UNP A0A2S6F4M7
D	-2	HIS	-	expression tag	UNP A0A2S6F4M7
D	-1	HIS	-	expression tag	UNP A0A2S6F4M7
D	0	HIS	-	expression tag	UNP A0A2S6F4M7

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	A	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	80	Total O 82 82	0	2
3	B	72	Total O 72 72	0	0
3	C	30	Total O 30 30	0	0
3	D	19	Total O 19 19	0	0

2 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Organic hydroperoxide resistance protein

Chain A: 



- Molecule 1: Organic hydroperoxide resistance protein

Chain B: 



- Molecule 1: Organic hydroperoxide resistance protein

Chain C: 



- Molecule 1: Organic hydroperoxide resistance protein

Chain D: 



3 Data and refinement statistics

EDS was not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.55Å 51.40Å 82.34Å 90.00° 110.09° 90.00°	Depositor
Resolution (Å)	42.81 – 1.75	Depositor
% Data completeness (in resolution range)	92.8 (42.81-1.75)	Depositor
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 1.75Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.181 , 0.209	Depositor
Wilson B-factor (Å ²)	33.4	Xtrriage
Anisotropy	0.088	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4181	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1062	0.60	0/1443
1	B	0.41	1/1056 (0.1%)	0.59	1/1436 (0.1%)
1	C	0.38	1/988 (0.1%)	0.51	0/1351
1	D	0.30	0/970	0.48	0/1325
All	All	0.38	2/4076 (0.0%)	0.55	1/5555 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	120	CYS	CB-SG	-5.34	1.73	1.81
1	C	56	CYS	CB-SG	5.30	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	LEU	CA-CB-CG	5.23	127.32	115.30

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1027	0	1006	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1026	0	994	5	0
1	C	970	0	907	6	0
1	D	952	0	868	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	82	0	0	4	0
3	B	72	0	0	0	0
3	C	30	0	0	1	0
3	D	19	0	0	0	0
All	All	4181	0	3775	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:124[B]:ASN:OD1	3:C:201:HOH:O	1.94	0.84
1:A:124[A]:ASN:ND2	3:A:302:HOH:O	2.21	0.73
1:A:68:GLN:HE22	1:A:115:LYS:HD2	1.58	0.68
1:A:83:LEU:HD22	1:A:91:PHE:HB3	1.80	0.63
3:A:366:HOH:O	1:B:56:CYS:SG	2.57	0.59

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/147 (97%)	141 (99%)	1 (1%)	0	100	100
1	B	142/147 (97%)	142 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	136/147 (92%)	135 (99%)	1 (1%)	0	100	100
1	D	138/147 (94%)	135 (98%)	3 (2%)	0	100	100
All	All	558/588 (95%)	553 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/114 (91%)	103 (99%)	1 (1%)	76	63
1	B	102/114 (90%)	99 (97%)	3 (3%)	42	19
1	C	95/114 (83%)	94 (99%)	1 (1%)	73	60
1	D	86/114 (75%)	84 (98%)	2 (2%)	50	28
All	All	387/456 (85%)	380 (98%)	7 (2%)	57	40

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	72	LEU
1	D	134	LEU
1	C	53	TYR
1	B	4	LEU
1	D	53	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

5.3 Carbohydrates

EDS was not executed - this section is therefore empty.

5.4 Ligands

EDS was not executed - this section is therefore empty.

5.5 Other polymers

EDS was not executed - this section is therefore empty.