



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2021 – 04:10 PM EDT

PDB ID : 7MJ1  
Title : LarB, a carboxylase/hydrolase involved in synthesis of the cofactor for lactate racemase, in complex with NAD  
Authors : Chatterjee, S.; Rankin, J.A.; Lagishetty, S.; Hu, J.; Hausinger, R.P.  
Deposited on : 2021-04-19  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	
1	B	256	
1	C	256	
1	D	256	
1	E	256	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	256	<div><div></div><div>6%</div><div>62%</div><div>15%</div><div>•</div><div>21%</div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridinium-3,5-biscarboxylic acid mononucleotide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1377	876	234	261	6			
1	C	200	Total	C	N	O	S	0	0	0
			1358	859	231	261	7			
1	D	201	Total	C	N	O	S	0	5	0
			1347	861	226	252	8			
1	E	211	Total	C	N	O	S	0	0	0
			1419	905	236	272	6			
1	F	201	Total	C	N	O	S	0	0	0
			1365	872	225	260	8			
1	B	203	Total	C	N	O	S	0	0	0
			1389	884	234	264	7			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	ALA	-	expression tag	UNP F9UST0
A	248	SER	-	expression tag	UNP F9UST0
A	249	TRP	-	expression tag	UNP F9UST0
A	250	SER	-	expression tag	UNP F9UST0
A	251	HIS	-	expression tag	UNP F9UST0
A	252	PRO	-	expression tag	UNP F9UST0
A	253	GLN	-	expression tag	UNP F9UST0
A	254	PHE	-	expression tag	UNP F9UST0
A	255	GLU	-	expression tag	UNP F9UST0
A	256	LYS	-	expression tag	UNP F9UST0
C	247	ALA	-	expression tag	UNP F9UST0
C	248	SER	-	expression tag	UNP F9UST0
C	249	TRP	-	expression tag	UNP F9UST0
C	250	SER	-	expression tag	UNP F9UST0
C	251	HIS	-	expression tag	UNP F9UST0
C	252	PRO	-	expression tag	UNP F9UST0
C	253	GLN	-	expression tag	UNP F9UST0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	254	PHE	-	expression tag	UNP F9UST0
C	255	GLU	-	expression tag	UNP F9UST0
C	256	LYS	-	expression tag	UNP F9UST0
D	247	ALA	-	expression tag	UNP F9UST0
D	248	SER	-	expression tag	UNP F9UST0
D	249	TRP	-	expression tag	UNP F9UST0
D	250	SER	-	expression tag	UNP F9UST0
D	251	HIS	-	expression tag	UNP F9UST0
D	252	PRO	-	expression tag	UNP F9UST0
D	253	GLN	-	expression tag	UNP F9UST0
D	254	PHE	-	expression tag	UNP F9UST0
D	255	GLU	-	expression tag	UNP F9UST0
D	256	LYS	-	expression tag	UNP F9UST0
E	247	ALA	-	expression tag	UNP F9UST0
E	248	SER	-	expression tag	UNP F9UST0
E	249	TRP	-	expression tag	UNP F9UST0
E	250	SER	-	expression tag	UNP F9UST0
E	251	HIS	-	expression tag	UNP F9UST0
E	252	PRO	-	expression tag	UNP F9UST0
E	253	GLN	-	expression tag	UNP F9UST0
E	254	PHE	-	expression tag	UNP F9UST0
E	255	GLU	-	expression tag	UNP F9UST0
E	256	LYS	-	expression tag	UNP F9UST0
F	247	ALA	-	expression tag	UNP F9UST0
F	248	SER	-	expression tag	UNP F9UST0
F	249	TRP	-	expression tag	UNP F9UST0
F	250	SER	-	expression tag	UNP F9UST0
F	251	HIS	-	expression tag	UNP F9UST0
F	252	PRO	-	expression tag	UNP F9UST0
F	253	GLN	-	expression tag	UNP F9UST0
F	254	PHE	-	expression tag	UNP F9UST0
F	255	GLU	-	expression tag	UNP F9UST0
F	256	LYS	-	expression tag	UNP F9UST0
B	247	ALA	-	expression tag	UNP F9UST0
B	248	SER	-	expression tag	UNP F9UST0
B	249	TRP	-	expression tag	UNP F9UST0
B	250	SER	-	expression tag	UNP F9UST0
B	251	HIS	-	expression tag	UNP F9UST0
B	252	PRO	-	expression tag	UNP F9UST0
B	253	GLN	-	expression tag	UNP F9UST0
B	254	PHE	-	expression tag	UNP F9UST0
B	255	GLU	-	expression tag	UNP F9UST0

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	256	LYS	-	expression tag	UNP F9UST0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	C	2	Total Mg 2 2	0	0
2	E	1	Total Mg 1 1	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C N O P 44 21 7 14 2	0	0


- Molecule 4 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula:  $C_{21}H_{29}N_7O_{14}P_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		










- Molecule 1: Pyridinium-3,5-biscarboxylic acid mononucleotide synthase



- Molecule 1: Pyridinium-3,5-biscarboxylic acid mononucleotide synthase



- Molecule 1: Pyridinium-3,5-biscarboxylic acid mononucleotide synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.44Å 120.44Å 213.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.76 – 3.40 41.76 – 3.40	Depositor EDS
% Data completeness (in resolution range)	92.0 (41.76-3.40) 92.4 (41.76-3.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.243 , 0.269 0.243 , 0.268	Depositor DCC
$R_{free}$ test set	986 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	99.9	Xtriage
Anisotropy	0.437	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 65.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8347	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	110.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, MG, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1399	0.49	1/1922 (0.1%)
1	B	0.26	0/1410	0.46	0/1934
1	C	0.27	0/1377	0.49	0/1889
1	D	0.25	0/1368	0.46	0/1887
1	E	0.28	0/1442	0.66	4/1985 (0.2%)
1	F	0.33	0/1385	0.50	0/1900
All	All	0.28	0/8381	0.52	5/11517 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	150	TYR	CA-C-N	-10.72	93.61	117.20
1	E	150	TYR	O-C-N	9.59	138.04	122.70
1	E	150	TYR	C-N-CA	-7.82	102.15	121.70
1	E	151	ASP	CB-CG-OD1	-6.25	112.67	118.30
1	A	213	ALA	N-CA-CB	5.71	118.09	110.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	150	TYR	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1377	0	1365	32	0
1	B	1389	0	1378	23	0
1	C	1358	0	1319	30	0
1	D	1347	0	1281	19	1
1	E	1419	0	1360	32	1
1	F	1365	0	1332	32	1
2	A	1	0	0	0	0
2	C	2	0	0	0	0
2	E	1	0	0	0	0
3	C	44	0	25	5	0
4	E	44	0	26	9	0
All	All	8347	0	8086	158	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:218:LEU:HD12	1:F:227:VAL:HG11	1.36	1.07
1:F:218:LEU:HD12	1:F:227:VAL:CG1	1.92	0.98
1:E:204:TYR:OH	1:F:219:ASN:HA	1.77	0.84
1:F:218:LEU:CD1	1:F:227:VAL:HG11	2.09	0.82
1:D:199:PRO:HG2	1:D:230:ILE:HA	1.64	0.79

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:234:PHE:CE2	1:F:223:SER:OG[3_545]	1.65	0.55

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:206:THR:OG1	1:D:229:ASN:OD1[3_445]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/256 (78%)	181 (91%)	17 (8%)	1 (0%)	29	61
1	B	201/256 (78%)	182 (90%)	19 (10%)	0	100	100
1	C	196/256 (77%)	180 (92%)	14 (7%)	2 (1%)	15	46
1	D	204/256 (80%)	178 (87%)	26 (13%)	0	100	100
1	E	209/256 (82%)	182 (87%)	25 (12%)	2 (1%)	15	46
1	F	197/256 (77%)	181 (92%)	16 (8%)	0	100	100
All	All	1206/1536 (78%)	1084 (90%)	117 (10%)	5 (0%)	34	67

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	74	LEU
1	E	48	PHE
1	E	39	ASP
1	A	181	GLY
1	C	202	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/191 (70%)	124 (92%)	10 (8%)	13	41
1	B	135/191 (71%)	129 (96%)	6 (4%)	28	58
1	C	128/191 (67%)	122 (95%)	6 (5%)	26	57
1	D	120/191 (63%)	109 (91%)	11 (9%)	9	31
1	E	131/191 (69%)	126 (96%)	5 (4%)	33	61
1	F	130/191 (68%)	122 (94%)	8 (6%)	18	48
All	All	778/1146 (68%)	732 (94%)	46 (6%)	20	49

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	73	THR
1	F	63	VAL
1	E	76	ILE
1	F	50	GLU
1	F	217	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAI	E	302	1,2	42,48,48	0.94	3 (7%)	47,73,73	1.09	4 (8%)
3	NAD	C	302	2	42,48,48	3.74	18 (42%)	50,73,73	2.49	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAI	E	302	1,2	-	10/25/72/72	0/5/5/5
3	NAD	C	302	2	-	9/26/62/62	0/5/5/5

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	302	NAD	C2D-C3D	-10.76	1.23	1.53
3	C	302	NAD	C3B-C4B	-8.68	1.30	1.53
3	C	302	NAD	O4B-C4B	7.70	1.62	1.45
3	C	302	NAD	O4D-C1D	7.42	1.51	1.41
3	C	302	NAD	O4B-C1B	-7.06	1.31	1.41

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	302	NAD	C1B-N9A-C4A	13.29	149.99	126.64
3	C	302	NAD	C5A-C6A-N6A	5.92	129.34	120.35
3	C	302	NAD	N3A-C2A-N1A	-5.46	120.14	128.68
3	C	302	NAD	N6A-C6A-N1A	-4.29	109.67	118.57
4	E	302	NAI	C1D-N1N-C2N	3.36	126.70	121.11

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	302	NAD	C5B-O5B-PA-O3
4	E	302	NAI	C5B-O5B-PA-O1A

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	E	302	NAI	O4D-C4D-C5D-O5D
4	E	302	NAI	O4D-C1D-N1N-C2N
4	E	302	NAI	C2N-C3N-C7N-N7N


There are no ring outliers.


2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	302	NAI	9	0
3	C	302	NAD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	201/256 (78%)	0.04	9 (4%) 33 33	39, 99, 196, 262	1 (0%)
1	B	203/256 (79%)	0.02	5 (2%) 57 55	42, 90, 159, 193	0
1	C	200/256 (78%)	-0.01	11 (5%) 25 25	45, 93, 181, 255	0
1	D	201/256 (78%)	0.17	15 (7%) 14 16	45, 110, 187, 245	0
1	E	211/256 (82%)	0.13	4 (1%) 66 65	65, 113, 173, 221	0
1	F	201/256 (78%)	0.33	16 (7%) 12 13	61, 121, 210, 294	0
All	All	1217/1536 (79%)	0.11	60 (4%) 29 29	39, 106, 189, 294	1 (0%)

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	96	ALA	4.9
1	F	95	THR	4.8
1	F	96	ALA	4.6
1	D	93	LEU	4.3
1	E	110	GLN	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	E	301	1/1	0.81	0.09	112,112,112,112	0
2	MG	A	301	1/1	0.85	0.41	67,67,67,67	0
2	MG	C	303	1/1	0.87	0.11	90,90,90,90	0
3	NAD	C	302	44/44	0.90	0.22	116,124,141,144	0
2	MG	C	301	1/1	0.91	0.17	76,76,76,76	0
4	NAI	E	302	44/44	0.92	0.19	99,108,115,122	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around NAI E 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.