



Full wwPDB X-ray Structure Validation Report ⓘ

May 12, 2020 – 11:12 pm BST

PDB ID : 4MKR
Title : Structure of the apo form of a Zingiber officinale double bond reductase
Authors : Buratto, J.; Langlois d'Estaintot, B.; Granier, T.; Gallois, B.; Willis, M.A.; Sang, Y.; Flores-Sanchez, I.J.; Gang, D.R.
Deposited on : 2013-09-05
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>8%</div> </div> </div>
1	B	358	<div> <div>11%</div> <div> <div></div> <div>58%</div> <div>6%</div> <div>36%</div> </div> </div>
1	C	358	<div> <div></div> <div> <div></div> <div>82%</div> <div>10%</div> <div>8%</div> </div> </div>
1	D	358	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>9%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zingiber officinale double bond reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	0	0	0
			2518	1643	406	457	12			
1	B	228	Total	C	N	O	S	0	0	0
			1644	1069	271	297	7			
1	C	330	Total	C	N	O	S	0	1	0
			2521	1644	409	456	12			
1	D	327	Total	C	N	O	S	0	0	0
			2442	1591	396	444	11			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	44	Total	O	0	0
			44	44		
2	B	12	Total	O	0	0
			12	12		
2	C	43	Total	O	0	0
			43	43		
2	D	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Zingiber officinale double bond reductase




- Molecule 1: Zingiber officinale double bond reductase



- Molecule 1: Zingiber officinale double bond reductase



● Molecule 1: Zingiber officinale double bond reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.93Å 76.34Å 93.45Å 80.15° 89.97° 85.66°	Depositor
Resolution (Å)	46.03 – 2.58 46.03 – 2.58	Depositor EDS
% Data completeness (in resolution range)	97.6 (46.03-2.58) 97.7 (46.03-2.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.58Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.228 , 0.282 0.230 , 0.283	Depositor DCC
R_{free} test set	2130 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.6	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9244	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/2575	0.87	3/3483 (0.1%)
1	B	0.63	0/1680	0.81	0/2278
1	C	0.78	0/2583	0.88	1/3500 (0.0%)
1	D	0.66	0/2498	0.78	1/3393 (0.0%)
All	All	0.73	0/9336	0.84	5/12654 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	GLY	N-CA-C	5.65	127.22	113.10
1	A	255	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	D	251	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	65	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	C	63	ARG	NE-CZ-NH1	-5.09	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2491	26	0
1	B	1644	0	1480	15	0
1	C	2521	0	2478	27	0
1	D	2442	0	2322	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	0	1	0
2	B	12	0	0	0	0
2	C	43	0	0	1	0
2	D	20	0	0	4	0
All	All	9244	0	8771	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:GLU:CB	1:A:275:ASN:HD22	2.15	0.59
1:D:65:ARG:NH1	1:D:76:ASP:O	2.35	0.59
1:A:132:ILE:HG22	1:A:133:PRO:HD2	1.84	0.58
1:D:144:VAL:HG23	2:D:407:HOH:O	2.05	0.57
1:B:254:GLY:C	1:B:255:ARG:HG3	2.25	0.56
1:B:244:GLU:O	1:B:275:ASN:ND2	2.37	0.56
1:A:254:GLY:C	1:A:255:ARG:HG3	2.25	0.56
1:C:62:MET:CE	1:C:331:LEU:HD13	2.36	0.55
1:A:244:GLU:HB3	1:A:275:ASN:HD22	1.71	0.55
1:A:139:GLY:O	1:A:142:GLY:N	2.31	0.55
1:D:62:MET:CE	1:D:331:LEU:HD13	2.37	0.55
1:C:63:ARG:NH1	1:C:67:THR:OG1	2.40	0.54
1:C:254:GLY:C	1:C:255:ARG:HG3	2.27	0.54
1:D:42:VAL:CG1	1:D:46:SER:OG	2.56	0.54
1:A:244:GLU:CA	1:A:275:ASN:HD22	2.22	0.53
1:D:241:PRO:HD2	2:D:415:HOH:O	2.07	0.53
1:D:57:SER:OG	1:D:342:LYS:NZ	2.32	0.53
1:C:244:GLU:HA	1:C:275:ASN:HD22	1.74	0.52
1:D:244:GLU:O	1:D:275:ASN:ND2	2.42	0.52
1:C:107:TRP:CZ2	1:C:125:PHE:CD2	2.97	0.52
1:A:276:LEU:HD22	1:A:279:ILE:HD12	1.92	0.51
1:D:62:MET:HE2	1:D:331:LEU:CD1	2.41	0.51
1:C:34:THR:HG22	1:C:34:THR:O	2.11	0.50
1:A:132:ILE:O	1:A:133:PRO:C	2.49	0.50
1:C:232:ASP:OD1	1:C:251:ARG:HD3	2.11	0.50
1:C:57:SER:OG	1:C:59:ASP:OD1	2.31	0.49
1:C:139:GLY:O	1:C:144:VAL:HB	2.12	0.49
1:C:293:ASP:OD2	1:D:283:ARG:HD3	2.12	0.49
1:D:244:GLU:CA	1:D:275:ASN:HD22	2.25	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:SER:OG	1:A:59:ASP:OD1	2.30	0.49
1:C:59:ASP:OD1	1:C:342:LYS:NZ	2.45	0.49
1:C:57:SER:HB2	1:C:138:THR:HB	1.95	0.48
1:B:244:GLU:CA	1:B:275:ASN:HD22	2.27	0.48
1:C:276:LEU:O	1:C:279:ILE:HB	2.14	0.48
1:D:244:GLU:HA	1:D:275:ASN:HD22	1.78	0.48
1:A:283:ARG:HD3	1:B:293:ASP:OD2	2.13	0.47
1:D:254:GLY:C	1:D:255:ARG:HG3	2.33	0.47
1:D:62:MET:HE2	1:D:331:LEU:HD13	1.95	0.47
1:A:293:ASP:OD2	1:B:283:ARG:HD3	2.14	0.47
1:C:283:ARG:HD3	1:D:293:ASP:OD2	2.13	0.47
1:D:127:ILE:HG23	1:D:137:TYR:CZ	2.50	0.47
1:D:11:ASN:HB2	1:D:38:ILE:HG13	1.95	0.47
1:D:172:ALA:HB2	1:D:342:LYS:HB2	1.96	0.47
1:A:182:LYS:NZ	1:A:208:ASP:OD2	2.47	0.47
1:B:88:GLY:HA2	1:B:138:THR:HG22	1.95	0.47
1:B:276:LEU:O	1:B:279:ILE:HB	2.15	0.47
1:D:232:ASP:OD1	1:D:251:ARG:HD3	2.15	0.47
1:A:47:ASN:O	1:A:97:HIS:HB2	2.15	0.47
1:A:277:PHE:N	1:B:261:MET:HE1	2.30	0.46
1:A:274:HIS:ND1	1:A:274:HIS:N	2.63	0.46
1:A:276:LEU:O	1:A:279:ILE:HB	2.16	0.46
1:C:65:ARG:NH1	2:C:431:HOH:O	2.34	0.46
1:A:59:ASP:HB2	1:A:61:TYR:CE2	2.51	0.46
1:D:38:ILE:HG22	2:D:413:HOH:O	2.16	0.45
1:D:11:ASN:OD1	1:D:114:GLU:HB2	2.17	0.45
1:C:241:PRO:HD3	1:C:271:GLU:HB2	1.99	0.45
1:C:62:MET:CE	1:C:331:LEU:CD1	2.95	0.45
1:D:58:CYS:HB2	1:D:343:GLN:HG2	1.99	0.45
1:C:244:GLU:O	1:C:275:ASN:ND2	2.49	0.44
1:A:129:HIS:HB2	1:A:137:TYR:OH	2.18	0.44
1:C:9:VAL:HG21	1:C:40:LEU:HD21	2.00	0.44
1:D:248:HIS:CE1	2:D:419:HOH:O	2.71	0.43
1:A:325:GLU:CD	1:A:325:GLU:H	2.20	0.43
1:C:244:GLU:CA	1:C:275:ASN:HD22	2.31	0.43
1:D:106:VAL:HG12	1:D:126:LYS:HA	2.00	0.43
1:A:57:SER:OG	1:A:342:LYS:NZ	2.37	0.43
1:B:198:VAL:HG21	1:B:212:ASN:HB2	2.01	0.43
1:D:9:VAL:HG21	1:D:40:LEU:HD11	2.01	0.43
1:B:298:TYR:O	1:B:299:PRO:C	2.57	0.43
1:A:234:TYR:HB3	1:A:256:ILE:HG12	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:336:GLU:OE1	1:C:338:ARG:NH1	2.48	0.42
1:D:234:TYR:HB3	1:D:256:ILE:HG12	2.00	0.42
1:A:34:THR:O	1:A:34:THR:HG22	2.20	0.42
1:C:327:ALA:N	1:C:328:PRO:CD	2.83	0.42
1:C:253:LYS:O	1:C:255:ARG:NH1	2.48	0.42
1:B:57:SER:CB	1:B:138:THR:HB	2.50	0.42
1:B:150:VAL:O	1:B:154:ASP:HB3	2.19	0.42
1:A:313:LYS:NZ	2:A:441:HOH:O	2.53	0.41
1:B:232:ASP:OD1	1:B:251:ARG:HD3	2.20	0.41
1:D:176:ILE:HG22	1:D:180:PHE:CE2	2.55	0.41
1:B:244:GLU:HA	1:B:275:ASN:HD22	1.84	0.41
1:A:20:ILE:HD11	1:A:66:MET:HE2	2.02	0.41
1:C:78:PHE:CZ	1:C:84:ILE:HG23	2.55	0.41
1:C:107:TRP:CH2	1:C:125:PHE:CE2	3.08	0.41
1:C:112:TRP:CZ3	1:C:327:ALA:HB1	2.56	0.41
1:C:283:ARG:CD	1:D:293:ASP:OD2	2.68	0.41
1:A:232:ASP:OD1	1:A:251:ARG:HD3	2.21	0.41
1:B:238:VAL:O	1:B:262:ILE:N	2.53	0.41
1:A:175:GLN:HE22	1:A:341:GLY:HA2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/358 (89%)	310 (97%)	10 (3%)	0	100	100
1	B	216/358 (60%)	206 (95%)	10 (5%)	0	100	100
1	C	325/358 (91%)	315 (97%)	10 (3%)	0	100	100
1	D	319/358 (89%)	312 (98%)	7 (2%)	0	100	100
All	All	1180/1432 (82%)	1143 (97%)	37 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/300 (87%)	257 (98%)	4 (2%)	65	82
1	B	145/300 (48%)	145 (100%)	0	100	100
1	C	259/300 (86%)	257 (99%)	2 (1%)	81	92
1	D	240/300 (80%)	238 (99%)	2 (1%)	81	92
All	All	905/1200 (75%)	897 (99%)	8 (1%)	78	90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ILE
1	A	44	GLU
1	A	76	ASP
1	A	348	SER
1	C	49	VAL
1	C	76	ASP
1	D	322	GLU
1	D	342	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	ASN
1	B	275	ASN
1	C	275	ASN
1	D	275	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/358 (91%)	0.23	3 (0%) 84 83	23, 39, 58, 78	0
1	B	228/358 (63%)	0.95	39 (17%) 1 1	29, 59, 112, 152	0
1	C	330/358 (92%)	0.18	1 (0%) 94 94	22, 37, 56, 79	0
1	D	327/358 (91%)	0.44	19 (5%) 23 19	31, 52, 76, 88	0
All	All	1213/1432 (84%)	0.41	62 (5%) 28 24	22, 45, 81, 152	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	194	SER	6.6
1	B	193	GLY	6.1
1	B	115	TYR	4.5
1	B	111	GLY	4.3
1	B	169	ALA	4.2
1	D	348	SER	4.1
1	D	170	ALA	4.0
1	B	344	VAL	4.0
1	B	135	SER	3.9
1	B	306	VAL	3.8
1	B	54	LEU	3.7
1	B	107	TRP	3.7
1	D	205	PHE	3.5
1	D	169	ALA	3.5
1	D	19	PHE	3.5
1	D	40	LEU	3.5
1	B	49	VAL	3.5
1	B	136	TYR	3.4
1	B	261	MET	3.4
1	B	200	LEU	3.4
1	B	143	MET	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	138	THR	3.4
1	D	340	VAL	3.3
1	B	53	ASN	3.1
1	B	199	ASN	3.1
1	B	56	LEU	3.1
1	B	86	GLY	3.1
1	D	194	SER	3.0
1	C	214	LYS	2.9
1	B	262	ILE	2.9
1	D	199	ASN	2.9
1	B	59	ASP	2.8
1	D	213	TYR	2.8
1	D	20	ILE	2.7
1	B	137	TYR	2.7
1	D	217	PRO	2.7
1	B	55	TYR	2.7
1	B	87	PHE	2.6
1	A	132	ILE	2.6
1	B	205	PHE	2.6
1	B	346	VAL	2.6
1	A	75	VAL	2.6
1	D	115	TYR	2.6
1	B	110	ILE	2.5
1	B	142	GLY	2.4
1	B	121	PRO	2.4
1	B	345	VAL	2.3
1	D	349	ARG	2.3
1	D	238	VAL	2.3
1	A	45	GLY	2.3
1	B	114	GLU	2.3
1	B	140	ILE	2.2
1	D	320	ILE	2.2
1	D	46	SER	2.1
1	D	319	ASP	2.1
1	B	106	VAL	2.1
1	B	88	GLY	2.1
1	D	210	ALA	2.1
1	B	294	HIS	2.1
1	B	212	ASN	2.0
1	B	347	VAL	2.0
1	B	117	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.