



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 05:31 pm BST

PDB ID : 4MLG  
Title : Structure of RS223-Beta-xylosidase  
Authors : Jordan, D.; Braker, J.; Wagschal, K.; Lee, C.; Dubrovskaya, I.; Anderson, S.;  
Wawrzak, Z.  
Deposited on : 2013-09-06  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>13%</div> </div> </div>
1	B	324	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>13%</div> </div> </div>
1	C	324	<div> <div>8%</div> <div> <div></div> <div>87%</div> <div>12%</div> </div> </div>
1	D	324	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>16%</div> </div> </div>
1	E	324	<div> <div>6%</div> <div> <div></div> <div>85%</div> <div>15%</div> </div> </div>
1	F	324	<div> <div>9%</div> <div> <div></div> <div>83%</div> <div>17%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	324	
1	H	324	
1	I	324	
1	J	324	
1	K	324	
1	L	324	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	402	-	-	-	X
3	SO4	F	402	-	-	-	X
3	SO4	G	402	-	-	-	X
3	SO4	G	403	-	-	-	X
3	SO4	I	403	-	-	-	X
3	SO4	J	403	-	-	-	X
3	SO4	K	402	-	-	-	X
3	SO4	K	403	-	-	-	X
3	SO4	L	403	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 32402 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-xylosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2640	1699	438	488	15			
1	B	323	Total	C	N	O	S	0	0	0
			2640	1699	438	488	15			
1	C	323	Total	C	N	O	S	0	0	0
			2636	1696	437	488	15			
1	D	322	Total	C	N	O	S	0	0	0
			2635	1696	437	487	15			
1	E	323	Total	C	N	O	S	0	0	0
			2640	1699	438	488	15			
1	F	323	Total	C	N	O	S	0	0	0
			2636	1696	437	488	15			
1	G	324	Total	C	N	O	S	0	0	0
			2641	1699	438	489	15			
1	H	324	Total	C	N	O	S	0	0	0
			2645	1701	438	491	15			
1	I	321	Total	C	N	O	S	0	0	0
			2615	1684	434	483	14			
1	J	323	Total	C	N	O	S	0	0	0
			2636	1696	437	488	15			
1	K	324	Total	C	N	O	S	0	0	0
			2645	1701	438	491	15			
1	L	322	Total	C	N	O	S	0	0	0
			2631	1694	436	486	15			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Ca	0	0
			1	1		
2	J	1	Total	Ca	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Ca	0	0
			1	1		
2	K	1	Total	Ca	0	0
			1	1		
2	E	1	Total	Ca	0	0
			1	1		
2	H	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	I	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	L	1	Total	Ca	0	0
			1	1		
2	F	1	Total	Ca	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		
3	I	1	Total	O	S	0	0
			5	4	1		
3	J	1	Total	O	S	0	0
			5	4	1		
3	J	1	Total	O	S	0	0
			5	4	1		
3	K	1	Total	O	S	0	0
			5	4	1		
3	K	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	72	Total	O	0	0
			72	72		
4	B	64	Total	O	0	0
			64	64		
4	C	45	Total	O	0	0
			45	45		
4	D	68	Total	O	0	0
			68	68		
4	E	50	Total	O	0	0
			50	50		
4	F	58	Total	O	0	0
			58	58		
4	G	47	Total	O	0	0
			47	47		
4	H	50	Total	O	0	0
			50	50		
4	I	49	Total	O	0	0
			49	49		
4	J	50	Total	O	0	0
			50	50		
4	K	44	Total	O	0	0
			44	44		
4	L	33	Total	O	0	0
			33	33		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Beta-xylosidase




#### • Molecule 1: Beta-xylosidase




#### • Molecule 1: Beta-xylosidase





#### • Molecule 1: Beta-xylosidase









• Molecule 1: Beta-xylosidase





• Molecule 1: Beta-xylosidase




• Molecule 1: Beta-xylosidase





• Molecule 1: Beta-xylosidase









• Molecule 1: Beta-xylosidase





• Molecule 1: Beta-xylosidase




• Molecule 1: Beta-xylosidase



• Molecule 1: Beta-xylosidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.53Å 148.70Å 125.65Å 90.00° 112.87° 90.00°	Depositor
Resolution (Å)	29.20 – 2.70 29.17 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.3 (29.20-2.70) 91.8 (29.17-2.69)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.68Å)	Xtriage
Refinement program	PHENIX 1.8.1 _1168	Depositor
R, $R_{free}$	0.210 , 0.257 0.211 , 0.258	Depositor DCC
$R_{free}$ test set	5761 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 37.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.033 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	32402	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.22	0/2729	0.39	0/3706
1	B	0.22	0/2729	0.38	0/3706
1	C	0.21	0/2725	0.38	0/3702
1	D	0.22	0/2724	0.38	0/3699
1	E	0.22	0/2729	0.38	0/3706
1	F	0.23	0/2725	0.38	0/3702
1	G	0.21	0/2730	0.37	0/3709
1	H	0.22	0/2734	0.39	0/3714
1	I	0.21	0/2704	0.37	0/3675
1	J	0.22	0/2725	0.38	0/3702
1	K	0.22	0/2734	0.38	0/3714
1	L	0.21	0/2720	0.37	0/3694
All	All	0.22	0/32708	0.38	0/44429

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2640	0	2490	20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2640	0	2490	21	0
1	C	2636	0	2479	25	0
1	D	2635	0	2485	26	0
1	E	2640	0	2490	28	0
1	F	2636	0	2479	41	0
1	G	2641	0	2481	34	0
1	H	2645	0	2485	30	0
1	I	2615	0	2455	26	0
1	J	2636	0	2479	27	0
1	K	2645	0	2485	30	0
1	L	2631	0	2479	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
3	D	10	0	0	0	0
3	E	10	0	0	0	0
3	F	10	0	0	0	0
3	G	10	0	0	0	0
3	H	10	0	0	0	0
3	I	10	0	0	0	0
3	J	10	0	0	0	0
3	K	10	0	0	0	0
3	L	10	0	0	0	0
4	A	72	0	0	0	0
4	B	64	0	0	0	0
4	C	45	0	0	0	0
4	D	68	0	0	0	0
4	E	50	0	0	2	0
4	F	58	0	0	0	0
4	G	47	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	50	0	0	2	0
4	I	49	0	0	1	0
4	J	50	0	0	1	0
4	K	44	0	0	1	0
4	L	33	0	0	0	0
All	All	32402	0	29777	328	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 328 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:265:LEU:HD12	1:G:266:THR:H	1.21	1.05
1:F:240:THR:CG2	1:F:243:THR:HB	2.13	0.79
1:G:265:LEU:HD12	1:G:266:THR:N	1.98	0.78
1:F:240:THR:HG22	1:F:243:THR:HB	1.69	0.74
1:L:266:THR:HG23	1:L:267:PRO:HD2	1.70	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/324 (99%)	302 (94%)	19 (6%)	0	100	100
1	B	321/324 (99%)	303 (94%)	18 (6%)	0	100	100
1	C	321/324 (99%)	303 (94%)	18 (6%)	0	100	100
1	D	320/324 (99%)	304 (95%)	16 (5%)	0	100	100
1	E	321/324 (99%)	304 (95%)	17 (5%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	321/324 (99%)	300 (94%)	21 (6%)	0	100	100
1	G	322/324 (99%)	298 (92%)	24 (8%)	0	100	100
1	H	322/324 (99%)	304 (94%)	18 (6%)	0	100	100
1	I	319/324 (98%)	296 (93%)	23 (7%)	0	100	100
1	J	321/324 (99%)	300 (94%)	21 (6%)	0	100	100
1	K	322/324 (99%)	302 (94%)	20 (6%)	0	100	100
1	L	320/324 (99%)	301 (94%)	19 (6%)	0	100	100
All	All	3851/3888 (99%)	3617 (94%)	234 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/281 (100%)	276 (99%)	4 (1%)	67	86
1	B	280/281 (100%)	275 (98%)	5 (2%)	59	83
1	C	279/281 (99%)	276 (99%)	3 (1%)	73	90
1	D	280/281 (100%)	274 (98%)	6 (2%)	53	80
1	E	280/281 (100%)	278 (99%)	2 (1%)	84	94
1	F	279/281 (99%)	277 (99%)	2 (1%)	84	94
1	G	279/281 (99%)	275 (99%)	4 (1%)	67	86
1	H	280/281 (100%)	277 (99%)	3 (1%)	73	90
1	I	276/281 (98%)	273 (99%)	3 (1%)	73	90
1	J	279/281 (99%)	276 (99%)	3 (1%)	73	90
1	K	280/281 (100%)	273 (98%)	7 (2%)	47	76
1	L	279/281 (99%)	273 (98%)	6 (2%)	52	79
All	All	3351/3372 (99%)	3303 (99%)	48 (1%)	67	86

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	220	ARG
1	H	135	MET
1	L	269	VAL
1	G	108	PHE
1	G	220	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	289	HIS
1	G	45	HIS
1	I	52	HIS
1	K	274	HIS
1	L	165	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 36 ligands modelled in this entry, 12 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	B	403	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	F	402	-	4,4,4	0.15	0	6,6,6	0.06	0
3	SO4	K	402	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	D	402	-	4,4,4	0.14	0	6,6,6	0.07	0
3	SO4	I	403	-	4,4,4	0.14	0	6,6,6	0.04	0
3	SO4	C	402	-	4,4,4	0.15	0	6,6,6	0.04	0
3	SO4	H	403	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	G	403	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	C	403	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	L	402	-	4,4,4	0.14	0	6,6,6	0.06	0
3	SO4	B	402	-	4,4,4	0.14	0	6,6,6	0.07	0
3	SO4	I	402	-	4,4,4	0.15	0	6,6,6	0.05	0
3	SO4	G	402	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	E	402	-	4,4,4	0.14	0	6,6,6	0.04	0
3	SO4	A	403	-	4,4,4	0.14	0	6,6,6	0.04	0
3	SO4	D	403	-	4,4,4	0.13	0	6,6,6	0.07	0
3	SO4	J	403	-	4,4,4	0.15	0	6,6,6	0.05	0
3	SO4	A	402	-	4,4,4	0.14	0	6,6,6	0.08	0
3	SO4	J	402	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	E	403	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	L	403	-	4,4,4	0.14	0	6,6,6	0.04	0
3	SO4	K	403	-	4,4,4	0.13	0	6,6,6	0.05	0
3	SO4	F	403	-	4,4,4	0.14	0	6,6,6	0.07	0
3	SO4	H	402	-	4,4,4	0.14	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	323/324 (99%)	0.27	10 (3%) 49 49	43, 60, 88, 99	0
1	B	323/324 (99%)	0.40	13 (4%) 38 37	45, 67, 86, 105	0
1	C	323/324 (99%)	0.55	25 (7%) 13 11	50, 78, 112, 146	0
1	D	322/324 (99%)	0.31	7 (2%) 62 63	43, 65, 91, 121	0
1	E	323/324 (99%)	0.38	18 (5%) 24 23	48, 66, 91, 120	0
1	F	323/324 (99%)	0.67	28 (8%) 10 8	49, 75, 104, 131	0
1	G	324/324 (100%)	0.57	17 (5%) 27 25	46, 78, 117, 183	0
1	H	324/324 (100%)	0.55	25 (7%) 13 11	51, 75, 105, 159	0
1	I	321/324 (99%)	0.76	29 (9%) 9 7	53, 76, 109, 148	0
1	J	323/324 (99%)	0.47	14 (4%) 35 33	50, 69, 99, 118	0
1	K	324/324 (100%)	0.70	18 (5%) 24 23	54, 77, 103, 140	0
1	L	322/324 (99%)	0.97	45 (13%) 2 1	58, 89, 123, 151	0
All	All	3875/3888 (99%)	0.55	249 (6%) 19 18	43, 72, 107, 183	0

The worst 5 of 249 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	323	ALA	9.0
1	H	324	GLU	6.6
1	L	42	ASN	5.4
1	J	322	THR	4.9
1	K	177	ASP	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	K	403	5/5	0.11	0.69	161,161,161,161	0
3	SO4	G	403	5/5	0.54	0.43	148,149,149,149	0
3	SO4	L	403	5/5	0.60	0.53	152,153,153,153	0
3	SO4	F	402	5/5	0.60	0.52	106,106,107,107	0
3	SO4	E	403	5/5	0.64	0.27	131,131,131,131	0
3	SO4	J	403	5/5	0.66	0.48	153,154,154,154	0
3	SO4	B	403	5/5	0.70	0.18	116,116,116,116	0
3	SO4	F	403	5/5	0.71	0.29	136,136,137,137	0
3	SO4	A	403	5/5	0.72	0.34	123,123,124,124	0
3	SO4	C	403	5/5	0.72	0.29	137,137,137,137	0
3	SO4	I	403	5/5	0.73	0.45	162,162,163,163	0
3	SO4	A	402	5/5	0.73	0.57	109,109,109,109	0
3	SO4	K	402	5/5	0.73	0.49	110,110,111,111	0
3	SO4	J	402	5/5	0.76	0.32	101,101,102,102	0
3	SO4	H	403	5/5	0.77	0.25	113,113,114,114	0
3	SO4	C	402	5/5	0.78	0.35	103,103,104,104	0
3	SO4	G	402	5/5	0.78	0.51	126,126,126,126	0
3	SO4	D	403	5/5	0.80	0.27	113,113,113,114	0
2	CA	B	401	1/1	0.80	0.51	73,73,73,73	0
3	SO4	I	402	5/5	0.81	0.41	105,105,105,106	0
2	CA	K	401	1/1	0.82	0.54	81,81,81,81	0
3	SO4	E	402	5/5	0.83	0.41	105,105,105,105	0
2	CA	H	401	1/1	0.84	0.49	71,71,71,71	0
2	CA	G	401	1/1	0.84	0.43	77,77,77,77	0
3	SO4	L	402	5/5	0.85	0.36	111,111,111,111	0
3	SO4	H	402	5/5	0.87	0.30	94,94,94,95	0
2	CA	J	401	1/1	0.88	0.37	70,70,70,70	0
2	CA	F	401	1/1	0.89	0.53	84,84,84,84	0
3	SO4	B	402	5/5	0.89	0.35	85,85,86,86	0
2	CA	A	401	1/1	0.91	0.50	62,62,62,62	0
3	SO4	D	402	5/5	0.91	0.36	92,93,93,93	0
2	CA	D	401	1/1	0.91	0.35	65,65,65,65	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	L	401	1/1	0.92	0.49	80,80,80,80	0
2	CA	C	401	1/1	0.92	0.49	77,77,77,77	0
2	CA	E	401	1/1	0.95	0.48	70,70,70,70	0
2	CA	I	401	1/1	0.96	0.61	78,78,78,78	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.