



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 11:55 pm BST

PDB ID : 2MRJ
Title : Structure of Fyn protein SH2 bound
Authors : Huculeci, R.; Buts, L.; Lenaerts, T.; Van Nuland, N.
Deposited on : 2014-07-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	112	

2 Ensemble composition and analysis

This entry contains 18 models. Model 17 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:149-A:244 (96)	0.41	17

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 6, 10, 12
2	15, 16, 17
3	7, 11, 13
4	1, 5
5	4, 9
Single-model clusters	3; 8; 14; 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1635 atoms, of which 812 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Tyrosine-protein kinase Fyn.

Mol	Chain	Residues	Atoms						Trace
1	A	100	Total	C	H	N	O	S	0
			1635	520	812	152	147	4	

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	137	SER	-	EXPRESSION TAG	UNP P06241
A	138	SER	-	EXPRESSION TAG	UNP P06241
A	139	GLY	-	EXPRESSION TAG	UNP P06241
A	140	LEU	-	EXPRESSION TAG	UNP P06241
A	141	VAL	-	EXPRESSION TAG	UNP P06241
A	142	PRO	-	EXPRESSION TAG	UNP P06241
A	143	ARG	-	EXPRESSION TAG	UNP P06241
A	144	GLY	-	EXPRESSION TAG	UNP P06241
A	145	SER	-	EXPRESSION TAG	UNP P06241
A	146	HIS	-	EXPRESSION TAG	UNP P06241
A	147	MET	-	EXPRESSION TAG	UNP P06241

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Tyrosine-protein kinase Fyn




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Tyrosine-protein kinase Fyn



4.2.2 Score per residue for model 2

- Molecule 1: Tyrosine-protein kinase Fyn






4.2.3 Score per residue for model 3

- Molecule 1: Tyrosine-protein kinase Fyn


Chain A: 58% 23% 11%



4.2.4 Score per residue for model 4

- Molecule 1: Tyrosine-protein kinase Fyn


Chain A: 55% 26% 11%



4.2.5 Score per residue for model 5

- Molecule 1: Tyrosine-protein kinase Fyn


Chain A: 58% 22% 5% 11%



4.2.6 Score per residue for model 6

- Molecule 1: Tyrosine-protein kinase Fyn

Chain A:  62% 21% • • 11%



4.2.7 Score per residue for model 7

- Molecule 1: Tyrosine-protein kinase Fyn

Chain A:  69% 15% • • 11%



4.2.8 Score per residue for model 8

- Molecule 1: Tyrosine-protein kinase Fyn


Chain A:  60% 21% • • 11%



4.2.9 Score per residue for model 9


- Molecule 1: Tyrosine-protein kinase Fyn

Chain A:  63% 21% • • 11%




4.2.10 Score per residue for model 10

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.11 Score per residue for model 11

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.12 Score per residue for model 12

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.13 Score per residue for model 13

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.14 Score per residue for model 14

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.15 Score per residue for model 15

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.16 Score per residue for model 16

- Molecule 1: Tyrosine-protein kinase Fyn




4.2.17 Score per residue for model 17 (medoid)

- Molecule 1: Tyrosine-protein kinase Fyn



4.2.18 Score per residue for model 18

- Molecule 1: Tyrosine-protein kinase Fyn



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 18 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TALOS	geometry optimization	
CYANA	structure solution	
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1386
Number of shifts mapped to atoms	1386
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	790	780	779	11±4
All	All	14220	14040	14022	195

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:200:VAL:C	1:A:201:LYS:N	1.45	1.68	18	1
1:A:200:VAL:CA	1:A:201:LYS:N	1.43	1.81	18	1
1:A:200:VAL:HA	1:A:201:LYS:N	1.30	1.33	18	1
1:A:200:VAL:C	1:A:201:LYS:CA	1.13	2.17	18	1
1:A:200:VAL:HA	1:A:201:LYS:H	0.90	1.14	18	1
1:A:200:VAL:C	1:A:201:LYS:HA	0.88	1.86	18	1
1:A:200:VAL:O	1:A:201:LYS:CA	0.84	2.24	18	1
1:A:200:VAL:O	1:A:201:LYS:HA	0.80	1.74	18	1
1:A:208:LEU:HD11	1:A:220:GLN:HB2	0.69	1.64	16	2
1:A:173:PHE:CG	1:A:189:ILE:HG22	0.69	2.23	14	1
1:A:156:ARG:NE	1:A:179:GLU:OE2	0.65	2.29	18	1
1:A:200:VAL:O	1:A:201:LYS:N	0.65	2.27	18	1
1:A:205:ILE:HD11	1:A:215:ILE:HG12	0.65	1.69	13	8
1:A:208:LEU:HG	1:A:214:TYR:CD1	0.61	2.30	6	10
1:A:232:SER:HA	1:A:242:LEU:O	0.58	1.97	3	6
1:A:208:LEU:HD13	1:A:210:ASN:HD21	0.56	1.60	16	1
1:A:193:ASP:OD1	1:A:196:LYS:HD3	0.56	2.01	13	6
1:A:176:ARG:O	1:A:185:TYR:HB2	0.55	2.01	11	12
1:A:170:ARG:HG2	1:A:192:TRP:CZ2	0.55	2.36	6	1
1:A:231:TYR:CE1	1:A:236:ALA:HB3	0.54	2.38	5	5
1:A:231:TYR:CE2	1:A:236:ALA:HB3	0.54	2.37	8	13
1:A:185:TYR:CZ	1:A:205:ILE:HG21	0.54	2.38	16	1
1:A:159:ALA:O	1:A:163:LEU:HB2	0.53	2.03	14	1
1:A:149:TRP:CH2	1:A:224:LEU:HB3	0.53	2.39	11	7
1:A:233:GLU:HG3	1:A:234:ARG:HG3	0.52	1.80	4	1
1:A:168:ASN:O	1:A:190:ARG:HD3	0.52	2.05	1	1
1:A:172:THR:HA	1:A:244:VAL:HG11	0.51	1.82	1	1
1:A:161:ARG:O	1:A:165:SER:HB2	0.51	2.06	13	8
1:A:193:ASP:OD1	1:A:196:LYS:HD2	0.50	2.06	3	1
1:A:230:HIS:O	1:A:233:GLU:HG2	0.49	2.06	4	1
1:A:149:TRP:CZ3	1:A:224:LEU:HB2	0.49	2.42	15	1
1:A:159:ALA:CB	1:A:174:LEU:HD21	0.49	2.38	18	1
1:A:172:THR:HA	1:A:244:VAL:HG22	0.48	1.86	3	1
1:A:155:GLY:O	1:A:159:ALA:HB3	0.48	2.09	6	7
1:A:158:ASP:O	1:A:161:ARG:HG2	0.48	2.09	2	3
1:A:182:LYS:O	1:A:182:LYS:HD2	0.47	2.10	10	1
1:A:191:ASP:HB3	1:A:201:LYS:CB	0.47	2.39	16	1
1:A:154:LEU:O	1:A:176:ARG:HB3	0.47	2.10	4	1
1:A:218:ARG:NE	1:A:218:ARG:HA	0.47	2.25	5	1
1:A:231:TYR:CB	1:A:242:LEU:HD23	0.46	2.41	17	1
1:A:241:ARG:HG3	1:A:243:VAL:HG13	0.46	1.87	8	3
1:A:207:LYS:HG2	1:A:213:TYR:CE1	0.46	2.46	14	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:192:TRP:CD1	1:A:197:GLY:HA2	0.46	2.45	18	1
1:A:178:SER:HB2	1:A:181:THR:O	0.46	2.10	18	1
1:A:169:PRO:HA	1:A:190:ARG:NH1	0.46	2.26	18	1
1:A:154:LEU:HB2	1:A:158:ASP:HB3	0.46	1.88	18	4
1:A:170:ARG:O	1:A:170:ARG:HD2	0.46	2.10	5	2
1:A:170:ARG:HD2	1:A:170:ARG:O	0.46	2.11	15	1
1:A:154:LEU:HB2	1:A:158:ASP:HB2	0.46	1.88	14	2
1:A:173:PHE:CD1	1:A:189:ILE:HG22	0.46	2.46	14	1
1:A:198:ASP:O	1:A:199:HIS:HB2	0.45	2.11	8	3
1:A:231:TYR:O	1:A:241:ARG:HB2	0.45	2.12	13	1
1:A:191:ASP:OD2	1:A:239:CYS:HB2	0.45	2.11	3	1
1:A:185:TYR:CZ	1:A:224:LEU:HD11	0.45	2.47	16	1
1:A:223:THR:OG1	1:A:226:GLN:HG2	0.45	2.12	2	6
1:A:158:ASP:HA	1:A:161:ARG:HG2	0.45	1.88	4	1
1:A:171:GLY:C	1:A:244:VAL:HB	0.45	2.31	6	2
1:A:175:ILE:HG21	1:A:224:LEU:HD21	0.45	1.89	1	1
1:A:164:LEU:CD2	1:A:199:HIS:HB2	0.44	2.42	18	1
1:A:229:GLN:O	1:A:233:GLU:HG3	0.44	2.12	12	1
1:A:171:GLY:HA2	1:A:241:ARG:O	0.44	2.12	10	3
1:A:189:ILE:HD11	1:A:203:TYR:CE1	0.44	2.47	3	1
1:A:203:TYR:CD2	1:A:238:LEU:HD22	0.44	2.47	11	1
1:A:192:TRP:HE3	1:A:192:TRP:O	0.44	1.95	10	3
1:A:155:GLY:HA3	1:A:176:ARG:CB	0.44	2.42	5	3
1:A:173:PHE:HA	1:A:188:SER:O	0.43	2.13	3	1
1:A:192:TRP:O	1:A:192:TRP:HE3	0.43	1.97	15	2
1:A:191:ASP:HB3	1:A:201:LYS:HB2	0.43	1.91	16	1
1:A:164:LEU:HD22	1:A:198:ASP:OD1	0.43	2.14	4	1
1:A:168:ASN:ND2	1:A:172:THR:HG21	0.43	2.29	6	2
1:A:185:TYR:HE2	1:A:213:TYR:CD1	0.43	2.31	2	1
1:A:173:PHE:CE1	1:A:242:LEU:HB3	0.43	2.49	13	2
1:A:205:ILE:CG1	1:A:215:ILE:HG12	0.43	2.44	3	1
1:A:192:TRP:CG	1:A:197:GLY:HA2	0.43	2.48	9	1
1:A:185:TYR:CE2	1:A:224:LEU:HD11	0.42	2.48	16	1
1:A:185:TYR:CZ	1:A:224:LEU:HD21	0.42	2.49	5	1
1:A:231:TYR:HB3	1:A:242:LEU:HD23	0.42	1.90	17	1
1:A:191:ASP:OD2	1:A:201:LYS:HD2	0.42	2.14	1	1
1:A:149:TRP:CZ3	1:A:224:LEU:HB3	0.42	2.50	5	1
1:A:170:ARG:HB2	1:A:240:CYS:SG	0.42	2.55	17	1
1:A:198:ASP:O	1:A:200:VAL:N	0.42	2.53	18	1
1:A:154:LEU:HB2	1:A:158:ASP:CG	0.41	2.35	9	2
1:A:185:TYR:O	1:A:205:ILE:HG22	0.41	2.15	11	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:153:LYS:HE3	1:A:177:GLU:OE1	0.41	2.16	15	1
1:A:208:LEU:HD11	1:A:220:GLN:HB3	0.41	1.93	1	1
1:A:182:LYS:HD2	1:A:182:LYS:O	0.41	2.16	15	1
1:A:183:GLY:O	1:A:207:LYS:HD3	0.41	2.16	12	1
1:A:215:ILE:HG21	1:A:238:LEU:HD21	0.41	1.93	7	1
1:A:187:LEU:O	1:A:202:HIS:HA	0.41	2.16	2	2
1:A:186:SER:HA	1:A:203:TYR:O	0.40	2.17	5	1
1:A:191:ASP:HB3	1:A:201:LYS:CG	0.40	2.46	17	1
1:A:159:ALA:HB1	1:A:174:LEU:HD21	0.40	1.92	18	1
1:A:185:TYR:CZ	1:A:205:ILE:HD13	0.40	2.52	2	1
1:A:156:ARG:HG2	1:A:176:ARG:NH2	0.40	2.32	11	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/112 (85%)	86±2 (91±2%)	7±2 (8±2%)	2±1 (2±1%)	13	57
All	All	1710/2016 (85%)	1549 (91%)	133 (8%)	28 (2%)	13	57

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	155	GLY	18
1	A	198	ASP	7
1	A	199	HIS	3

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/96 (85%)	70±2 (85±3%)	12±2 (15±3%)	6	44
All	All	1476/1728 (85%)	1257 (85%)	219 (15%)	6	44

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	192	TRP	18
1	A	234	ARG	17
1	A	162	GLN	16
1	A	157	LYS	15
1	A	166	PHE	13
1	A	196	LYS	13
1	A	187	LEU	12
1	A	150	TYR	12
1	A	241	ARG	11
1	A	223	THR	11
1	A	170	ARG	10
1	A	176	ARG	9
1	A	214	TYR	7
1	A	205	ILE	7
1	A	224	LEU	6
1	A	182	LYS	5
1	A	158	ASP	4
1	A	244	VAL	4
1	A	193	ASP	4
1	A	179	GLU	4
1	A	229	GLN	3
1	A	190	ARG	2
1	A	206	ARG	2
1	A	163	LEU	2
1	A	204	LYS	2
1	A	243	VAL	1
1	A	198	ASP	1
1	A	200	VAL	1
1	A	153	LYS	1
1	A	242	LEU	1
1	A	199	HIS	1
1	A	177	GLU	1
1	A	168	ASN	1
1	A	161	ARG	1
1	A	210	ASN	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	18-A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
18	A	200:VAL	C	201:LYS	N	1.68
18	A	199:HIS	C	200:VAL	N	1.65

6 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 87% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

6.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1386
Number of shifts mapped to atoms	1386
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

6.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	112	-0.37 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	101	-0.05 ± 0.21	None needed (< 0.5 ppm)
$^{13}\text{C}'$	106	0.13 ± 0.19	None needed (< 0.5 ppm)
^{15}N	106	-0.65 ± 0.29	Should be applied

6.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 1094 atoms were assigned a chemical shift out of a possible 1250. 14 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	470/478 (98%)	189/191 (99%)	188/192 (98%)	93/95 (98%)
Sidechain	513/643 (80%)	320/379 (84%)	186/223 (83%)	7/41 (17%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	111/129 (86%)	57/68 (84%)	52/56 (93%)	2/5 (40%)
Overall	1094/1250 (88%)	566/638 (89%)	426/471 (90%)	102/141 (72%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 87%, i.e. 1140 atoms were assigned a chemical shift out of a possible 1303. 14 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	487/496 (98%)	196/198 (99%)	195/200 (98%)	96/98 (98%)
Sidechain	540/671 (80%)	338/397 (85%)	195/232 (84%)	7/42 (17%)
Aromatic	113/136 (83%)	58/72 (81%)	53/58 (91%)	2/6 (33%)
Overall	1140/1303 (87%)	592/667 (89%)	443/490 (90%)	105/146 (72%)

6.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

6.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

