



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 05:40 PM BST

PDB ID : 6MRJ
Title : Crystal structure of H.pylori NikR in complex with DNA
Authors : Pozharski, E.; de Serrano, V.
Deposited on : 2018-10-13
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>14%</div> <div>9%</div> </div> </div>
1	B	148	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>• •</div> </div> </div>
1	C	148	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>22%</div> <div>• •</div> </div> </div>
1	D	148	<div> <div></div> <div> <div></div> <div>75%</div> <div>18%</div> <div>7%</div> </div> </div>
2	L	36	<div> <div></div> <div> <div></div> <div>69%</div> <div>31%</div> </div> </div>
3	M	36	<div> <div></div> <div> <div></div> <div>72%</div> <div>28%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nickel-responsive regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	0	5	0
			1086	668	198	215	5			
1	B	142	Total	C	N	O	S	0	2	0
			1145	710	208	222	5			
1	C	142	Total	C	N	O	S	0	2	0
			1168	721	212	230	5			
1	D	138	Total	C	N	O	S	0	0	0
			1115	689	204	217	5			

- Molecule 2 is a DNA chain called DNA (36-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	36	Total	C	N	O	P	0	0	0
			739	359	127	218	35			

- Molecule 3 is a DNA chain called DNA (36-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	36	Total	C	N	O	P	0	0	0
			731	355	131	210	35			

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ni	0	0
			2	2		
4	A	2	Total	Ni	0	0
			2	2		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	12	Total O 12 12	0	0
6	B	10	Total O 10 10	0	0
6	C	5	Total O 5 5	0	0
6	D	3	Total O 3 3	0	0
6	L	3	Total O 3 3	0	0
6	M	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nickel-responsive regulator




- Molecule 1: Nickel-responsive regulator



- Molecule 1: Nickel-responsive regulator



- Molecule 1: Nickel-responsive regulator




- Molecule 2: DNA (36-MER)





- Molecule 3: DNA (36-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.37Å 94.18Å 204.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.47 – 2.80 39.47 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.47-2.80) 100.0 (39.47-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.81Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.207 , 0.241 0.238 , 0.273	Depositor DCC
R_{free} test set	1721 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	76.6	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6025	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1107	0.66	0/1496
1	B	0.49	0/1170	0.68	0/1579
1	C	0.46	0/1189	0.69	0/1601
1	D	0.48	0/1129	0.71	0/1520
2	L	0.99	0/828	1.01	0/1278
3	M	0.98	0/820	1.00	0/1262
All	All	0.65	0/6243	0.79	0/8736

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1086	0	1041	8	0
1	B	1145	0	1104	14	0
1	C	1168	0	1139	25	0
1	D	1115	0	1106	20	0
2	L	739	0	415	8	0
3	M	731	0	411	7	0
4	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	12	0	0	0	0
6	B	10	0	0	0	0
6	C	5	0	0	0	0
6	D	3	0	0	0	0
6	L	3	0	0	0	0
6	M	2	0	0	0	0
All	All	6025	0	5216	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99:HIS:HB3	1:C:97:THR:HG23	1.61	0.82
1:B:92:THR:HG23	1:B:119:GLU:HB3	1.66	0.78
1:C:144:PHE:HB2	1:D:134:LYS:HD3	1.79	0.64
1:C:38:SER:HA	1:D:15:VAL:HG13	1.80	0.64
1:C:137:LYS:HD3	1:D:139:THR:HG23	1.79	0.63
1:B:99:HIS:HB3	1:C:97:THR:CG2	2.31	0.57
2:L:31:DA:H2''	2:L:32:DT:H5''	1.87	0.56
1:C:82:ARG:HG2	1:C:130:LEU:HD21	1.89	0.55
2:L:14:DA:H2''	2:L:15:DA:H5''	1.90	0.53
1:B:82[B]:ARG:HE	1:B:86:ILE:HD11	1.74	0.53
2:L:22:DT:H2''	2:L:23:DA:C8	2.47	0.50
1:A:42[A]:ARG:HD3	1:B:17:LEU:HB3	1.93	0.50
1:B:95:LEU:HD12	1:B:112:ILE:HG22	1.93	0.50
1:C:25:LEU:HD11	1:C:40:LEU:HD23	1.93	0.50
1:C:112:ILE:HG21	1:D:100:ILE:HD13	1.94	0.49
1:C:76:GLN:HB3	1:C:79:LEU:HD12	1.94	0.49
1:D:11:ILE:HG13	1:D:12:ARG:N	2.27	0.48
1:D:54:TRP:CE3	1:D:121:GLN:HG2	2.49	0.48
3:M:32:DA:H2''	3:M:33:DT:H5''	1.96	0.48
1:C:34:TYR:CE2	1:D:129:GLY:HA2	2.50	0.47
1:C:37:ARG:HH12	3:M:27:DT:H3'	1.79	0.47
1:B:29:ILE:HA	1:B:34:TYR:HD2	1.80	0.47
1:D:54:TRP:HB2	1:D:117:SER:HB2	1.97	0.47
2:L:9:DA:C2	3:M:28:DA:C2	3.03	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:29:ILE:O	1:D:34:TYR:HB2	2.14	0.47
1:D:65:ILE:HB	1:D:143:SER:HB2	1.96	0.47
1:B:86:ILE:HG21	1:B:127:ILE:HG13	1.97	0.46
1:C:71:ILE:CD1	1:D:141:ALA:HB1	2.46	0.46
1:A:28:ARG:HA	1:A:31:LYS:HE2	1.98	0.46
1:C:12:ARG:HH11	3:M:26:DA:H62	1.64	0.46
1:C:15:VAL:O	1:D:12:ARG:HA	2.15	0.45
1:A:134:LYS:HD3	1:B:144:PHE:HB2	1.99	0.45
1:C:70:VAL:HG13	1:C:133:VAL:HG13	1.98	0.45
1:D:18:GLN:HB2	1:D:21:LEU:HD12	1.99	0.45
1:D:113:LEU:HD22	1:D:123:LEU:HD13	1.99	0.45
1:D:11:ILE:HG13	1:D:12:ARG:H	1.82	0.45
1:C:145:GLU:HB2	1:C:148:GLU:HG3	1.98	0.44
1:A:22:LEU:HD12	1:A:25:LEU:HD23	1.99	0.44
1:A:75:HIS:CE1	1:A:104:GLU:O	2.71	0.43
1:A:100:ILE:HD13	1:D:96:CYS:HB3	2.00	0.43
1:B:116:ASN:HD21	1:B:118:PHE:HB2	1.83	0.43
1:C:95:LEU:HD11	1:C:114:GLN:HG3	2.01	0.43
1:C:141:ALA:HB1	1:D:71:ILE:HD11	2.01	0.43
1:C:46:ARG:O	1:C:50:VAL:HG23	2.19	0.43
1:D:22:LEU:HD12	1:D:25:LEU:HD23	2.01	0.42
1:A:125:LEU:HD22	1:B:32:ASN:HB3	2.01	0.42
1:B:42:ARG:HH22	2:L:24:DG:P	2.43	0.42
1:A:110:THR:HG21	1:B:112:ILE:HD11	2.02	0.42
1:C:95:LEU:HD12	1:C:112:ILE:HG22	2.02	0.41
1:C:39:GLU:HG2	1:C:42:ARG:HH21	1.85	0.41
1:C:58:ASN:HD22	1:C:60:ASN:ND2	2.18	0.41
1:C:112:ILE:HG21	1:D:100:ILE:CD1	2.51	0.41
3:M:16:DT:H2"	3:M:17:DT:H5"	2.02	0.41
1:C:71:ILE:HD11	1:D:141:ALA:CB	2.51	0.41
3:M:18:DC:H2"	3:M:19:DA:C8	2.55	0.41
2:L:5:DT:H2'	2:L:6:DA:H8	1.86	0.40
1:C:99:HIS:CD2	1:C:101:HIS:CE1	3.09	0.40
2:L:23:DA:C2	3:M:14:DA:C2	3.10	0.40
2:L:30:DT:H2"	2:L:31:DA:H8	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/148 (93%)	128 (93%)	8 (6%)	1 (1%)	22	53
1	B	142/148 (96%)	137 (96%)	5 (4%)	0	100	100
1	C	142/148 (96%)	138 (97%)	4 (3%)	0	100	100
1	D	136/148 (92%)	129 (95%)	7 (5%)	0	100	100
All	All	557/592 (94%)	532 (96%)	24 (4%)	1 (0%)	47	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/137 (88%)	111 (92%)	10 (8%)	11	32
1	B	127/137 (93%)	123 (97%)	4 (3%)	40	74
1	C	132/137 (96%)	124 (94%)	8 (6%)	18	48
1	D	127/137 (93%)	123 (97%)	4 (3%)	40	74
All	All	507/548 (92%)	481 (95%)	26 (5%)	24	55

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	12	ARG
1	A	17	LEU
1	A	35	SER
1	A	43	ASP
1	A	52	ASP
1	A	90[A]	SER
1	A	90[B]	SER
1	A	113	LEU
1	A	121	GLN
1	B	17	LEU
1	B	19	GLN
1	B	85	ASP
1	B	142	SER
1	C	8	ASP
1	C	17	LEU
1	C	34	TYR
1	C	60	ASN
1	C	79	LEU
1	C	119	GLU
1	C	123	LEU
1	C	147	ASN
1	D	8	ASP
1	D	17	LEU
1	D	30	ILE
1	D	114	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	116	ASN
1	C	58	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	134/148 (90%)	0.12	3 (2%) 62 52	55, 98, 141, 157	0
1	B	142/148 (95%)	-0.24	1 (0%) 87 84	56, 79, 122, 139	0
1	C	142/148 (95%)	0.00	5 (3%) 44 34	47, 91, 134, 145	0
1	D	138/148 (93%)	0.06	0 100 100	50, 89, 131, 154	0
2	L	36/36 (100%)	-0.52	0 100 100	105, 124, 140, 145	0
3	M	36/36 (100%)	-0.29	0 100 100	100, 133, 145, 147	0
All	All	628/664 (94%)	-0.06	9 (1%) 75 70	47, 93, 139, 157	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	59	PRO	3.9
1	A	54	TRP	3.1
1	C	7	ASP	2.9
1	A	58	ASN	2.7
1	A	59	PRO	2.6
1	C	10	ILE	2.5
1	B	15	VAL	2.1
1	C	146[A]	TYR	2.1
1	C	133	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	C	201	1/1	0.92	0.13	96,96,96,96	0
5	MG	A	202	1/1	0.95	0.19	116,116,116,116	0
4	NI	A	201	1/1	0.99	0.24	63,63,63,63	0
4	NI	B	201	1/1	0.99	0.12	88,88,88,88	0
4	NI	B	202	1/1	0.99	0.16	62,62,62,62	0
4	NI	A	203	1/1	0.99	0.20	86,86,86,86	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 202:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NI A 201:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NI B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.