



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 02:06 am BST

PDB ID : 6MSV
Title : Structure of the 6th type III domain from human fibronectin
Authors : Loa, S.; Mou, T.C.; Sprang, S.R.; Briknarova, K.
Deposited on : 2018-10-18
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	92	<div> <div>68%</div> <div>25%</div> <div>7%</div> </div>
1	B	92	<div> <div>70%</div> <div>21%</div> <div>10%</div> </div>
1	C	92	<div> <div>60%</div> <div>33%</div> <div>8%</div> </div>
1	D	92	<div> <div>2%</div> <div>73%</div> <div>24%</div> <div>..</div> </div>
1	E	92	<div> <div>5%</div> <div>54%</div> <div>36%</div> <div>•</div> <div>9%</div> </div>
1	F	92	<div> <div>2%</div> <div>79%</div> <div>12%</div> <div>9%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	92	<div> <div>2%</div> <div>61%</div> <div>29%</div> <div>10%</div> </div>
1	H	92	<div> <div>1%</div> <div>78%</div> <div>18%</div> <div>3%</div> </div>
1	I	92	<div> <div>2%</div> <div>54%</div> <div>33%</div> <div>12%</div> </div>
1	J	92	<div> <div>4%</div> <div>51%</div> <div>38%</div> <div>10%</div> </div>
1	K	92	<div> <div>3%</div> <div>53%</div> <div>37%</div> <div>9%</div> </div>
1	L	92	<div> <div>12%</div> <div>57%</div> <div>33%</div> <div>10%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibronectin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	86	Total	C	N	O	0	0	0
			645	407	109	129			
1	B	83	Total	C	N	O	0	0	0
			629	399	106	124			
1	C	85	Total	C	N	O	0	0	0
			641	405	108	128			
1	D	90	Total	C	N	O	0	0	0
			673	425	114	134			
1	E	84	Total	C	N	O	0	0	0
			635	402	107	126			
1	F	84	Total	C	N	O	0	0	0
			635	402	107	126			
1	G	83	Total	C	N	O	0	0	0
			629	399	106	124			
1	H	89	Total	C	N	O	0	0	0
			669	423	113	133			
1	I	81	Total	C	N	O	0	0	0
			618	393	103	122			
1	J	83	Total	C	N	O	0	0	0
			629	399	106	124			
1	K	84	Total	C	N	O	0	0	0
			635	402	107	126			
1	L	83	Total	C	N	O	0	0	0
			629	399	106	124			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1082	GLY	-	expression tag	UNP P02751
A	1083	SER	-	expression tag	UNP P02751
A	1084	GLY	-	expression tag	UNP P02751
B	1082	GLY	-	expression tag	UNP P02751
B	1083	SER	-	expression tag	UNP P02751

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1084	GLY	-	expression tag	UNP P02751
C	1082	GLY	-	expression tag	UNP P02751
C	1083	SER	-	expression tag	UNP P02751
C	1084	GLY	-	expression tag	UNP P02751
D	1082	GLY	-	expression tag	UNP P02751
D	1083	SER	-	expression tag	UNP P02751
D	1084	GLY	-	expression tag	UNP P02751
E	1082	GLY	-	expression tag	UNP P02751
E	1083	SER	-	expression tag	UNP P02751
E	1084	GLY	-	expression tag	UNP P02751
F	1082	GLY	-	expression tag	UNP P02751
F	1083	SER	-	expression tag	UNP P02751
F	1084	GLY	-	expression tag	UNP P02751
G	1082	GLY	-	expression tag	UNP P02751
G	1083	SER	-	expression tag	UNP P02751
G	1084	GLY	-	expression tag	UNP P02751
H	1082	GLY	-	expression tag	UNP P02751
H	1083	SER	-	expression tag	UNP P02751
H	1084	GLY	-	expression tag	UNP P02751
I	1082	GLY	-	expression tag	UNP P02751
I	1083	SER	-	expression tag	UNP P02751
I	1084	GLY	-	expression tag	UNP P02751
J	1082	GLY	-	expression tag	UNP P02751
J	1083	SER	-	expression tag	UNP P02751
J	1084	GLY	-	expression tag	UNP P02751
K	1082	GLY	-	expression tag	UNP P02751
K	1083	SER	-	expression tag	UNP P02751
K	1084	GLY	-	expression tag	UNP P02751
L	1082	GLY	-	expression tag	UNP P02751
L	1083	SER	-	expression tag	UNP P02751
L	1084	GLY	-	expression tag	UNP P02751

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		
3	B	18	Total	O	0	0
			18	18		
3	C	16	Total	O	0	0
			16	16		
3	D	8	Total	O	0	0
			8	8		
3	E	12	Total	O	0	0
			12	12		
3	F	17	Total	O	0	0
			17	17		
3	G	12	Total	O	0	0
			12	12		
3	H	14	Total	O	0	0
			14	14		
3	I	14	Total	O	0	0
			14	14		
3	J	11	Total	O	0	0
			11	11		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	K	11	Total	O	0	0
			11	11		
3	L	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibronectin

Chain A: 



• Molecule 1: Fibronectin

Chain B: 





• Molecule 1: Fibronectin

Chain C: 




• Molecule 1: Fibronectin

Chain D: 





• Molecule 1: Fibronectin

Chain E: 




- Molecule 1: Fibronectin



- Molecule 1: Fibronectin




- Molecule 1: Fibronectin



- Molecule 1: Fibronectin




- Molecule 1: Fibronectin




- Molecule 1: Fibronectin





● Molecule 1: Fibronectin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.52Å 79.46Å 80.89Å 112.33° 95.88° 94.48°	Depositor
Resolution (Å)	26.64 – 2.40 26.64 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.8 (26.64-2.40) 84.7 (26.64-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.29Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.235 , 0.295 0.235 , 0.295	Depositor DCC
R_{free} test set	1992 reflections (4.25%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.076 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7827	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/659	0.47	0/905
1	B	0.24	0/643	0.47	0/884
1	C	0.24	0/655	0.47	0/900
1	D	0.25	0/688	0.50	0/945
1	E	0.26	0/649	0.54	0/892
1	F	0.44	0/649	0.59	0/892
1	G	0.35	0/643	0.53	0/884
1	H	0.25	0/684	0.50	0/940
1	I	0.24	0/631	0.51	0/867
1	J	0.25	0/643	0.50	0/884
1	K	0.25	0/649	0.48	0/892
1	L	0.29	0/643	0.58	0/884
All	All	0.28	0/7836	0.51	0/10769

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	645	0	648	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	629	0	635	17	0
1	C	641	0	645	23	0
1	D	673	0	677	13	0
1	E	635	0	640	33	0
1	F	635	0	640	5	0
1	G	629	0	635	19	0
1	H	669	0	674	13	0
1	I	618	0	625	23	0
1	J	629	0	635	32	0
1	K	635	0	640	20	0
1	L	629	0	635	29	0
2	B	6	0	8	2	0
2	F	6	0	8	0	0
3	A	13	0	0	0	0
3	B	18	0	0	0	0
3	C	16	0	0	1	0
3	D	8	0	0	0	0
3	E	12	0	0	0	0
3	F	17	0	0	0	0
3	G	12	0	0	0	0
3	H	14	0	0	1	0
3	I	14	0	0	0	0
3	J	11	0	0	0	0
3	K	11	0	0	0	0
3	L	2	0	0	1	0
All	All	7827	0	7745	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:1157:ARG:HB3	1:G:1162:ARG:HD2	1.49	0.92
1:L:1120:ARG:HB3	1:L:1128:PRO:HB3	1.50	0.91
1:I:1157:ARG:HD2	1:I:1162:ARG:HE	1.36	0.88
1:G:1116:LYS:CE	1:G:1130:GLU:OE1	2.21	0.88
1:A:1168:ASN:HD22	1:J:1170:VAL:HA	1.39	0.88

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	84/92 (91%)	83 (99%)	1 (1%)	0	100	100
1	B	81/92 (88%)	81 (100%)	0	0	100	100
1	C	83/92 (90%)	79 (95%)	4 (5%)	0	100	100
1	D	88/92 (96%)	83 (94%)	5 (6%)	0	100	100
1	E	82/92 (89%)	78 (95%)	4 (5%)	0	100	100
1	F	82/92 (89%)	79 (96%)	3 (4%)	0	100	100
1	G	81/92 (88%)	79 (98%)	2 (2%)	0	100	100
1	H	87/92 (95%)	83 (95%)	4 (5%)	0	100	100
1	I	77/92 (84%)	75 (97%)	2 (3%)	0	100	100
1	J	81/92 (88%)	77 (95%)	3 (4%)	1 (1%)	13	19
1	K	82/92 (89%)	78 (95%)	3 (4%)	1 (1%)	13	19
1	L	81/92 (88%)	79 (98%)	1 (1%)	1 (1%)	13	19
All	All	989/1104 (90%)	954 (96%)	32 (3%)	3 (0%)	41	55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	1125	GLY
1	J	1113	ILE
1	L	1128	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	74/78 (95%)	73 (99%)	1 (1%)	67	82
1	B	72/78 (92%)	71 (99%)	1 (1%)	67	82
1	C	74/78 (95%)	74 (100%)	0	100	100
1	D	77/78 (99%)	75 (97%)	2 (3%)	46	66
1	E	73/78 (94%)	72 (99%)	1 (1%)	67	82
1	F	73/78 (94%)	71 (97%)	2 (3%)	44	65
1	G	72/78 (92%)	71 (99%)	1 (1%)	67	82
1	H	77/78 (99%)	77 (100%)	0	100	100
1	I	72/78 (92%)	69 (96%)	3 (4%)	30	47
1	J	72/78 (92%)	71 (99%)	1 (1%)	67	82
1	K	73/78 (94%)	70 (96%)	3 (4%)	30	48
1	L	72/78 (92%)	70 (97%)	2 (3%)	43	63
All	All	881/936 (94%)	864 (98%)	17 (2%)	57	75

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	1137	SER
1	I	1112	ARG
1	K	1156	LEU
1	F	1157	ARG
1	K	1157	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1160	GLN
1	A	1168	ASN
1	F	1160	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	F	1201	-	5,5,5	0.89	0	5,5,5	1.00	0
2	GOL	B	1201	-	5,5,5	0.98	0	5,5,5	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	F	1201	-	-	2/4/4/4	-
2	GOL	B	1201	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	1201	GOL	C1-C2-C3-O3
2	B	1201	GOL	O1-C1-C2-C3
2	F	1201	GOL	O2-C2-C3-O3
2	B	1201	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1201	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	86/92 (93%)	0.12	0 100 100	31, 49, 87, 92	0
1	B	83/92 (90%)	-0.04	0 100 100	33, 50, 73, 82	0
1	C	85/92 (92%)	0.13	0 100 100	35, 52, 80, 92	0
1	D	90/92 (97%)	0.24	2 (2%) 62 60	37, 59, 87, 106	0
1	E	84/92 (91%)	0.42	5 (5%) 21 20	38, 53, 91, 102	0
1	F	84/92 (91%)	-0.00	2 (2%) 59 57	34, 49, 81, 93	0
1	G	83/92 (90%)	-0.06	2 (2%) 59 57	33, 46, 77, 91	0
1	H	89/92 (96%)	0.17	1 (1%) 80 79	32, 51, 83, 102	0
1	I	81/92 (88%)	0.18	2 (2%) 57 55	39, 51, 88, 102	0
1	J	83/92 (90%)	0.44	4 (4%) 30 29	34, 58, 108, 122	0
1	K	84/92 (91%)	0.14	3 (3%) 42 42	34, 51, 81, 97	0
1	L	83/92 (90%)	0.64	11 (13%) 3 3	38, 61, 102, 108	0
All	All	1015/1104 (91%)	0.20	32 (3%) 47 46	31, 53, 90, 122	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	1127	ALA	6.2
1	F	1111	PRO	5.3
1	J	1157	ARG	4.6
1	L	1091	ILE	4.6
1	E	1155	VAL	4.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	F	1201	6/6	0.92	0.11	55,57,64,64	0
2	GOL	B	1201	6/6	0.94	0.12	50,59,63,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.