



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 11, 2022 – 10:13 AM EST

PDB ID : 7MSQ  
Title : Complex between the Fab arm of AB-3467 and the SARS-CoV-2 receptor binding domain (RBD)  
Authors : Langley, D.B.; Christ, D.  
Deposited on : 2021-05-12  
Resolution : 2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.25  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.25


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	<div> <div>13%</div> <div>81%</div> <div>13%</div> <div>.</div> </div>
1	B	204	<div> <div>13%</div> <div>80%</div> <div>11%</div> <div>7%</div> </div>
2	D	238	<div> <div>2%</div> <div>82%</div> <div>11%</div> <div>7%</div> </div>
2	H	238	<div> <div>%</div> <div>85%</div> <div>7%</div> <div>7%</div> </div>
3	E	214	<div> <div>%</div> <div>90%</div> <div>9%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	L	214	<div><div>%</div><div><div></div><div>91%</div><div>8%</div></div></div>
4	C	5	<div><div><div></div><div>20%</div><div>80%</div></div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1541	988	257	288	8			
1	B	189	Total	C	N	O	S	0	0	0
			1492	955	248	281	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	529	GLY	-	expression tag	UNP P0DTC2
A	530	SER	-	expression tag	UNP P0DTC2
A	531	HIS	-	expression tag	UNP P0DTC2
A	532	HIS	-	expression tag	UNP P0DTC2
A	533	HIS	-	expression tag	UNP P0DTC2
A	534	HIS	-	expression tag	UNP P0DTC2
A	535	HIS	-	expression tag	UNP P0DTC2
A	536	HIS	-	expression tag	UNP P0DTC2
B	529	GLY	-	expression tag	UNP P0DTC2
B	530	SER	-	expression tag	UNP P0DTC2
B	531	HIS	-	expression tag	UNP P0DTC2
B	532	HIS	-	expression tag	UNP P0DTC2
B	533	HIS	-	expression tag	UNP P0DTC2
B	534	HIS	-	expression tag	UNP P0DTC2
B	535	HIS	-	expression tag	UNP P0DTC2
B	536	HIS	-	expression tag	UNP P0DTC2


- Molecule 2 is a protein called AB-3467 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	221	Total	C	N	O	S	0	3	0
			1698	1082	279	331	6			
2	H	221	Total	C	N	O	S	0	4	0
			1702	1084	280	332	6			

- Molecule 3 is a protein called AB-3467 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	213	Total	C	N	O	S	0	0	0
			1637	1027	274	332	4			
3	L	214	Total	C	N	O	S	0	0	0
			1640	1027	274	334	5			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	5	Total	C	N	O	0	0	0
			60	34	2	24			

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		
5	D	2	Total	Cl	0	0
			2	2		
5	E	2	Total	Cl	0	0
			2	2		
5	H	1	Total	Cl	0	0
			1	1		

- Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	S	0	0
			3	1	1	1		
6	E	1	Total	C	N	S	0	0
			3	1	1	1		
6	L	1	Total	C	N	S	0	0
			3	1	1	1		
6	L	1	Total	C	N	S	0	0
			3	1	1	1		
6	L	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	10	Total	O	0	0
			10	10		
8	B	18	Total	O	0	0
			18	18		
8	D	32	Total	O	0	0
			32	32		
8	E	35	Total	O	0	0
			35	35		
8	H	38	Total	O	0	0
			38	38		
8	L	31	Total	O	0	0
			31	31		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 1: Spike protein S1



- Molecule 2: AB-3467 Fab Heavy Chain




- Molecule 2: AB-3467 Fab Heavy Chain






● Molecule 3: AB-3467 Fab Light Chain



● Molecule 3: AB-3467 Fab Light Chain



● Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.76Å 115.49Å 240.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.00 – 2.29 48.00 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.00-2.29) 99.8 (48.00-2.29)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.194 , 0.245 0.198 , 0.246	Depositor DCC
$R_{free}$ test set	3368 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 34.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.023 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9962	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SCN, CL, BMA, FUC, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.78	0/1585	0.90	0/2158
1	B	0.74	0/1533	0.92	0/2084
2	D	0.77	1/1744 (0.1%)	0.96	2/2382 (0.1%)
2	H	0.76	1/1748 (0.1%)	0.94	0/2388
3	E	0.79	0/1673	0.94	0/2273
3	L	0.78	1/1676 (0.1%)	0.90	0/2277
All	All	0.77	3/9959 (0.0%)	0.93	2/13562 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	162	GLU	CD-OE1	6.51	1.32	1.25
2	D	16	GLU	CD-OE1	5.44	1.31	1.25
3	L	123	GLU	CD-OE2	5.10	1.31	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	66	ARG	NE-CZ-NH2	-7.75	116.42	120.30
2	D	66	ARG	NE-CZ-NH1	5.50	123.05	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1541	0	1454	14	0
1	B	1492	0	1395	17	0
2	D	1698	0	1645	11	0
2	H	1702	0	1644	11	0
3	E	1637	0	1590	11	0
3	L	1640	0	1584	10	0
4	C	60	0	52	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	D	2	0	0	0	0
5	E	2	0	0	0	0
5	H	1	0	0	0	0
6	B	3	0	0	0	0
6	E	3	0	0	0	0
6	L	9	0	0	0	0
7	D	6	0	8	0	0
8	A	10	0	0	0	0
8	B	18	0	0	0	0
8	D	32	0	0	0	0
8	E	35	0	0	0	0
8	H	38	0	0	2	0
8	L	31	0	0	0	0
All	All	9962	0	9372	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:213[A]:ASN:OD1	8:H:401:HOH:O	1.85	0.94
1:B:365:TYR:HB3	1:B:387:LEU:HD13	1.60	0.82
1:B:371:SER:O	1:B:373:SER:N	2.29	0.65
3:L:11:LEU:CD2	3:L:104:LEU:HD13	2.27	0.63
3:L:11:LEU:HD22	3:L:104:LEU:HD13	1.82	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/204 (95%)	174 (90%)	17 (9%)	2 (1%)	15	17
1	B	185/204 (91%)	174 (94%)	9 (5%)	2 (1%)	14	15
2	D	220/238 (92%)	210 (96%)	7 (3%)	3 (1%)	11	11
2	H	221/238 (93%)	213 (96%)	5 (2%)	3 (1%)	11	11
3	E	211/214 (99%)	204 (97%)	7 (3%)	0	100	100
3	L	212/214 (99%)	202 (95%)	10 (5%)	0	100	100
All	All	1242/1312 (95%)	1177 (95%)	55 (4%)	10 (1%)	25	23

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	372	ALA
1	A	519	HIS
1	B	387	LEU
2	D	65	SER
1	A	483	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/176 (95%)	160 (96%)	7 (4%)	30	42
1	B	161/176 (92%)	153 (95%)	8 (5%)	24	34
2	D	190/207 (92%)	183 (96%)	7 (4%)	34	48

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	190/207 (92%)	185 (97%)	5 (3%)	46	63
3	E	186/187 (100%)	184 (99%)	2 (1%)	73	86
3	L	186/187 (100%)	184 (99%)	2 (1%)	73	86
All	All	1080/1140 (95%)	1049 (97%)	31 (3%)	42	58

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	516	GLU
2	H	175	SER
2	D	122	MET
3	L	11	LEU
2	H	50	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	C	1	1,4	14,14,15	0.89	1 (7%)	17,19,21	1.10	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	C	2	4	14,14,15	0.62	0	17,19,21	1.35	3 (17%)
4	BMA	C	3	4	11,11,12	0.61	0	15,15,17	1.70	4 (26%)
4	MAN	C	4	4	11,11,12	0.43	0	15,15,17	1.09	2 (13%)
4	FUC	C	5	4	10,10,11	0.52	0	14,14,16	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1
4	BMA	C	3	4	-	0/2/19/22	0/1/1/1
4	MAN	C	4	4	-	0/2/19/22	0/1/1/1
4	FUC	C	5	4	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1	NAG	O5-C1	-2.15	1.40	1.43

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	3	BMA	C1-O5-C5	3.88	117.45	112.19
4	C	3	BMA	O2-C2-C1	-3.48	102.04	109.15
4	C	3	BMA	O5-C1-C2	2.56	114.72	110.77
4	C	2	NAG	O5-C5-C6	2.44	111.04	107.20
4	C	3	BMA	O5-C5-C6	2.41	110.98	107.20


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 7 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	SCN	L	301	-	1,2,2	0.74	0	0,1,1	-	-
7	GOL	D	303	-	5,5,5	0.21	0	5,5,5	0.56	0
6	SCN	L	303	-	1,2,2	0.78	0	0,1,1	-	-
6	SCN	L	302	-	1,2,2	0.91	0	0,1,1	-	-
6	SCN	E	303	-	1,2,2	0.53	0	0,1,1	-	-
6	SCN	B	602	-	1,2,2	0.32	0	0,1,1	-	-



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	D	303	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	D	303	GOL	O1-C1-C2-O2
7	D	303	GOL	O1-C1-C2-C3
7	D	303	GOL	C1-C2-C3-O3
7	D	303	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	195/204 (95%)	0.68	27 (13%) 2 4	31, 63, 119, 146	0
1	B	189/204 (92%)	0.74	27 (14%) 2 3	35, 63, 106, 133	0
2	D	221/238 (92%)	0.04	5 (2%) 60 67	34, 46, 71, 101	0
2	H	221/238 (92%)	-0.01	3 (1%) 75 80	33, 47, 66, 100	0
3	E	213/214 (99%)	-0.12	2 (0%) 84 88	31, 46, 68, 96	0
3	L	214/214 (100%)	-0.14	2 (0%) 84 88	36, 48, 66, 124	0
All	All	1253/1312 (95%)	0.18	66 (5%) 26 33	31, 49, 94, 146	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	486	PHE	6.6
1	B	476	GLY	6.0
3	L	214	CYS	5.4
1	A	477	SER	5.3
1	B	477	SER	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FUC	C	5	10/11	0.83	0.21	112,117,121,121	0
4	BMA	C	3	11/12	0.84	0.14	74,76,80,83	0
4	NAG	C	1	14/15	0.87	0.17	56,70,82,90	0
4	MAN	C	4	11/12	0.90	0.12	77,81,85,86	0
4	NAG	C	2	14/15	0.93	0.17	70,75,79,81	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CL	D	302	1/1	0.56	0.17	89,89,89,89	0
7	GOL	D	303	6/6	0.75	0.17	44,55,63,67	0
6	SCN	L	303	3/3	0.82	0.28	46,46,63,75	0
5	CL	A	601	1/1	0.82	0.09	84,84,84,84	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CL	B	601	1/1	0.89	0.23	77,77,77,77	0
6	SCN	L	301	3/3	0.90	0.13	60,60,69,88	0
6	SCN	L	302	3/3	0.94	0.15	55,55,58,59	0
5	CL	E	302	1/1	0.94	0.11	64,64,64,64	0
5	CL	D	301	1/1	0.94	0.09	63,63,63,63	0
6	SCN	E	303	3/3	0.95	0.10	47,47,54,60	0
6	SCN	B	602	3/3	0.96	0.09	43,43,45,53	0
5	CL	E	301	1/1	0.97	0.10	59,59,59,59	0
5	CL	H	301	1/1	0.99	0.11	63,63,63,63	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.