



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 11:41 am BST

PDB ID : 4MT9
Title : Co-crystal structure of tankyrase 1 with compound 49
Authors : Huang, X.
Deposited on : 2013-09-19
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	 77% 19% • •
1	B	217	 2% 64% 25% • 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3776 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1688	1063	308	305	12			
1	B	202	Total	C	N	O	S	0	0	0
			1613	1012	295	294	12			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1315	HIS	-	EXPRESSION TAG	UNP O95271
A	1316	HIS	-	EXPRESSION TAG	UNP O95271
A	1317	HIS	-	EXPRESSION TAG	UNP O95271
A	1318	HIS	-	EXPRESSION TAG	UNP O95271
A	1319	HIS	-	EXPRESSION TAG	UNP O95271
A	1320	HIS	-	EXPRESSION TAG	UNP O95271
B	1315	HIS	-	EXPRESSION TAG	UNP O95271
B	1316	HIS	-	EXPRESSION TAG	UNP O95271
B	1317	HIS	-	EXPRESSION TAG	UNP O95271
B	1318	HIS	-	EXPRESSION TAG	UNP O95271
B	1319	HIS	-	EXPRESSION TAG	UNP O95271
B	1320	HIS	-	EXPRESSION TAG	UNP O95271

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is N-[trans-4-(4-cyanophenoxy)cyclohexyl]-3-[(4-oxo-3,4-dihydroquinazolin-2-yl)sulfanyl]propanamide (three-letter code: 2D6) (formula: C₂₄H₂₄N₄O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			32	24	4	3	1		
3	B	1	Total	C	N	O	S	0	0
			32	24	4	3	1		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total	O	0	0
			227	227		
4	B	182	Total	O	0	0
			182	182		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tankyrase-1

Chain A: 



• Molecule 1: Tankyrase-1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.01Å 76.78Å 148.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 41.63 – 1.88	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.00) 99.1 (41.63-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.88Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.212 , 0.258 0.218 , 0.236	Depositor DCC
R_{free} test set	1979 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3776	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 2D6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1731	0.67	1/2327 (0.0%)
1	B	0.59	1/1652 (0.1%)	0.67	0/2221
All	All	0.57	1/3383 (0.0%)	0.67	1/4548 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1160	GLU	CG-CD	5.15	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1251	PHE	N-CA-C	-5.03	97.41	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1622	46	0
1	B	1613	0	1527	63	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	32	0	24	3	0
3	B	32	0	24	0	0
4	A	227	0	0	22	0
4	B	182	0	0	17	0
All	All	3776	0	3197	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1114:GLU:CG	4:A:1537:HOH:O	2.19	0.88
1:B:1176:HIS:HD2	4:B:1551:HOH:O	1.55	0.87
1:B:1144:ARG:HD2	1:B:1311:MET:SD	2.17	0.84
1:A:1175:ASN:OD1	4:A:1661:HOH:O	1.96	0.82
1:B:1131:GLU:OE1	4:B:1682:HOH:O	1.98	0.80
1:B:1290:ALA:N	4:B:1599:HOH:O	2.15	0.79
1:B:1268:MET:HE1	1:B:1278:VAL:HG21	1.67	0.77
1:A:1201:HIS:HD2	4:A:1655:HOH:O	1.69	0.76
1:A:1266:MET:CE	4:A:1702:HOH:O	2.33	0.75
1:B:1131:GLU:CD	4:B:1682:HOH:O	2.27	0.73
1:A:1172:GLU:OE2	4:A:1662:HOH:O	2.07	0.73
1:A:1298:GLU:CG	4:A:1522:HOH:O	2.37	0.72
1:B:1115:ASP:OD1	4:B:1627:HOH:O	2.07	0.71
1:A:1253:ARG:HG3	1:A:1303:GLU:HG2	1.73	0.70
1:B:1205:GLY:O	4:B:1620:HOH:O	2.09	0.69
1:B:1197:PHE:HB3	1:B:1212:ILE:HD13	1.74	0.69
3:A:1402:2D6:H23	4:A:1608:HOH:O	1.92	0.68
1:A:1298:GLU:H	1:A:1298:GLU:CD	1.96	0.67
1:A:1144:ARG:HG2	1:A:1145:TYR:N	2.07	0.67
1:B:1176:HIS:CD2	4:B:1551:HOH:O	2.37	0.66
1:A:1208:PHE:CE2	1:A:1228:ILE:HD12	2.31	0.66
1:A:1112:ALA:HB1	1:A:1114:GLU:OE2	1.97	0.65
1:A:1114:GLU:CD	4:A:1537:HOH:O	2.37	0.62
1:B:1268:MET:CE	1:B:1278:VAL:HG21	2.31	0.61
1:B:1144:ARG:HD3	1:B:1145:TYR:O	2.00	0.61
1:A:1130:ARG:NE	4:A:1708:HOH:O	2.01	0.59
1:A:1131:GLU:OE1	1:A:1131:GLU:HA	1.99	0.59
1:B:1268:MET:HE1	1:B:1278:VAL:CG2	2.32	0.59
1:A:1281:ARG:HH11	1:A:1281:ARG:HG3	1.68	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1266:MET:O	1:B:1268:MET:HG3	2.05	0.57
1:A:1216:GLU:HG2	4:A:1547:HOH:O	2.05	0.57
1:A:1253:ARG:HG3	1:A:1303:GLU:CG	2.35	0.56
1:B:1208:PHE:HE2	1:B:1262:GLN:HB3	1.71	0.56
1:A:1281:ARG:NH1	1:A:1281:ARG:HG3	2.20	0.56
1:B:1291:GLU:HG3	4:B:1675:HOH:O	2.05	0.56
1:A:1208:PHE:CZ	1:A:1228:ILE:HD12	2.42	0.55
1:A:1114:GLU:HG2	4:A:1537:HOH:O	1.96	0.55
1:B:1128:THR:HB	1:B:1217:ASN:HA	1.88	0.55
1:A:1266:MET:HG3	1:A:1267:LYS:HG2	1.89	0.54
1:B:1122:GLU:HG3	1:B:1147:VAL:HG21	1.89	0.54
1:A:1128:THR:HB	1:A:1217:ASN:HA	1.88	0.54
1:A:1114:GLU:H	1:A:1114:GLU:CD	2.10	0.53
1:A:1263:PHE:CZ	1:B:1261:LEU:HG	2.44	0.53
1:A:1298:GLU:HG3	4:A:1522:HOH:O	2.04	0.53
1:B:1260:PHE:CE2	1:B:1262:GLN:HG2	2.44	0.52
1:B:1112:ALA:O	1:B:1115:ASP:HB3	2.09	0.52
1:B:1180:ARG:NH2	1:B:1292:TYR:OH	2.43	0.52
1:B:1156:LYS:O	1:B:1160:GLU:HG3	2.09	0.52
1:B:1132:HIS:C	1:B:1134:ASP:H	2.13	0.52
1:B:1252:CYS:HB3	1:B:1301:TYR:O	2.09	0.52
1:B:1117:GLU:CD	1:B:1152:LYS:HZ1	2.14	0.52
1:B:1180:ARG:NH1	1:B:1180:ARG:HG3	2.25	0.52
1:B:1262:GLN:HE21	1:B:1263:PHE:N	2.08	0.51
1:B:1114:GLU:H	1:B:1114:GLU:CD	2.13	0.51
1:A:1266:MET:HE3	4:A:1702:HOH:O	2.04	0.51
1:B:1180:ARG:NE	4:B:1609:HOH:O	1.94	0.51
1:B:1159:ARG:NH2	1:B:1303:GLU:OE1	2.38	0.51
1:A:1167:LYS:O	1:A:1171:GLU:HG3	2.11	0.51
1:B:1208:PHE:CE2	1:B:1262:GLN:HB3	2.46	0.51
1:A:1180:ARG:NE	4:A:1544:HOH:O	1.94	0.50
1:A:1143:ASN:ND2	4:A:1611:HOH:O	2.45	0.50
1:B:1148:ILE:HD11	1:B:1307:THR:HG22	1.93	0.50
1:B:1270:HIS:HD2	4:B:1510:HOH:O	1.95	0.49
1:B:1117:GLU:OE2	1:B:1152:LYS:NZ	2.40	0.49
1:B:1180:ARG:NH2	4:B:1609:HOH:O	2.43	0.49
1:B:1248:GLN:HA	1:B:1306:ILE:O	2.12	0.49
1:B:1156:LYS:HG3	4:B:1587:HOH:O	2.11	0.49
1:A:1159:ARG:HH12	1:A:1253:ARG:HE	1.61	0.48
1:A:1239:ASP:HB3	1:A:1242:CYS:HB2	1.94	0.48
1:B:1281:ARG:NH1	1:B:1281:ARG:HB3	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1114:GLU:CG	4:B:1595:HOH:O	2.60	0.48
1:A:1159:ARG:NH2	1:A:1303:GLU:OE2	2.47	0.47
1:B:1124:GLU:HG2	1:B:1251:PHE:CZ	2.49	0.47
1:B:1149:ARG:HB3	1:B:1307:THR:HB	1.97	0.47
1:B:1180:ARG:HG3	1:B:1180:ARG:HH11	1.80	0.47
1:A:1114:GLU:OE2	4:A:1537:HOH:O	2.20	0.47
1:A:1158:LEU:HD11	1:A:1196:GLY:CA	2.45	0.46
1:B:1134:ASP:C	1:B:1136:GLY:H	2.17	0.46
1:B:1189:ILE:HD12	1:B:1250:LEU:HG	1.96	0.46
1:B:1190:ASN:O	1:B:1194:HIS:CD2	2.68	0.46
1:A:1174:HIS:N	4:A:1689:HOH:O	2.27	0.46
1:B:1232:THR:N	4:B:1526:HOH:O	2.33	0.46
1:B:1117:GLU:OE1	1:B:1152:LYS:NZ	2.49	0.45
1:B:1253:ARG:O	1:B:1300:ALA:HA	2.17	0.45
1:B:1281:ARG:HH11	1:B:1281:ARG:HB3	1.82	0.45
1:B:1161:ARG:HD2	4:B:1606:HOH:O	2.17	0.45
1:B:1260:PHE:HE2	1:B:1262:GLN:HG2	1.80	0.45
1:A:1166:GLN:HG3	1:A:1177:HIS:HB2	1.98	0.44
1:B:1280:GLY:O	1:B:1290:ALA:HA	2.17	0.44
1:B:1180:ARG:HG2	4:B:1651:HOH:O	2.16	0.44
1:B:1190:ASN:O	1:B:1194:HIS:HD2	2.01	0.44
1:A:1188:PHE:O	1:A:1192:ILE:HG13	2.17	0.44
1:A:1114:GLU:HG3	4:A:1537:HOH:O	2.03	0.44
1:B:1117:GLU:CD	1:B:1152:LYS:NZ	2.71	0.44
1:A:1263:PHE:CD1	1:A:1283:SER:HB3	2.53	0.43
1:B:1220:LYS:NZ	1:B:1220:LYS:HB2	2.32	0.43
3:A:1402:2D6:C7	4:A:1608:HOH:O	2.65	0.43
1:A:1180:ARG:NH2	1:A:1216:GLU:OE1	2.45	0.43
1:A:1248:GLN:HA	1:A:1306:ILE:O	2.18	0.43
1:B:1215:ALA:HB1	4:B:1578:HOH:O	2.19	0.42
1:A:1174:HIS:CD2	4:A:1689:HOH:O	2.72	0.42
1:B:1262:GLN:NE2	1:B:1263:PHE:N	2.67	0.42
3:A:1402:2D6:H20	3:A:1402:2D6:H21	1.73	0.42
1:B:1268:MET:HE2	1:B:1295:TYR:OH	2.19	0.42
1:B:1155:ASN:OD1	1:B:1158:LEU:HB2	2.20	0.42
1:A:1148:ILE:H	1:A:1148:ILE:HG12	1.73	0.41
1:A:1144:ARG:HG2	1:A:1145:TYR:H	1.82	0.41
1:B:1295:TYR:N	1:B:1295:TYR:CD2	2.89	0.41
1:A:1298:GLU:HG2	4:A:1522:HOH:O	2.15	0.41
1:B:1132:HIS:C	1:B:1134:ASP:N	2.74	0.41
1:A:1114:GLU:OE2	4:A:1706:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/217 (96%)	204 (98%)	3 (1%)	1 (0%)	29	23
1	B	198/217 (91%)	191 (96%)	7 (4%)	0	100	100
All	All	406/434 (94%)	395 (97%)	10 (2%)	1 (0%)	47	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1133	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/185 (96%)	173 (98%)	4 (2%)	50	53
1	B	167/185 (90%)	157 (94%)	10 (6%)	19	14
All	All	344/370 (93%)	330 (96%)	14 (4%)	30	28

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1131	GLU
1	A	1158	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1170	SER
1	A	1314	GLU
1	B	1115	ASP
1	B	1134	ASP
1	B	1144	ARG
1	B	1158	LEU
1	B	1159	ARG
1	B	1180	ARG
1	B	1204	ILE
1	B	1220	LYS
1	B	1262	GLN
1	B	1298	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1143	ASN
1	A	1166	GLN
1	A	1173	ASN
1	A	1178	ASN
1	A	1201	HIS
1	B	1143	ASN
1	B	1174	HIS
1	B	1176	HIS
1	B	1178	ASN
1	B	1194	HIS
1	B	1201	HIS
1	B	1223	GLN
1	B	1262	GLN
1	B	1270	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	2D6	A	1402	-	34,35,35	2.60	17 (50%)	43,47,47	2.97	14 (32%)
3	2D6	B	1402	-	34,35,35	2.86	19 (55%)	43,47,47	3.03	18 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2D6	A	1402	-	-	3/16/26/26	0/4/4/4
3	2D6	B	1402	-	-	3/16/26/26	0/4/4/4

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1402	2D6	C14-C13	6.64	1.54	1.41
3	A	1402	2D6	C14-C13	5.74	1.52	1.41
3	B	1402	2D6	C16-C15	5.57	1.49	1.36
3	B	1402	2D6	C10-S1	5.25	1.97	1.81
3	A	1402	2D6	C16-C15	4.81	1.47	1.36
3	A	1402	2D6	C17-C18	4.70	1.47	1.36
3	B	1402	2D6	C6-C1	4.61	1.47	1.38
3	B	1402	2D6	C2-C1	4.53	1.47	1.38
3	A	1402	2D6	O1-C1	4.46	1.47	1.38
3	A	1402	2D6	C5-C4	4.23	1.48	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1402	2D6	C2-C1	3.96	1.46	1.38
3	B	1402	2D6	C17-C18	3.91	1.45	1.36
3	A	1402	2D6	C6-C1	3.84	1.46	1.38
3	B	1402	2D6	C5-C4	3.80	1.47	1.39
3	A	1402	2D6	C17-C16	3.71	1.47	1.38
3	B	1402	2D6	C17-C16	3.64	1.47	1.38
3	B	1402	2D6	C6-C5	3.34	1.44	1.38
3	B	1402	2D6	C23-C24	3.02	1.59	1.51
3	B	1402	2D6	C3-C4	3.01	1.45	1.39
3	B	1402	2D6	C15-C14	2.90	1.46	1.41
3	A	1402	2D6	C12-N2	2.89	1.38	1.33
3	A	1402	2D6	C10-S1	2.85	1.90	1.81
3	B	1402	2D6	C18-C13	2.82	1.46	1.41
3	B	1402	2D6	C12-N2	2.75	1.37	1.33
3	A	1402	2D6	C11-N2	2.73	1.38	1.34
3	A	1402	2D6	C18-C13	2.42	1.46	1.41
3	B	1402	2D6	C22-C21	2.41	1.57	1.52
3	B	1402	2D6	C11-S1	-2.34	1.71	1.75
3	A	1402	2D6	C3-C2	2.33	1.43	1.38
3	A	1402	2D6	C15-C14	2.28	1.45	1.41
3	A	1402	2D6	C6-C5	2.26	1.42	1.38
3	B	1402	2D6	C11-N3	-2.25	1.30	1.34
3	A	1402	2D6	C20-C21	2.21	1.57	1.52
3	B	1402	2D6	C11-N2	2.12	1.37	1.34
3	B	1402	2D6	O1-C1	2.06	1.42	1.38
3	A	1402	2D6	C8-N4	2.03	1.38	1.34

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1402	2D6	C14-C12-N2	-8.79	118.27	124.40
3	B	1402	2D6	C14-C12-N2	-8.66	118.36	124.40
3	B	1402	2D6	C12-N2-C11	8.45	125.47	115.79
3	A	1402	2D6	C3-C4-C7	-8.03	106.61	119.99
3	B	1402	2D6	C12-C14-C13	-7.58	111.99	118.59
3	A	1402	2D6	C12-N2-C11	7.41	124.28	115.79
3	A	1402	2D6	C5-C4-C7	5.55	129.23	119.99
3	B	1402	2D6	C15-C14-C12	5.49	129.67	121.61
3	B	1402	2D6	C21-N4-C8	-5.00	115.55	122.92
3	A	1402	2D6	C14-C13-N3	-4.57	121.12	123.60
3	A	1402	2D6	C21-N4-C8	-4.54	116.22	122.92
3	A	1402	2D6	C12-C14-C13	-4.43	114.74	118.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1402	2D6	N3-C11-N2	-3.69	120.19	126.95
3	B	1402	2D6	C5-C4-C3	3.58	125.42	118.96
3	A	1402	2D6	C19-C20-C21	3.46	115.31	111.48
3	A	1402	2D6	N3-C11-N2	-3.40	120.72	126.95
3	B	1402	2D6	C20-C21-N4	-3.34	103.66	110.56
3	B	1402	2D6	C14-C13-N3	-3.33	121.79	123.60
3	B	1402	2D6	C23-C22-C21	3.05	114.86	111.48
3	B	1402	2D6	C19-C20-C21	3.03	114.83	111.48
3	A	1402	2D6	C23-C22-C21	2.91	114.70	111.48
3	A	1402	2D6	C5-C4-C3	2.88	124.15	118.96
3	B	1402	2D6	C5-C6-C1	-2.73	116.39	119.73
3	B	1402	2D6	C2-C3-C4	-2.72	116.83	120.35
3	B	1402	2D6	C22-C23-C24	2.71	114.99	110.82
3	B	1402	2D6	C6-C1-C2	2.50	124.02	120.18
3	A	1402	2D6	C20-C19-C24	2.47	114.61	110.82
3	B	1402	2D6	C22-C21-N4	2.41	115.55	110.56
3	B	1402	2D6	C20-C19-C24	2.33	114.41	110.82
3	A	1402	2D6	C22-C23-C24	2.28	114.33	110.82
3	A	1402	2D6	C15-C14-C12	2.22	124.87	121.61
3	B	1402	2D6	O2-C8-N4	2.19	126.65	122.95

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1402	2D6	S1-C10-C9-C8
3	B	1402	2D6	O2-C8-C9-C10
3	B	1402	2D6	S1-C10-C9-C8
3	A	1402	2D6	N4-C8-C9-C10
3	B	1402	2D6	N4-C8-C9-C10
3	A	1402	2D6	O2-C8-C9-C10


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1402	2D6	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/217 (96%)	-0.23	1 (0%) 91 90	14, 21, 34, 38	0
1	B	202/217 (93%)	-0.09	4 (1%) 65 63	15, 24, 40, 48	0
All	All	412/434 (94%)	-0.16	5 (1%) 79 78	14, 23, 37, 48	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1281	ARG	3.9
1	B	1135	GLY	3.1
1	B	1133	ARG	3.0
1	A	1133	ARG	2.4
1	B	1207	MET	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	2D6	A	1402	32/32	0.87	0.18	16,23,33,35	0
3	2D6	B	1402	32/32	0.90	0.15	18,25,29,37	0
2	ZN	A	1401	1/1	0.99	0.07	22,22,22,22	0
2	ZN	B	1401	1/1	0.99	0.06	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.