



# wwPDB NMR Structure Validation Summary Report ⓘ

May 29, 2020 – 12:03 am BST

PDB ID : 2MV3  
Title : The N-domain of the AAA metalloproteinase Yme1 from *Saccharomyces cerevisiae*  
Authors : Scharfenberg, F.; Serek-Heuberger, J.; Martin, J.; Lupas, A.N.; Coles, M.  
Deposited on : 2014-09-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	88	

## 2 Ensemble composition and analysis ⓘ

This entry contains 23 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:103-A:165 (63)	0.07	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 4, 6, 7, 8, 9, 10, 11, 13, 14, 17, 18, 19, 22, 23
2	1, 3, 12, 16
3	5, 20
Single-model clusters	15; 21

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1375 atoms, of which 670 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Mitochondrial inner membrane i-AAA protease supercomplex subunit YME1.

Mol	Chain	Residues	Atoms						Trace
1	A	88	Total	C	H	N	O	S	0
			1375	436	670	130	134	5	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	MET	-	INITIATING METHIONINE	UNP P32795
A	177	GLU	-	EXPRESSION TAG	UNP P32795
A	178	HIS	-	EXPRESSION TAG	UNP P32795
A	179	HIS	-	EXPRESSION TAG	UNP P32795
A	180	HIS	-	EXPRESSION TAG	UNP P32795
A	181	HIS	-	EXPRESSION TAG	UNP P32795
A	182	HIS	-	EXPRESSION TAG	UNP P32795
A	183	HIS	-	EXPRESSION TAG	UNP P32795



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, NOESY back-calculation*.

Of the 100 calculated structures, 23 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.9.4
X-PLOR NIH	refinement	2.9.4
NMR-SPIRIT	refinement	1.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	501	492	492	1±1
All	All	11523	11316	11316	33

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 6 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:156:TYR:CE2	1:A:160:LEU:HD11	0.46	2.45	20	15
1:A:129:LEU:HD12	1:A:155:LEU:HD22	0.43	1.91	20	2
1:A:107:ARG:HD2	1:A:127:LEU:HD11	0.42	1.90	5	1
1:A:125:TYR:CB	1:A:155:LEU:HD13	0.42	2.44	23	13
1:A:156:TYR:CZ	1:A:160:LEU:HD11	0.42	2.50	23	1

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	63/88 (72%)	62±0 (98±0%)	1±0 (2±0%)	0±0 (0±0%)	100	100
All	All	1449/2024 (72%)	1426 (98%)	23 (2%)	0 (0%)	100	100

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	54/75 (72%)	54±0 (100±0%)	0±0 (0±0%)	100	100
All	All	1242/1725 (72%)	1242 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.6 Ligand geometry ⓘ

There are no ligands in this entry.

### 6.7 Other polymers ⓘ

There are no such molecules in this entry.



## 6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 72% for the entire structure.

### 7.1 Chemical shift list 1

File name: input\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	79	$-0.16 \pm 0.22$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	77	$0.42 \pm 0.18$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	52	$-0.12 \pm 0.28$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	52	$0.48 \pm 0.52$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 679 atoms were assigned a chemical shift out of a possible 784. 9 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	273/307 (89%)	112/122 (92%)	112/126 (89%)	49/59 (83%)
Sidechain	356/427 (83%)	222/253 (88%)	131/153 (86%)	3/21 (14%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	50/50 (100%)	26/26 (100%)	24/24 (100%)	0/0 (—%)
Overall	679/784 (87%)	360/401 (90%)	267/303 (88%)	52/80 (65%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	108	GLU	HA	2.17	6.30 – 2.20	-5.1

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

